

HISTIOSTOMA BLOMQUISTI N. SP. (ACARI: ASTIGMATA: HISTIOSTOMATIDAE), A
PHORETIC MITE OF THE RED IMPORTED FIRE ANT, *SOLENOPTIS INVICTA* BUREN
(HYMENOPTERA: FORMICIDAE) — ACAROLOGIA, 50(3): 357-371

(Erratum — *Acarologia*, 50(3) p. 366)

Stefan WIRTH¹ and John C. MOSER²

¹ Museum für Naturkunde, Leibniz-Institut für Evolutions- und Biodiversitätsforschung an der Humboldt-Universität zu Berlin, Invalidenstraße
43, 10115 Berlin. wirthstef@web.de (corresponding author)

² Southern Research Station, USDA Forest Service, 2500 Shreveport Hwy., Pineville LA, USA 7136. johnmoser@fs.fed.us

Please note the following text corrections for the legends of figures 6 and 7 (page 366):

Monthly Totals of Red Imported Fire Ant Alates and *Histiostoma blomquisti*




FIGURE 6: Frequency of *H. blomquisti* n. sp. deutonymphs phoretic on red imported fire ant alates from a 17-month study in 2004 and 2005.




FIGURE 7: Frequency of *H. blomquisti* n. sp. Deutonymphs per ant alate (*S. invicta*, n = 62 female alates) collected in Pineville/LA/USA from a 2 months study in March and April 2009.

COPYRIGHT

 Wirth and Moser. *Acarologia* is under free license.