Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Supporting agricultural research
for sustainable development

**Acarologia is under free license and distributed under the terms of the
Creative Commons-BY**

CHEYLOSTIGMAEUS HASSANPOURI N. SP., A NEW SPECIES OF THE GENUS
CHEYLOSTIGMAEUS WILLMANN, 1951 (ACARI: TROMBIDIFORMES:
STIGMAEIDAE) FROM NORTHWEST IRAN

Mohammad BAGHERI¹, Reza NAVAEE-BONAB⁴, Edward A. UECKERMANN^{2,3}, Hamed GHORBANI^{1,5}

(Received 16 May 2011; accepted 23 August 2011; published online 23 September 2011)

¹ Department of Plant Protection, Faculty of Agriculture, University of Maragheh, Maragheh, Iran. mbagheri20022002@yahoo.com; mobagheri2012@gmail.com; reza_kami2005@yahoo.com; hg.ghorbani@gmail.com

² ARC-PPRI, Private Bagx134, Queenswood, Pretoria 0121, South Africa UeckermannE@arc.agric.za

³ School of Environmental Sciences and Development, North-West University, Potchefstroom Campus 2520, South Africa

⁴ Young Researchers Club, Islamic Azad University, Marand Branch, Marand, Iran

⁵ Young Researchers Club, Islamic Azad University, Maragheh Branch, Maragheh, Iran

ABSTRACT — A new species of the genus *Cheylostigmaeus* (Acari: Stigmaeidae) is described and illustrated based on females and males collected from soil under apple trees in Marand, East Azerbaijan Province, Iran.

KEYWORDS — Acari; Prostigmata; Stigmaeidae; *Cheylostigmaeus*; Iran

INTRODUCTION

Members of the family Stigmaeidae are ecologically diverse, acting as predators of spider mites, scale insects, especially their eggs in leaf litter and on trees, and as herbivores on moss (Akyol and Koç 2010). This family is one of the most important predatory mite family after Phytoseiidae (Santos and Laing 1985, Croft 1994, Khanjani *et al.* 2010). To date 26 species of the genus *Cheylostigmaeus* have been described worldwide (Fan and Zhang 2004, Koç 2005, Khanjani *et al.* 2010, Dönel and Dogan 2011), among them two species namely *C. iranensis* Khanjani and Ueckermann 2002 and *C. ferdowsi* Khanjani *et al.* 2010 have been described from Iran. In this paper we described the third species, collected from soil

in apple orchards from East Azerbaijan province.

MATERIALS AND METHODS

Mites were extracted from the soil using a Berlese funnel; specimens were cleared in Nesbitt's fluid, mounted in Hoyer's medium (Walter and Krantz 2009) and were examined under 1000x magnification of Olympus Bx40 phase contrast microscope. The length of the idiosoma was measured from the base of chelicerae to the posterior margin of suranal shield, the width of the idiosoma at the broadest part of the idiosoma and setae were measured from their insertions to their tips; distances between setae were measured between their insertions. The terminology and abbreviations are based on Kethley

FIGURE 1: *Cheylostigmaeus hassanpouri* n. sp. (Male) – Dorsal view of body.

FIGURE 2: *Cheylostigmaeus hassanpouri* n. sp. (Male) – Ventral view of body.

(1990). All measurements are given in micrometers (μm). The first measurement is that of the holotype male or allotype female followed by the variation in measurements of the paratype specimens in parentheses including the holotype and allotype.

STIGMAEIDAE OUDEMANS, 1931

Cheyllostigmaeus Willmann, 1951

Cheyllostigmaeus hassanpourii n. sp. Bagheri

(Figures 1–11)

Diagnosis — Dorsal setae are thick and barbed sparsely in male and female; palpfemur without apophyses; ratios of setae *sce* : *sci* = 1.9, *ve* : *sci* = 2.6 in female; endopodal shield divided and smooth in both sex; femura I–IV and genua II–IV of male and female each with one sheathed seta; subcapitulum of male with one pair of bicuspidate lamellae; aedeagus without bulb, reversed comma-shape sclerotized appendages present on both sides of shaft distally.

Holotype male ($n = 4$) — Idiosoma oval, idiosoma length 292 (285 – 300); idiosoma width 214 (210 – 217); length of leg I 330 (325 – 331); leg II 280 (275 – 285); leg III 230 (265 – 270); leg IV 271 (267 – 274).

Dorsum — (Figure 1): Dorsum covered with three weakly sclerotized and smooth shields; no obvious dimples or micro-punctations observed; prodorsal shield truncated anteriorly, triangular and bearing four pairs of setae (*vi*, *ve*, *sci* and *sce*) and a pair of eyes; setae *ve* are the longest and setae *sci* the shortest; opisthosomal shield with six pairs of setae (*c1*, *d1*, *d2*, *e1*, *e2* and *f1*) and suranal shield with two pairs of setae (*h1* and *h2*); *h2* two-times longer than *h1*; most of dorsal setae fairly rigid, sparsely barbed and hyaline sheaths cover their tips; humeral shield bearing setae *c2* situated ventro-laterally. Length of setae : *vi* 28 (26 – 30); *ve* 67 (65 – 73); *sci* 19 (18 – 21); *sce* 49 (46 – 51); *c1* 34 (33 – 35); *c2* 52 (50 – 55); *d1* 34 (34 – 35); *d2* 40 (39 – 41); *e1* 33 (32 – 35); *e2* 42 (40 – 45); *f1* 65 (62 – 68); *h1* 23 (22 – 25); *h2* 41 (40 – 45); distances between dorsal setae: *vi-vi* 32 (30 – 35); *ve-ve* 72 (70 – 75); *vi-ve* 41 (39 – 43); *ve-sci* 22 (20 – 24); *sci-sci* 108 (106 – 110); *sci-sce*

22 (20 – 24); *sce-sce* 126 (125 – 130); *c1-d1* 55 (53 – 57); *c1-d2* 53 (50 – 55); *d1-d1* 93 (93 – 95); *d1-d2* 39 (37 – 42); *d1-e1* 50 (47 – 52); *d1-e2* 31 (30 – 33); *d2-e2* 52 (50 – 54); *e1-e1* 69 (67 – 71); *e1-e2* 35 (33 – 38); *e1-f1* 25 (23 – 27); *f1-f1* 63 (60 – 65); *h1-h1* 31 (30 – 33); *h1-h2* 11 (10 – 12); *h2-h2* 60 (57 – 63).

Venter — (Figure 2): Coxae I–II and III–IV surrounded by smooth and divided endopodal shields; length of setae *1a* 23 (21 – 25), *3a* 22 (20 – 25), and *4a* 20 (20 – 22); anogenital shield with three pairs of subequal aggenital setae (*ag1-ag3*) and three pairs of pseudanal (*ps1-ps3*) setae on tubercles; measurements of setae : *ag1* 16 (15 – 17); *ag2* 16 (15 – 18); *ag3* 20 (19 – 22); *ps1* 11 (11 – 12); *ps2* 12 (12 – 13); *ps3* 15 (14 – 15).

FIGURE 3: *Cheyllostigmaeus hassanpourii* n. sp. (Male) – Subcapitulum.

Gnathosoma — Chelicerae finely punctated, fixed digit 80 and moveable digit 60; palpi (Figure 5) five-segmented; palptarsus with five tactile setae plus one solenidiom (ω) and one trifold eupathidium distally; palptibia with two setae plus one claw and one spine-like accessory claw; palpgenu with two dorsal and palpfemur with three setae; palptrochanter without setae. Subcapitulum (Figure 3) finely punctated and with two pairs of subcapitular setae (*m* and *n*), *m* 19 (18 – 21), *n* 20 (18 – 24) and two pairs of adoral setae (*or1* and *or2*), *or1* 17

(17 – 19), *or*2 19 (17 – 20); subcapitulum bears a pair of strongly developed wing-like lamellae and one pair of acute teeth in front of lamellae, each lamellae thick and bicuspidate in front.

FIGURE 4: *Cheylostigmaeus hassanpourii* n. sp. (Male) – Aedeagus.

Aedeagus — (Figure 4) Calyx of the aedeagus funnel-shaped, shaft straight and pointed distally, no obvious bulb; reversed comma-shaped sclero-

tized appendages present on both sides of shaft distally.

FIGURE 5: *Cheylostigmaeus hassanpourii* n. sp. (Male) – Palp.

Legs — (Figures 6): Counts of setae (excluding solenidia ; solenidia in parentheses) on legs I–IV: coxae 2-2-2-2; trochanter 1-1-2-1; femora 6-5-3-2; genua 4-3+κ-1-1; tibiae 5(+1φ+1φp)-5(+1φp)-5(+1φp)-5(+1φp); tarsi 13(+2ω)-9(+2ω)-7(+2ω)-7(+2ω); dorsal setae are rigid and sparsely barbed; femora I–IV and genua II–IV each with one distinct sheathed seta; lengths of solenidia: Iω1 29 (28 – 30), Iω2 42 (40 – 44); IIω1 20 (20 – 21); IIω2 50 (48 – 52); IIIω1 9 (8 – 10); IIIω2 41 (40 – 44); IVω1 6 (5 – 8); IVω2 37 (35 – 40); Iφ 11 (10 – 12); φp 21 (20 – 22); IIφ 19 (19 – 21); IIIφ 21 (20 – 23); IVφ 19 (18 – 20).

Female (n = 5) — Idiosoma oval, idiosoma length: 350 (340 – 365); idiosoma width 280 (270 – 292); length of leg I 360 (350 – 370); leg II 300 (294 – 305); leg III 265 (260 – 270); leg IV 320 (312 – 325).

Dorsum — (Figure 7): Dorsum covered with two shields, the shields weakly sclerotized; no obvious dimples or micro-punctations; suranal shield is displaced to ventral position; prodorsal shield with four pairs (*vi*, *ve*, *sci* and *sce*) of setae and a pair of eyes, setae *ve* are the longest and setae *sci* the shortest; opisthosomal shield with six pairs of setae (*c1*, *d1*, *d2*, *e1*, *e2* and *f1*); suranal shield with two pairs of setae (*h1* and *h2*); dorsal setae fairly

FIGURE 6: *Cheylostigmaeus hassanpouri* n. sp. (Male): A – Leg I, femur-tarsus; B – Leg II, femur-tarsus; C – Leg III, femur-tarsus; D – Leg IV, femur-tarsus

rigid, sparsely barbed and hyaline sheaths cover their tips; humeral shields bearing setae *c2* situated ventro-laterally. Length of setae: *vi* 50 (48 – 50); *ve* 90 (85 – 90); *sci* 35 (33 – 36); *sce* 65 (62 – 67); *c1* 50 (50 – 51); *c2* 70 (70 – 75); *d1* 50 (49 – 50); *d2* 60 (54 – 60); *e1* 55 (55 – 60); *e2* 60 (57 – 63); *f1* 80 (72 – 81); *h1* 55 (50 – 55); *h2* 50 (48 – 50); distances between dorsal setae: *vi* – *vi* 40 (37 – 41); *ve* – *ve* 95 (92 – 98); *vi* – *ve* 65 (64–66); *ve* – *sci* 35 (33 – 35); *sci* – *sci* 150 (145 – 152); *sci* – *sce* 30 (27 – 32); *sce* – *sce* 180 (176 – 182); *c1* – *c1* 120 (118 – 125); *c1* – *d1* 76 (75 – 77); *c1* – *d2* 60 (59 – 63); *d1* – *d1* 115 (111 – 116); *d1* – *d2* 65 (62 – 67); *d1* – *e1* 80 (79 – 82); *d1* – *e2* 55 (53 – 55); *d2* – *e2* 80 (77 – 81); *e1* – *e1* 95 (92 – 96); *e1* – *e2* 55 (55 – 57); *e1* – *f1* 60 (60 – 63); *f1* – *f1* 75 (74 – 77); *h1* – *h1* 30 (30 – 34); *h1* – *h2* 20 (19 – 22); *h2* – *h2* 50 (49 – 53); setal ratios: *vi*: *vi* – *vi* 1.25; *c1*: *c1* – *c1* 0.42; *d1*: *d1* – *d1* 0.43; *e1*: *e1* – *e1* 0.58; *f1*: *f1* – *f1* 1.06; *h1*: *h1* – *h1* 1.83; *h2*: *h2* – *h2* 1; *c1* – *c1*: *d1* – *d1*: *e1* – *e1*: *f1* – *f1* 1.60: 1.53: 1.27: 1.00.

Venter — (Figure 8): Ventral cuticle striated; coxae I-II and III-IV surrounded by smooth and divided endopodal shields; length of setae *1a* 30 (28 – 30), *3a* 30 (28 – 31), and *4a* 32 (30 – 32); anogenital

shield with three pairs of subequal aggenital setae (*ag1*–*ag3*) and three pairs of pseudanal (*ps1*–*ps3*) setae; measurements of setae: *ag1* 15 (15 – 21); *ag2* 20 (18 – 21); *ag3* 20 (19 – 22); *ps1* 20 (19 – 20); *ps2* 20 (19 – 21); *ps3* 22 (19 – 22).

Gnathosoma — Chelicerae finely punctated, fixed digit 80 and moveable digit 60; palpi (Figure 9) five-segmented; palptarsus with five tactile setae plus one solenidiom (ω) and one trifid eupathidium distally; palptibia with two setae plus one claw and one spine-like accessory claw; palpgenu with two setae and palpfemur with three setae; palptrochanter without setae. Subcapitulum (Figure 10) finely punctated and with two pairs of subcapitular setae (*m* and *n*), *m* 35 (33 – 36), *n* 25 (25 – 27) and two pairs of adoral setae (*or1* and *or2*), *or1* 17 (15 – 17), *or2* 30 (27 – 31).

Legs — (Figures 11): Counts of setae (excluding solenidia; solenidia in parentheses) on legs I-IV: coxae 2-2-2-2; trochanters 1-1-2-1; femora 6-5-3-2; genua 4-3+ κ -1-1; tibiae 5(+1 ϕ +1 ϕp)-5(+1 ϕp)-5(+1 ϕp)-5(+1 ϕp); tarsi 13(+1 ω)-9(+1 ω)-7(+1 ω)-7(+1 ω); dorsal setae are rigid and sparsely barbed;

FIGURE 7: *Cheylostigmaeus hassanpouri* n. sp. (Female) – Dorsal view of body.

FIGURE 8: *Cheylostigmaeus hassanpouri* n. sp. (Female) – Ventral view of body.

femora I–IV and genua II–IV each with one distinct sheathed seta; lengths of solenidia: I ω 27 (25 – 27), II ω 19 (18 – 20); III ω 10 (10 – 11); IV ω 4 (4 – 5); I φ 12 (12 – 13); I φp 25 (23 – 25); II φp 20 (18 – 20); III φp 17 (16 – 17); IV φp 20 (18 – 21).

FIGURE 9: *Cheylostigmaeus hassanpouri* n. sp. (Female) – Palp.

Etymology — This species is named in honor of Dr. Mehdi Hassanpour, University of Mohaghegh Ardabili who is the friend of the first author.

Material examined — Holotype male, allotype female, paratype males (3) and paratype females (n= 4) were collected from soil in apple orchards, 20 August 2009, Marand, East Azerbaijan province, Iran, by Reza Navaei-Bonab. The holotype and one paratype female are deposited in the Arachnida Collection of Plant Protection Research Institute, Pretoria, South Africa and the rest of the paratypes are deposited in the Acarological Collection, Department of Plant Protection, Faculty of Agriculture, University of Maragheh, Maragheh, Iran.

FIGURE 10: *Cheylostigmaeus hassanpouri* n. sp. (Female) – Subcapitulum.

Remarks — The new species is unique in having the following characters:

1. setae *sci* are the smallest dorsal setae and 1/3 length of *ve*, the longest setae,
2. shape of the male aedeagus.

The new species closely resembles *Cheylostigmaeus luxtoni* Wood (Wood, 1968) by having one pair of bicuspidate lamellae on subcapitulum and absence of bulb on aedeagus but differs from the latter by:

1. the length of setae *sci* 18 – 21 in new species vs 34 – 49 in *C. luxtoni*,
2. the distance *vi-vi* in the new species (30 – 35) is about the half (59 – 69) than in *C. luxtoni*,
3. shape of copulatory appendages of aedeagus in the two species distinctively differ.

Cheylostigmaeus hassanpouri n. sp. is also close to *C. pannonicus* Willmann (Summers and Ehara, 1965) by the following characters: femur of palpus without apophyse and having one pair of bicuspidate lamellae on subcapitulum. But it differs from this latter species by:

FIGURE 11: *Cheylostigmaeus hassanpouri* n. sp. (Female): A – Leg I, femur-tarsus; B – Leg II, femur-tarsus; C – Leg III, femur-tarsus; D – Leg IV, femur-tarsus.

1. aedeagus of *C. pannonicus* has a capacious bulb which is absent in the new species, additionally the copulatory appendages have different shapes in the two species,
2. ratio $ve : sci = 2.6$ in female of the new species vs 1.3 in those of *C. pannonicus*.

The new species also resembles to *Cheylostigmaeus urhani* Dönel, G. and Dogan as in both species the setae *sci* are the smallest prodorsal setae and there is a bicuspidate lamellae on subcapitulum.

However these species could be differentiated by the following characters:

1. in all specimens of the new species apophyses on the palp femur are absent whereas *C. urhani* has apophyses on the palp femur,
2. ratio of $sce : sci = 1.9$ [sce 65 (62 – 67), sci 35 (33 – 36)] in female of the new species vs 1.4 [sce 110 (105 – 125), sci 77 (65 – 97)] in *C. urhani*,

3. there is not obvious bulb on aedeagus of the new species vs capacious bulb in *C. urhani*,
4. endopodal shield of the new species is smooth vs reticulated in *C. urhani*.

ACKNOWLEDGEMENTS

This project was supported by the research division of the University of Maragheh, Maragheh, Iran, which is greatly appreciated. The authors also wish to thank the reviewers of this paper for their inputs. Thanks also to Dr. Salih Dogan, Atatürk University, Kazim Karabekir Education Faculty, Department of Biology Education, Erzurum, Turkey for some literatures.

REFERENCES

- Akyol, M. and Koç, K. 2010 — Contributions to the raphignathoid fauna of Turkey, with a description of a new species of *Cryptognathus* Kramer (Acari: Actinedida: Raphignathoidea) — Turk. J. Zool., 34: 159-167.

- Croft, B.A. 1994 — Biological control of apple mites by a phytoseiid mite complex and *Zetzellia mali* (Acari: Stigmaeidae): long-term effects and impact of azinphosmethyle on colonization by *Amblyseius andersoni* (Acari: Phytoseiidae) — Environ. Entomol., 23: 1317-1325.
- Dönel, G. and Dogan, S. 2011 — The stigmaeid mites (Acari: Stigmaeidae) of Kelkit Valley (Turkey) — Zootaxa, 2942: 1-56.
- Fan, Q. H. and Zhang, Z. Q. 2004 — Revision of raphignathoid mites (Acari: Prostigmata) in the collection of Habeeb — Zootaxa, 736: 1- 28.
- Kethley, J. 1990 — Acarina: Prostigmata (Actinedida). In: Dindal, D.L. (Ed.), Soil Biology Guide - John Wiley and Sons, New York, pp. 667-756.
- Khanjani, M., Raisii, H., Izadi, H. and Ueckermann, E. A. 2010 — A new species of the genus *Cheylostigmaeus* Willmann (Acari: Stigmaeidae) from eastern Iran — Internat. J. Acarol., 36: 7-13. doi:10.1080/01647950903473323
- Khanjani, M. and Ueckermann, E. A. 2002 — The stigmaeid mites of Iran (Acari: Stigmaeidae) — Internat. J. Acarol., 28: 317-339. doi:10.1080/01647950208684309
- Koç, K. 2005 — A new species of the genus *Cheylostigmaeus* (Acari: Stigmaeidae) from Turkey — Biologia, 60: 483- 487.
- Summers, F.M. and Ehara, S. 1965 — Revaluation of the taxonomic characters in four species of the genus *Cheylostigmaeus* Willmann (Acarina: Stigmaeidae) — Acarologia, 7: 49-62.
- Santos, M. A. and Laing, J. E. 1985 — Other predaceous mites and spiders 2.2.1 Stigmaeid predators. pp. 197-203. Helle, W. and Sabelis, M. W. (Eds.). Spider Mites, Their Biology, Natural Enemies and Control. World Crop Pests, vol. LB, 458 pp. Elsevier, Amsterdam, The Netherlands.
- Walter, D. E. and Krantz, G. W. 2009 — Collection, rearing and preparing specimens: 83-96. In: (Eds.) Krantz, G.W. and Walter, D.W. A Manual of Acarology. Third Edition. 807 pp.
- Wood, T.G. 1968 — A new species of *Cheylostigmaeus* Willmann (Acari: Stigmaeidae) from New Zealand — N. Z. Jl Sci, 11: 276-279.

COPYRIGHT

Bagheri *et al.*. Acarologia is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.