

A new species of *Myrmozercon* Berlese (Acari: Mesostigmata: Laelapidae) associated with an arboreal ant (Formicidae: *Crematogaster*) from Iran

Omid JOHARCHI^{1✉}, Elham ARJOMANDI² and Viacheslav A. TRACH³

(Received 25 November 2016; accepted 03 February 2017; published online 06 July 2017; edited by Farid FARAJI)

¹ Department of Plant Protection, Yazd Branch, Islamic Azad University, Yazd, Iran. (✉) joharchi@iauyazd.ac.ir, j.omid2000@gmail.com

² Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, 14115-336, Tehran, Iran. elham.arjomandi@yahoo.com

³ Department of Zoology, I. I. Mechnikov Odessa National University, Shampanskij Al., 2, 65058, Odessa, Ukraine. vatrach@gmail.com

ABSTRACT — This paper reports on a new species of mite of the genus *Myrmozercon* Berlese, 1902 associated with arboreal ants in Iran – *Myrmozercon brachytrichos* **n. sp.** was collected in association with *Crematogaster inermis* Mayr (Formicidae) in Alborz province, Iran. This new species is described and illustrations provided.

KEYWORDS — Laelapidae; *Myrmozercon*; arboreal ant; taxonomy; Iran; myrmecophiles

ZOOBANK — [F51400D4-BA00-47F7-B823-37D0463A30BB](https://zoobank.org/F51400D4-BA00-47F7-B823-37D0463A30BB)

INTRODUCTION

Many species Laelapidae have been reported from ants or their nests, but the genus *Myrmozercon* Berlese, has a more intimate association with ants. Currently, *Myrmozercon* comprises 29 described species from Europe, Transcaucasia, Middle East, Central Asia, Africa, North America and Australia (Karawajew 1909; Vitzthum 1930; Hunter and Hunter 1963; Rosario and Hunter 1988; Walter 2003; Shaw and Seeman 2009; Trach and Khaustov 2011; Joharchi *et al.* 2011; Ghafarian *et al.* 2013; Joharchi and Moradi 2013; Babaeian *et al.* 2013, 2014; Joharchi *et al.* 2015). The most recent taxonomic work on the genus (Joharchi *et al.* 2015) revised the generic concept and its morphological attributes, provided a key to the well-described species of *Myrmozercon* and reviewed the poorly described species. They suggested that the most highly mod-

ified species of *Myrmozercon* from a monophyletic group, termed *Myrmozercon sensu stricto*, because it includes the type species *Myrmozercon brevipes* Berlese, 1902 (redescribed by Kontschán & Seeman 2015). The members of *Myrmozercon sensu stricto* not only share a multitude of reductive or specialized character states, but also tend to associate with ants in the genus *Crematogaster* (Joharchi *et al.* 2015).

MATERIALS AND METHODS

Laelapidae associated with ants were collected in different regions of Iran over a period of eight years (2008 – 2016). Mites were removed from ants' nests by individual hand picking and by extraction from ant nesting material using Tullgren funnels. Mites were cleared in Nesbitt's solution and mounted in Hoyer's medium (Walter and Krantz 2009). The line drawings and examinations of the specimens were

performed with a compound microscope (XS-2100 series) equipped with a drawing tube. Pencil line drawings were then scanned and traced over using Microsoft Office © Powerpoint 2003. Measurements of structures are expressed as minimum-maximum ranges in micrometres (µm). Dorsal shield length and width were taken from the anterior to posterior margins along the midline, and at its broadest point, respectively. Length and width of the sternal shield were measured from the anterior point to the posterior point at the full length and broadest point, respectively. Genital shield length and width were measured along the midline from the posterior margin of the sternal shield to the posterior margin of the genital shield, and at the level of setae *st5*, respectively. Leg lengths were measured from base of the coxa to the apex of the tarsus, excluding the pre-tarsus. Lengths for the fixed and movable cheliceral digits were taken from the base of the movable digit to their tips. The nomenclature used for the dorsal idiosomal chaetotaxy is that of Lindquist and Evans (1965), the leg chaetotaxy is that of Evans (1963), and names of other anatomical structures mostly follow Evans and Till (1979). We use the term "lyrifissure" to refer to slit-shaped sensilli, and "pore" for circular or oval-shaped cuticular openings of unspecified function. The holotype and one paratype of the new species are deposited in the Acarological collection, Department of Plant Protection, Yazd Branch, Islamic Azad University (YIAU). Paratypes are also deposited in the Jalal Afshar Zoological Museum, College of Agriculture, University of Tehran, Iran (JAZM) and in Museum für Naturkunde (MfN), Berlin, Germany.

RESULTS

Genus *Myrmozercon* Berlese, 1902

Myrmozercon Berlese, 1902: 699. Type species *Myrmozercon brevipes* Berlese, 1902, by monotypy. *Myrmonyssus* Berlese, 1903: 16. Type species *Myrmonyssus diplogeni* Berlese, 1903, designated by Berlese, 1904 (synonymy by Rosario and Hunter 1988). *Myrmonyssus* (*Laelaspulus*) Berlese, 1904: 437. Type species *Myrmozercon acuminatus* Berlese, 1903, by original designation (synonymy by Shaw and Seeman 2009).

Parabisternalis Ueckermann and Loots, 1995: 35. Type species *Parabisternalis yemeni* Ueckermann and Loots, 1995, by original designation (synonymy by Shaw and Seeman 2009).

The diagnosis of the genus *Myrmozercon* used here is based on Joharchi *et al.* (2015) and Shaw and Seeman (2009).

Myrmozercon brachytrichos n. sp. (Figures 1–2)

Myrmozercon tauricus Trach and Khaustov, 2011: 23 – Joharchi and Moradi, 2013: 250, (misidentification).

Zoobank: [FB546BDF-8B25-4594-AD15-8CEF1D1CE9A4](https://zoobank.org/FB546BDF-8B25-4594-AD15-8CEF1D1CE9A4)

Type material — Holotype, female, Iran, Alborz province, Savojbolagh, Khoznan, 36°71' N, 50°32' E, alt. 1595 m, 15 August 2016, O. Joharchi coll., clinging to the head of *Crematogaster inermis* Mayr (Formicidae) (misidentification in Joharchi and Moradi, 2013) in the bark of grapevine (in YIAU). Paratypes: one female same data as holotype (in JAZM), two females, Iran, Damavand mountain, 35°52' N, 52°07' E, alt. 2422 m, 18 May 2013, O. Joharchi coll., clinging to the head of *Crematogaster inermis* (one in YIAU and one in MfN) (host ants are also in YIAU).

Description of female (n = 4) — Dorsal idiosoma (Figure 1A). Length 470 – 475. Dorsal shield length 440 – 449, width 316 – 325. Shield posteriorly truncate, not covering entire idiosoma, leaving a strip of striate cuticle posterior to setae *J5*, shield without distinct reticulate ornamentation over whole surface; with 34 pairs of setae, 20 podonotal (*z2* and *z6* absent), 14 opisthonotal, *Z5* in soft skin posterior to shield, almost all setae slightly barbed in apical third or less, all setae on shield minute, uniform in length 12 – 15 and thickness, without unpaired or asymmetrical setae. Setae in *R*-series absent. Shield with 12 pairs of pore-like structures; lyrifissures near base of *j1* large and slit-like.

Ventral idiosoma (Figure 1B) — Tritosternum with short broad base (25 – 30 × 14 – 16 wide), laciniae 46 – 48 forked a short distance above suture, with smooth edges, strap-like and broad at base; pre-sternal shields apparently fused with sternal shield. Sternal shield (length 200 – 210) narrowest between coxae II (106 – 108) widest between




FIGURE 1: *Myrmozercon brachytrichos* n. sp., female: A – dorsal idiosoma; B – ventral idiosoma; C – epistome; D – hypostome; E – chelicerae.

coxae II and III (166 – 172), with biconvex anterior margin and extending beyond level of *st1*, lateral margins thickened and posterolateral corners fused with endopodal shields; posterior margin concave; shield not eroded between *st1* – *st1*, bearing three pairs of smooth pointed setae (*st1* 14 – 16, *st2* 20 – 22, *st3* 20 – 22) and two pairs of lyrifissures, lyrifissures *iv1* adjacent to setae *st1*, lyrifissures *iv2* between *st2* and *st3*; weak lines near central and posterior margin, anterior area smooth. Seta *st4* absent, metasternal pores also on extent of sternal shield but metasternal plates apparently absent. Genito-ventral shield wide, inverted U-shape, strongly tapering posteriorly, 240 – 250 long, 106 – 115 maximum width. Surface of shield has characteristic ornamentation consisting of longitudinal striae; with one pair of simple setae *st5* (20 – 22). Anal shield (52 – 60 × 100 – 107 wide) with anterolateral extensions, its anterior without lineate ornamentation, cribrum a narrow band without discernible rows, with a pair of circular lateral gland pores, bearing post-anal seta 8 – 10 long and a pair of para-anal setae 8 – 10 long. Opisthogastric skin with long, narrow and oval metapodal plates (34 – 40 × 8 – 10 wide) and 12 pairs of simple setae 10 – 14 long. Peritreme short (35 – 40), extending from coxa IV to posterior level of coxa II. Peritrematal shields absent, post-stigmatal section conspicuous and narrow, bearing lyrifissure near stigmata and two pore-like structures of post-stigmatal pores.

Gnathosoma — Epistome triangular, irregularly denticulate (Figure 1C). Hypostomal groove with seven rows of denticles, three to six very fine denticles per row (Figure 1D). Hypostome with four pairs of setae, internal posterior hypostomal setae *h3* longest. Palp chaetotaxy: trochanter 1, femur 5, genu 6, tibia 11; all palp setae pointed, palp tarsal claw two-tined, dorsodistal edge of palp femur without swelling. Chelicera hyaline, fixed digit of chelicera reduced and edentate, with a low median bulge, pilus dentilis and dorsal lyrifissure absent (Figure 1E); movable digit weakly sclerotised, with one strong terminal hook, cheliceral seta, arthrodial corona and its filaments absent (Figure 1E). Corniculi long, membranous. Lateral malae arms absent (Figure 1D).

Legs (Figure 2 A–D) — Length of leg I 200 – 210, leg II 190 – 202, leg III 178 – 190, leg IV 190 – 205. Setal counts for legs I–IV: 2–4–8–8–8, 2–4–8–8–7, 2–4–6–8–7, 0–5–5–8–7. Chaetotaxy (both sides examined): Leg I: coxa 0 0/1 0/1 0, trochanter 1 0/1 0/1 1, femur 1 2/1 2/1 1, genu 1 2/1 2/1 1, tibia 1 2/1 2/1 1 (Figure 2A). Leg II: coxa 0 0/1 0/1 0, trochanter 1 0/1 0/1 1, femur 1 2/1 2/1 1, genu 1 2/1 2/1 1, tibia 1 1/1 2/1 1 (Figure 2B). Leg III: coxa 0 0/1 0/1 0, trochanter 1 0/1 0/1 1, femur 1 2/1 1/0 1, genu 1 2/1 2/1 1, tibia 1 1/1 2/1 1 (Figure 2C). Leg IV: coxa 0 0/0 0/0 0, trochanter 1 1/2 0/1 0, femur 1 2/1 1/0 0, genu 1 2/1 2/1 1, tibia 1 1/1 2/1 1 (Figure 2D); all setae smooth. Tarsi II–IV with 18 setae 3 3/2 3/2 3 + mv, md. All pre-tarsi with a pair of thin claws and a membranous ambulatorium.

Genital structures — Insemination ducts opening on posterior margin of coxa III, sacculus indistinct, apparently unsclerotised.

Male. Unknown.

Etymology — The name *brachytrichos* (Greek brachy, short and trichos, hair) refers to the minute dorsal shield setae.

Remarks — According to the key to species provided by Joharchi *et al.* (2015), *Myrmozercon brachytrichos* n. sp. most resembles *M. tauricus* Trach and Khaustov, 2011. The new species differs from *M. tauricus* by having: opisthonotal shield with 14 pairs of setae and Z5 in soft skin posterior to shield (versus Z5 in opisthonotal shield in *M. tauricus*), all setae on shield minute and uniform in length (versus posterior dorsal setae longer than anterior dorsal setae in *M. tauricus*); epistome triangular, irregularly denticulate (versus smooth in *M. tauricus*), sternal shield not eroded between *st1*–*st1* (versus eroded between *st1*–*st1* in *M. tauricus*), setal counts for trochanter I–III (four setae in *M. brachytrichos* versus five setae in *M. tauricus*) and for genu IV (eight setae in *M. brachytrichos* versus seven setae in *M. tauricus*). *Myrmozercon brachytrichos* is also very similar to three other species: *Myrmozercon eidmanni* (Sellnick), *Myrmozercon liguricus* (Vitzthum) and *Myrmozercon minor* (Sellnick). The new species differs from *M. eidmanni* by having no seta on coxa IV and all dorsal shield setae minute and uniform in length (*vs* coxa IV seta present and seta *j1* longer




FIGURE 2: *Myrmozercon brachytrichos* n. sp., female: A – trochanter, femur, genu and tibia I, dorsal aspect; B – trochanter, femur, genu and tibia II, dorsal aspect; C – trochanter, femur, genu and tibia III; D – trochanter, femur, genu and tibia IV, dorsal aspect.

than other body setae in *M. eidmanni*); it differs from *M. liguricus* by the slightly reduced fixed cheliceral digit (obsolete fixed digit in *M. brachytrichos*); and it differs from *Myrmozercon minor* by having fewer setae on the dorsal shield (34 pairs in *M. brachyserose*; approximately 31 pairs in *M. minor*).

Discussion — *Myrmozercon brachytrichos* n. sp. belongs to the *Myrmozercon sensu stricto* species group, which has the following character states: leg I femur, genu and tibia 8–8–8; leg II 8–8–7; leg III genu and tibia 8–7; leg IV genu with seven or eight setae, tibia with seven setae; legs short, palp trochanter with one seta; palpcoxal seta sometimes absent; corniculi membranous, dorsal shield smooth, sternal shield an inverted U-shape, setae *st4* usually absent and anal shield usually with anterolateral extensions.

Myrmozercon s. str. may have a more intimate association with their hosts because they are almost always collected while they cling onto ants (see Joharchi and Moradi 2013). Almost all have been reported in association with arboreal ants of the genus *Crematogaster* but nothing is known of their feeding behavior, or any other aspect of their biology. However, their specialized morphology (e.g. edentate chelicerae, short peritremes, corniculi membranous etc.) suggest that *Myrmozercon s. str.* could be parasitic on its ant hosts, and not simply a commensal in its host's nests, but this has not been established experimentally.


ACKNOWLEDGEMENTS

We gratefully acknowledge Dr. Owen D. Seeman (Queensland Museum, South Brisbane, Queensland, Australia) for reviewing and constructive comments on an earlier version of the manuscript. The authors wish to thank Dr. Farid Faraji (Mitox, Amsterdam, Netherlands) and the other anonymous reviewer for their valuable suggestions.

REFERENCES

- Babaeian E., Joharchi O., Saboori A. 2013 — A new species of *Myrmozercon* Berlese (Acari: Mesostigmata: Laelapidae) associated with ant from Iran — *Acarologia*, 53: 453-460. doi:10.1051/acarologia/20132109
- Babaeian E., Seraj A., Nemati A. 2014 — Description of a new ant-associated species (Acari: Mesostigmata: Laelapidae) from Iran — *Acarologia*, 54: 441-448. doi:10.1051/acarologia/20142129
- Berlese A. 1902 — Specie di Acari nuovi — *Zool. Anz.*, 25: 697-700.
- Berlese A. 1903 — Diagnosi di alcuninuove specie di Acari italiani, mirmecofili e liberi — *Zool. Anz.*, 27: 12-28.
- Berlese A. 1904 — Illustrazione iconografica degli Acari mirmecofili — *Redia*, 1: 299-474 + Plates 7-20.
- Evans G.O. 1963 — Observations on the chaetotaxy of the legs in the free-living Gamasina (Acari: Mesostigmata) — *Bull. Br. Mus. (Nat. Hist.) Zool.*, 10: 277-303.
- Evans G.O., Till W.M. 1979 — Mesostigmatic mites of Britain and Ireland (Chelicerata: Acari: Parasitiformes). An introduction to their external morphology and classification — *Trans. Zool. Soc. Lond.*, 35: 145-270. doi:10.1111/j.1096-3642.1979.tb00059.x
- Ghafarian A., Joharchi O., Jalalizand A., Jalaieian M. 2013 — A new species of *Myrmozercon* Berlese (Acari: Mesostigmata: Laelapidae) associated with ants from Iran — *ZooKeys*, 272: 21-28.
- Hunter P.E., Hunter C.A. 1963 — The genus *Myrmonyssus* with descriptions of two new species (Acarina: Laelapidae) — *Acarologia*, 5: 335-341.
- Joharchi O., Babaeian E., Seeman O.D. 2015 — Review of the genus *Myrmozercon* Berlese (Acari: Laelapidae), with description of a new species from Iran — *Zootaxa*, 3955(4): 549-560.
- Joharchi O., Halliday B., Saboori A., Kamali K. 2011 — New species and new records of mites of the family Laelapidae (Acari: Mesostigmata) associated with ants in Iran — *Zootaxa*, 2972: 22-36.
- Joharchi O., Moradi M. 2013 — Review of the genus *Myrmozercon* Berlese (Acari: Laelapidae), with description of two new species from Iran — *Zootaxa*, 3686: 244-254.
- Karawajew W. 1909 — Myrmekophilen aus Transkaspien — *Russkoe Entomologicheskoe Obozrenie*, 9: 227-237.
- Lindquist E.E., Evans G.O. 1965 — Taxonomic concepts in the Ascidae, with a modified setal nomenclature for the idiosoma of the Gamasina (Acarina: Mesostigmata) — *Mem. Ent. Soc. Can.*, 47: 1-64. doi:10.4039/entm9747fv
- Rosario R.M.T., Hunter P.E. 1988 — The genus *Myrmozercon* Berlese, with descriptions of two new species (Acari: Mesostigmata: Laelapidae) — *J. Parasitol.*, 74: 466-470. doi:10.2307/3282057
- Sellnick M. 1941 — Milben von Fernando Poo — *Zool. Anz.*, 136: 221-228.
- Shaw M.D., Seeman O.D. 2009 — Two new species of *Myrmozercon* (Acari: Laelapidae) from Australian ants (Hymenoptera: Formicidae) — *Zootaxa*, 2025: 43-55.
- Trach V.A., Khaustov A.A. 2011 — A myrmecophilous mite *Myrmozercon tauricus* sp.n. of the family Laelapidae (Acari, Mesostigmata) from Ukraine — *Vestn. Zool.*, 45: 23-27.
- Ueckermann E.A., Loots G.C. 1995 — A new laelapid genus and species (Acari: Parasitiformes: Laelapidae) from Yemen — *Afr. Entomol.*, 3: 35-38.
- Vitzthum H.G. 1930 — Ein Ameisengast (Acar.) — *Mitt. Deut. Entomol. Gesell. Ber.*, 6: 89-94.
- Walter D.E. 2003 — A new mite from an arboreal ant (Formicidae: *Polyrachis* sp.): *Myrmozercon iainkayi* n. sp. (Mesostigmata: Laelapidae) — *Int. J. Acarol.*, 29: 81-85. doi:10.1080/01647950308684325
- Walter D.E., Krantz G.W. 2009 — Collecting, rearing and preparing specimens. In: Krantz G.W., Walter D.E. (eds) *A Manual of Acarology*, Third Edition, Texas Tech University Press, Lubbock, Texas, pp. 83-95.

COPYRIGHT

 Joharchi O. *et al.* *Acarologia* is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.