Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Supporting agricultural research
for sustainable development

Acarologia is under **free license** and distributed under the terms of the
Creative Commons-BY

Psittophagus hollandicus n. sp., a new feather mite species (Acariformes: Pterolichidae) from the cockatiel *Nymphicus hollandicus* (Kerr, 1792) (Psittaciformes: Cacatuidae) in Brazil

Fabio Akashi HERNANDES

(Received 03 March 2017; accepted 09 May 2017; published online 29 September 2017; edited by Michel BERTRAND)

Departamento de Zoologia, Av. 24-A, 1515, 13506-900, Universidade Estadual Paulista, Rio Claro, São Paulo State, Brazil. abakashi@gmail.com;
Phone (55) 19-3526-4296.

ABSTRACT — A new feather mite species, *Psittophagus hollandicus* n. sp. (Acariformes: Pterolichidae), is described from the cockatiel *Nymphicus hollandicus* (Kerr, 1792) (Psittaciformes: Cacatuidae) kept in captivity in Brazil.

KEYWORDS — Acari; Psoroptidia; Pterolichoidea; systematics; household pets

ZOOBANK — [D14DD129-3984-4DBE-BC0D-C971F50538B3](https://zoobank.org/D14DD129-3984-4DBE-BC0D-C971F50538B3)

INTRODUCTION

The cockatiel *Nymphicus hollandicus* (Kerr, 1792) (Psittaciformes: Cacatuidae) is native to Australia but owing to maintaining as a household pet it distributed worldwide (Engelbreton, 2006). The only other feather mite species previously described from this host is *Nymphicilichus perezae* Mironov & Galloway, 2002 (Astigmata: Pterolichidae) in New Zealand.

Pterolichid feather mites associated with parrots (Aves: Psittaciformes) are represented by three morphologically distinct generic groups: *Psittophagus* (4 genera), *Rhytidelasma* (12 genera), and *Protolichus* (24 genera) (Gaud and Mouchet, 1959; Gaud, 1980; Atyeo and Pérez, 1982, 1990; Atyeo *et al.*, 1984; Atyeo, 1985, 1988, 1989a, b, c; Atyeo and Gaud, 1987; Gaud and Atyeo, 1996; Mironov and Galloway, 2002, Mironov and Pérez, 2003; Mironov *et al.*, 2003a,b, 2005, 2011, 2014; Dabert *et al.*, 2004, 2006, 2008; Mironov and Dabert, 2007, 2010). From these, the *Psittophagus* generic group is the least diverse, comprising currently 9 species (including the new one described herein) in four genera: *Nymphicilichus* Mironov & Galloway, 2002, *Micropsittophagus* Mironov & Pérez, 2003, *Psittaculobius* Mironov, Dabert & Ehrnsberger, 2003, and *Psittophagus* Gaud & Atyeo, 1996; the three former genera include only one species each.

To date, the genus *Psittophagus* has included five species associated with Old World parrots of the family Cacatuidae (Psittaciformes) from the Australasian region: *Psittophagus ornatus* (Mégnin & Trouessart, 1884) (type species), *P. obsoletus* (Mégnin & Trouessart, 1884), *P. galahi* Mironov, Dabert & Proctor, 2003, *P. calyptorhynchi* Mironov, Dabert & Ehrnsberger, 2003, and *P. lacunosus* Dabert, Badek & Skoracki 2007 (Mégnin and Trouessart, 1884;

Mironov *et al.*, 2003a, b; Dabert *et al.*, 2007). A table summarizing the mite-host relationships of the *Psittophagus* generic group, in addition to a key to genera and species of this group, were provided by Mironov *et al.* (2003a). In this paper, a new species of *Psittophagus* is described from the cockatiel in Brazil.

MATERIALS AND METHODS

The material was collected from a specimen of *Nymphicus hollandicus* which died in captivity and sent to the laboratory of Acari of the Department of Zoology, Universidade Estadual Paulista, Rio Claro, São Paulo, Brazil. The bird specimen was placed in a freezer and then washed with water containing detergent; the liquid was then filtered and mites were collected with a fine brush from the filter papers under stereomicroscope, cleared in 30% lactic acid for 24 h at 50°C, and mounted in Hoyer's medium (Krantz and Walter, 2009). After five days at 50 °C, the slides were sealed with varnish. Drawings and measuring of mites were made with a Leica DM3000 microscope equipped with differential interference contrast (DIC) optics and a camera lucida. Pencil sketches were scanned at 300 dpi and grayscale mode; line drawings were created with Adobe Illustrator CS6 and a Wacom Bamboo Create tablet. The idiosomal and leg chaetotaxies follow Griffiths *et al.* (1990) and Atyeo and Gaud (1966), respectively, with corrections for coxal setae proposed by Norton (1998). Type specimens are deposited at DZUnesp-RC — Collection of Acari of Department of Zoology of the Universidade Estadual Paulista, Rio Claro, São Paulo, Brazil.

Family Pterolichidae Trouessart & Mégnin, 1884
Subfamily Pterolichinae Trouessart & Mégnin,
1884

Genus *Psittophagus* Gaud & Atyeo, 1996

***Psittophagus hollandicus* n. sp.**

Zoobank: [F5817710-74E3-4711-ABEF-F4A935AC21F7](https://doi.org/10.5281/zenodo.1000000)

Type material: holotype ♂ (DZUnesp-RC#3759), paratypes 10 ♂ and 4 ♀ (#3760–3773) ex *Nymphicus hollandicus* (Kerr, 1792) (Psittaciformes: Cacatuidae),

captivity specimen from Pedreira, São Paulo State, BRAZIL, 17 August 2007, col. David V. Boas Filho (#203).

Differential diagnosis — The new species is morphologically close to *P. lacunosus* Dabert, Badek & Skoracki, 2007 in having, in males, setae *e2* thick spiculiform, and the hysterosomal shield reaching to or close to the level of setae *c1*. *Psittophagus hollandicus* n. sp. is distinguished from that species in having, in males, the dorsal shields with weak ornamentation, the prodorsal shield split transversally at the level of scapular setae and free from the scapular shields, solenidion σ of genu III about half as long as the segment (Fig. 3C), setae *si*, *c2* filiform, and the anal suckers without indentations. Females of the new species have piliform setae *d2* and *e2*, setae *d2* being 30µm long and barely reaching the halfway to bases of piliform setae *e2*. In contrast, males of *P. lacunosus* have the dorsal shields covered with numerous rounded lacunae, the prodorsal shield entire and fused with the scapular shields, solenidion σ of genu III equal to or even longer than the segment, setae *si*, *c2* spiculiform, and the suckers with indentations; females of *P. lacunosus* have setae *d2* and *e2* long spiculiform, the former setae are much longer (100µm) and distinctly surpass the bases of corresponding setae *e2*.

Description

Male (Figures 1, 3A–E) (holotype, range for 6 paratypes in parenthesis) — Idiosoma short and wide, 252 (241–260) in length, 171 (174–190) in width. Gnathosoma 41 (41–43) × 52 (50–56). Prodorsal shield split into two pieces at the level of scapular setae, length of the anterior part along midline 47 (40–47), maximum width of posterior part 76 (74–80) (Fig. 1A). Distance between scapular setae *se-se* 58 (57–61). Setae *si* thin piliform. Scapular shields present, with poorly distinct borders, not fused with prodorsal shield. Humeral shields small, setae *c2* on soft tegument, mesal to these shields. Hysteronotal shield: greatest length 148 (142–149), width at anterior margin 110 (107–116), anterior end almost extending to level of setae *c1*, anterior margin roughly convex, surface with faint circular ornamentation. Hysteronotal gland openings *gl* situated at half distance between setae *d2* and *e2*. Setae

FIGURE 1: *Psittophagus hollandicus* n. sp. male: dorsal (A) and ventral (B) views.

c3 long, about half the idiosomal width. Subtegumental sclerotized structures in sejugal area absent. Opisthosomal lobes roughly rounded, with small extensions bearing bases of setae *h2*, *h3*. Terminal cleft small, as a wide-based triangle with rounded anterior end, 14 (10–14) in length. Supranal concavity semi-ovate. Setae *e2* spiculiform, noticeably thickened in distal third (somewhat shaped as base-

ball bat), 30 (28–32); setae *ps1* small, narrowly lanceolate, 15 (14–16) long. Dorsal measurements (distances between setal rows and seta bases): *c1-c1* 41 (40–46), *d2-d2* 86 (86–94), *e2-e2* 49 (49–54), *c1-d2* 66 (65–71), *d2-e2* 63 (61–63), *h2-h2* 51 (50–54), *ps1-ps1* 23 (23–26).

Epimerites I, II with inflated and heavily sclerotized bases, epimerites I free. Genital apparatus 15

FIGURE 2: *Psittophagus hollandicus* n. sp. female: dorsal (A) and ventral (B) views

(14–16) × 11 (9–12), aedeagus slightly shorter than genital arch. Setae 4a at base of genital arch. Setae 4b anterior to level of inner tips of epimerites IIIb. Adanal shields absent. Opisthoventral shields large, with projection towards anal suckers, carrying cupules *ih* and bases of setae *ps2* (Fig. 2B). Ventral measurements: 4b–3a 10 (7–10), 3a–g 31 (29–38), g–4a 34 (28–34), 4a–ps3 22 (22–24), ps3–ps3 18 (15–19).

Legs IV extending beyond opisthosoma by entire tarsus. Tarsus I with three long setae (*f*, *la*, *ra*), exceeding length of solenidion ω3, tarsus II with one long seta (*d*). Legs IV with tarsus entirely extending beyond level of lobar apices. Solenidion φ of tibia IV about ¾ the length of tarsus IV. Length of tarsi: I 30 (30–35), II 40 (39–43), III 34 (33–37), IV 37 (32–37).

Female (Figures 2, 3F–G) (range for 4 paratypes)

FIGURE 3: *Psittophagus hollandicus* n. sp.: dorsal view of male genua, tibiae and tarsi of legs I-IV (A-D) and opisthosoma (E); dorsal view of female genua, tibiae and tarsi III and IV (F-G).

— Idiosoma 350–386 in length, 224–250 in width. Gnathosoma 58–60 × 76–80. Prodorsal shield present only anterior to scapular setae, length along midline 54–61. Setae *si* thick spiculiform, 48–58 long. Distance between scapular setae *se-se* 76–83. Setae *si* thick spiculiform, 48–58 long. Scapular and humeral shields absent. Setae *c2* on striated tegument, slightly shorter than humeral seta *cp*. Setae *c3* long, about 1/2 of idiosomal width. Hysteronotal shield: occupying posterior half of hysterosoma, anterior margin convex, extending slightly anterior to level of setae *d2*, posterior margin indistinct, greatest length 139–148, surface with faint ornamentation. Setae *d2* piliform, about 30 long, situated on striated tegument, barely extending to level of openings *gl*; setae *e2* piliform, 20 long. Subtegumental sclerotized structures in sejugal region very large, ball-shaped, and dark-coloured; subtegumental structures of opisthosoma roughly rectangular and less sclerotized. Hysteronotal gland openings *gl* on soft tegument at level of trochanters IV. Setae *c3* long, about 1/2 of idiosomal width. Posterior end of opisthosoma wide and bluntly rounded. Dorsal measurements: *c1-c1* 62–65, *d2-d2* 110–125, *e2-e2* 108–115, *c1-d2* 85–89, *d2-e2* 65–72, *h2-h2* 66–76.

Epimerites I, II as in the male. Oviporus anterior to level of trochanters III, folds of this opening not sclerotized, epigynum absent. Legs IV with entire tarsus extending beyond posterior margin of opisthosoma. Length of tarsi: I 41–48, II 53–57, III 61–64, IV 74–77.

Etymology — The species name is taken from the specific epithet of the type host.

Remarks — 1) In addition to *Nymphicilichus perezae* from the cockatiel in Brazil, Albuquerque *et al.* (2012) reported an undetermined *Psittophagus* species quite probably corresponding to the new species described herein. However, the SEM photo of the female applied to that undetermined species (Albuquerque *et al.*, 2012: page 577, figure 7B) seemingly depicts the female of *N. perezae*, rather than that of *Psittophagus* species. This is suggested based on the length of setae *si* and the ornamentation of the hysteronotal shield.

2) The illustrations of *P. galahi* and *N. perezae* females in Mironov *et al.* (2003) were accidentally mixed up.

The correct legend for figure 9 in that paper should actually be: Fig. 9, a – *N. perezae*, b – *P. galahi*.

ACKNOWLEDGEMENTS

The author thanks David Vilas Boas Filho (Faculdade Jaguariúna, Jaguariúna, São Paulo State, Brazil) for collecting the dead specimen of cockatiel. The study was funded by FAPESP—São Paulo Research Foundation (2011/50145-0).

REFERENCES

- Albuquerque D.A.D., Brener B., Menna-Barreto R.F.S., Brun S.F. 2012 — The first identification of *Nymphicilichus perezae* Mironov and Galloway, 2002 in cockatiels in Brazil and the first record of *Psittophagus* sp. Gaud and Atyeo, 1996 and cf. *Dubininia* sp. Vassilev, 1958 in cockatiels (*Nymphicus hollandicus* Kerr, 1792) — *Parasitol. Int.*, 61: 572–578. doi:10.1016/j.parint.2012.05.006
- Atyeo W.T. 1985 — A new genus of pterolichid feather mites (Acari) from parrots Psittacidae — *J. Med. Entomol.*, 22: 54–57. doi:10.1093/jmedent/22.1.54
- Atyeo W.T. 1988 — Feather mites of the *Aralichus canestrinii* (Trouessart) complex (Acarina, Pterolichidae) from New World parrots (Psittacidae). I. From the genera *Ara* Lacépède and *Anodorhynchus* Spix. *Fieldiana, N.S., Zoology*, 47: 1–26. doi:10.5962/bhl.title.2982
- Atyeo W.T. 1989a — *Aralichus hastifolia* (Mégnin and Trouessart), a species of feather mite (Acarina, Pterolichidae) restricted to species of the parrot genus *Enicognathus* Gray (Aves, Psittacidae) — *J. Kans. Entomol. Soc.*, 62: 329–334.
- Atyeo W.T. 1989b — *Aralichus porrectus* (Mégnin & Trouessart) and related feather mite species (Acarina, Pterolichidae) from parrots of the genus *Brotoeris* Vigors (Aves, Psittacidae). — *Syst. Parasitol.*, 14: 101–111. doi:10.1007/BF00016904
- Atyeo W.T. 1989c — *Pararalichus* gen.n. (Acarina, Pterolichidae) from New World parrots (Aves, Psittacidae) — *Zool. Scripta*, 18: 331–346. doi: 10.1111/j.1463-6409.1989.tb00459.x
- Atyeo W.T., Gaud J. 1966 — The chaetotaxy of sarcotiform feather mites (Acarina: Analgoidea) — *J. Kans. Entomol. Soc.*, 39: 337–346. doi: http://www.jstor.org/stable/25083524
- Atyeo W.T., Gaud J. 1987 — Feather mites (Acarina) of the parakeet, *Melopsittacus undulatus* (Shaw)

- (Aves: Psittacidae) — J. Parasitol., 73: 203-206. doi:10.2307/3282367
- Atyeo W.T., Gaud J., Pérez T.M. 1984 — *Distigmesikya*, a new genus, and five species of feather mites (Acarina: Pterolichidae) from New World parrots (Aves: Psittacidae) — *Acarologia*, 25: 67-76.
- Atyeo W.T., Pérez T.M. 1982 — New taxa of psittacine feather mites (Acarina, Pterolichidae) — J. Parasitol., 68: 1158-1161. doi:10.2307/3281112
- Atyeo W.T., Pérez T.M. 1990 — Feather mites of the *Aralichus canestrinii* (Trouessart) complex (Acarina, Pterolichidae) from New World parrots (Psittacidae). II. From the genera *Aratinga* Spix, *Deroptyus* Wagler, *Leptopsittaca* Berlepsch & Stolzmann, *Ognorhynchus* Bonaparte, *Pionites* Heine, and *Pyrrhura* Bonaparte, and conclusions to the study — *Fieldiana*, N.S., Zoology, 62: 1-30. doi:10.5962/bhl.title.3205
- Dabert J., Mironov S.V., Ehrnsberger R. 2004 — New feather mite taxa of the *Rhytidelasma* generic group (Astigmata; Pterolichidae) from the Red-flanked Lorikeet *Charmosyna placensis* (Psittacidae) — *Syst. Parasitol.*, 58: 91-104. doi:10.1023/B:SYPA.0000029424.64857.ce
- Dabert J., Mironov S.V., Ehrnsberger R. 2008 — Systematic revision of the feather mite genera *Apexolichus* Gaud et Atyeo and *Titanolichus* Gaud et Atyeo (Astigmata: Pterolichidae), parasites of parrots of the Old World (Psittaciformes: Psittacidae) — *Acta Parasit.*, 53: 46-80. doi:10.2478/s11686-008-0007-x
- Dabert J., Mironov S.V., Proctor H.C. 2006 — Systematics of the feather mite genus *Titanolichus* Gaud & Atyeo, 1996 (Acari: Pterolichoidea) with description of one new species from the orange-bellied parrot *Neophema chrysogaster* (Aves: Psittaciformes) in Tasmania — *Aust. J. Entomol.*, 45: 206-214. doi:10.1111/j.1440-6055.2006.00535.x
- Dabert J., Badek A., Skoracki M. 2007 — New feather mite species (Acari, Astigmata) from the Sulphur-crested Cockatoo *Cacatua galerita* and Yellow-crested Cockatoo *C. sulphurea* (Psittaciformes, Cacatuidae) — *Acta Parasitol.*, 52(3): 250-267. doi:10.2478/s11686-007-0034-z
- Engelbreton M. 2006 — The welfare and suitability of parrots as companion animals: a review — *Anim. Welf.*, 15: 263-276.
- Gaud J. 1980 — Acariens Sarcopitiformes plumicoles parasites sur les oiseaux Psittaciformes, Strigiformes et Caprimulgiformes en Afrique — *Ann. Mus. Roy. Afr. Centr., Sér. In-8, Sces. Zool.*, 230: 1-106.
- Gaud J., Atyeo W.T. 1996 — Feather mites of the World (Acarina, Astigmata): the supraspecific taxa — *Ann. Mus. Roy. Afr. Centr., Sér. In-8, Sces. Zool.*, 277: 1-193 (Pt. 1, text), 1-436 (Pt. 2, illustrations).
- Gaud J., Mouchet J. 1959. — Acariens Plumicoles des oiseaux du Cameroun. V. Pterolichidae. — *Ann. Paras. Hum. Comp.*, 34: 493-545, 631-675.
- Griffiths D.A., Atyeo W.T., Norton R.A., Lynch C.A. 1990 — The idiosomal chaetotaxy of astigmatid mites — *J. Zool.*, 220: 1-32. doi: 10.1111/j.1469-7998.1990.tb04291.x
- Krantz G.W., Walter D.E. 2009 — *A Manual of Acarology*. 3rd ed. — Texas Tech University Press, Lubbock, 807 PP.
- Mégnin P., Trouessart E.L. 1884 — Les Sarcopitiformes plumicoles. — *J. Micrograph.*, 8: 92-101, 150-157, 211-219, 257-266, 331-338, 380-385, 428-436.
- Mironov S.V., Dabert J. 2007 — Three new feather mite genera of the *Protolichus* generic group (Astigmata: Pterolichidae) from parrots (Aves: Psittaciformes) of the Old World — *Acta Parasitol.*, 52: 386-402. doi:10.2478/s11686-007-0042-z
- Mironov S.V. & Dabert J. 2010 — A revision of the feather mite genus *Protolichus* Trouessart (Astigmata, Pterolichidae) — *Zootaxa*, 2526: 1-36. doi:10.11646/zootaxa.2526.1.1
- Mironov S.V., Dabert J., Ehrnsberger R. 2003a — A review of feather mites of the *Psittophagus* generic group (Astigmata: Pterolichidae) with descriptions of new taxa from parrots (Aves: Psittaciformes) of the Old World — *Acta Parasitol.*, 48: 280-293.
- Mironov S.V., Dabert J., Ehrnsberger R. 2005 — Six new feather mite species (Acari: Astigmata) from the carolina parakeet *Conuropsis carolinensis* (Psittaciformes: Psittacidae), an extinct parrot of North America. — *J. Nat. Hist.*, 39: 2257-2278. doi:10.1080/00222930400014155
- Mironov S.V., Dabert J., Ehrnsberger R. 2011 — A new feather mite species of the genus *Loriprotolichus* Mironov et Dabert, 2007 (Acariformes: Pterolichidae) from the ultramarine lorikeet *Vini ultramarina* (Aves: Psittaciformes) — *Acarina*, 19: 77-82.
- Mironov S.V., Dabert J., Proctor H.C. 2003b — New feather mites of the family Pterolichidae (Acari: Pterolichoidea) from parrots (Aves: Psittaciformes) in Australia — *Aust. J. Entomol.*, 42: 185-202. doi:10.1046/j.1440-6055.2003.00348.x
- Mironov S.V., Ehrnsberger R., Dabert J. 2014 — New species of the feather mite genus *Protolichus* Trouessart, 1884 (Astigmata, Pterolichidae) from lorries and lorikeets (Aves: Psittaciformes). — *Zootaxa*, 3774: 131-151. doi:10.11646/zootaxa.3774.2.2
- Mironov S.V., Galloway T.D. 2002 — *Nymphilichus perezae* gen. nov., n. sp., a new feather mite (Astigmata: Pterolichidae) from the cockatiel, *Nymphicus hollandicus* (Psittaciformes: Cacatuidae) — *J. R. Soc. N. Z.*, 32: 1-6. doi:10.1080/03014223.2002.9517682

Hernandes F.A.

Mironov S.V., Pérez T.M. 2003 — A review of feather mites of the *Rhytidelasma* generic group (Pterolichoidea Pterolichidae), specific parasites of parrots (Aves Psittaciformes) — Bull. Inst. R. Sci. Nat. Belg. Entomol., 73: 135-176.

Norton R. 1998 — Morphological evidence for the evolutionary origin of Astigmata (Acari: Acariformes) — Exp. Appl. Acarol., 22: 559-594. doi: [10.1023/A:1006135509248](https://doi.org/10.1023/A:1006135509248)

COPYRIGHT

Hernandez F.A. Acarologia is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.