

First records of Phytoseiidae (Acari: Mesostigmata) from one island of the Comoros Archipelago

Serge Kreiter^a, Rose-My Payet^b, Jacques Fillâtre^c, Hamza Abdou Azali^d

^a CBGP, Montpellier SupAgro, INRA, CIRAD, IRD, Univ Montpellier, Montpellier, France.

^b CIRAD, UPR Hortsys, Station de Bassin-Plat, 97410, Saint-Pierre, Réunion, France.

^c Armefflor, 1 Chemin de l'IRFA, 97410 Saint-Pierre, Réunion, France.

^d INRAPE, Moroni, Grande Comore, Union des Comores.

ABSTRACT

The Comoros Archipelago is constituted of four islands. These islands are located in the North Mozambique Channel in the Indian Ocean, one of the world's hotspots of biodiversity. Despite this status of hotspot, only one species of Phytoseiidae was known from this Archipelago, from Mayotte: *Phytoseius mayottae*. No species were recorded from the three other main islands. We report in this paper the results of a preliminary survey in Great Comoro or "Grande Comore" Island also called Ngazidja in the Comorian language (= Shikomori) with five species recorded.

Keywords Survey, collection, taxonomy, systematics, Grande Comore

Zoobank <http://zoobank.org/BABDD080-D2D0-46F6-8172-442298192D42>

Introduction

Several species in the family Phytoseiidae are important natural enemies controlling phytophagous mite and small insects in natural areas and crops all around the world (McMurtry and Croft 1997; McMurtry *et al.* 2013).

This family is widespread all over the world and consists of 2,479 valid species dispatched in three sub-families and 94 genera (Demite *et al.* 2017). Most of areas of the Indian Ocean constitute one of the world's hotspots of biodiversity. The hotspot of biodiversity concept was defined by Myers (1988) in order to identify the most immediately important areas for conservation of biodiversity. These hotspots hold high endemism levels and have lost at least 70% of their original natural vegetation (Myers *et al.* 2000). The characterization of the phytoseiid mite diversity in these areas is thus contributing to this general topic of conservation.

Located in the Indian Ocean in the North of Mozambique Channel, Comoros Archipelago is constituted of four islands: "Grande Comore" called also Ngazidja in Comorian language (= Shikomori), Mohéli called also Mweli, Anjouan called also Nzouani, and Mayotte called also Maoré. The climate of the Comoros Archipelago, south of the Equator, is tropical, with a hot and rainy season from December to April, and a relatively cool and dry season from May to November. The average daily temperature goes from around 27°C in the warmest period (January to April), to around 23°C in the coolest months (July, August and September). The two smaller islands (Mohéli and Anjouan) are covered by hills in the interior, while Grande Comore has the volcano Karthala, 2,360 meters high. This main island is the rainiest. On Mount Karthala, even 5,000 mm of rain / year fall, while the capital Moroni, which is located on the windward side receives 2,700 mm of rain / year. On the Comoros there are numerous tropical ecosystems that are primarily dependent on the altitude. Many kinds of tropical plants can be found, large numbers of which are endemic. Like most islands, the diversity of the local

Received 28 November 2017

Accepted 10 January 2018

Published 01 June 2018

Corresponding author

Serge Kreiter:

serge.kreiter@supagro.fr

Academic editor

Marie-Stéphane Tixier

DOI

10.24349/acarologia/20184256

© Copyright

Kreiter S. *et al.*

Distributed under

Creative Commons CC-BY 4.0



How to cite this article Kreiter S. *et al.* (2018), First records of Phytoseiidae (Acari: Mesostigmata) from one island of the Comoros Archipelago. *Acarologia* 58(3): 529-545; DOI 10.24349/acarologia/20184256

flora suffers from two pressures, on one hand the reduction of available space caused by the reduction of biotopes due to the invasion of humans in wildest areas and on the other hand, the invasion of exotic plant species such as guava trees.

Presently, only one species of Phytoseiidae is known from this part of the world, from Mayotte: *Phytoseius mayottae* Schicha. This species was collected by Jean Gutierrez and described by Schicha (1984). No species are recorded from the three other islands.

We report in this paper result of a preliminary survey in Grande Comore Island (Ngazidja).

Material and methods

The survey took place in “Grande Comore” (or Ngazidja) in February 2017. Plant inhabiting mites were collected for this preliminary survey only from various crops in two locations. Mites were directly collected from leaves with a fine brush and then transferred into small plastic vials containing 70% and 98% ethanol. Plant species were identified by Jacques Fillâtre (Armefflor).

Mites in vials with 70% ethanol were then all mounted on slides using Hoyer’s medium and all identified using a phase and interferential contrast microscope (Leica DMLB, Leica Microsystems SAS, Nanterre, France). Mites in vials with 98% ethanol will be used for barcoding (in progress).

Morphological characters were measured using a graduate eyepiece (Leica, see above). Chant and McMurtry’s (1994, 2007) concepts of the taxonomy for the family Phytoseiidae and the world catalogue database of Demite *et al.* (2017) were used for faunistical and biogeographical aspects. The chaetotaxy terminologies used in this paper followed those proposed by Lindquist and Evans (1965) as adapted by Rowell *et al.* (1978) for Phytoseiidae for dorsal and by Chant and Yoshida-Shaul (1991) for ventral idiosomal setae, respectively. Adenotaxy and poroidotaxy terminologies are those proposed by Athias-Henriot (1975).

Numbers of teeth on the fixed and movable cheliceral digits do not include the respective apical teeth. Setae not referred to in the Results section should be considered as absent.

All measurements are given in micrometers (μm) and presented as the mean in bold followed by the range in parenthesis. All individuals collected were measured.

Specimens of each species are deposited in the mite collections of Montpellier SupAgro conserved in UMR CBGP INRA/IRD/CIRAD/SupAgro.

The following abbreviations are used in this paper for morphological characters: **dsl** = dorsal shield length; **dsw** = dorsal shield width; **lisl** = Largest inguinal sigilla (= “metapodal plate”) length; **lisw** = Largest inguinal sigilla (= “metapodal plate”) width; **sisl** = smallest inguinal sigilla (= “metapodal plate”) length; **vsl** = ventrianal shield length (or ventral shield length for *Iphiseius degenerans*); **asl** = anal shield length; **vsw ZV2** and **vsw anus** = ventrianal shield width at ZV2 level and at anus level; **scl** = spermatheca cervix length; **scw** = spermatheca cervix width; **fdl** = fixed digit length; **mdl** = movable digit length.

The following abbreviations are used in this paper for institutions: **Armefflor** = Association Réunionnaise pour la Modernisation de l’Economie Fruitière, Légumière et HORTICOLE; **CBGP** = Centre de Biologie pour la Gestion des Populations; **CIRAD** = Centre International de Recherche Agronomique pour le Développement; **INRA** = Institut National de la Recherche Agronomique; **INRAPE** = Institut National de Recherche pour l’Agriculture, la Pêche et l’Environnement; **IRD** = Institut de Recherche pour le Développement; **MSA** = Montpellier SupAgro, France; **UMR** = Unité Mixte de Recherche; **UPR** = Unité Propre de Recherche.

Results and discussion

A total of five species have been collected and identified, four species of the sub-family Amblyseinae and one species of the sub-family Phytoseiinae. Three of them are biological control agents, may have great interest for agriculture of this part of the world and some data from the literature are provided here.

Subfamily Amblyseinae Muma

Amblyseinae Muma, 1961: 273.

Tribe Amblyseini Muma

Amblyseini, Muma, 1961: 68.

Subtribe Amblyseina Muma

Amblyseina Muma, 1961: 69.

Genus *Amblyseius* Berlese

Amblyseius Berlese, 1914: 143.

Amblyseius herbicolus (Chant)

Typhlodromus (*Amblyseius*) *herbicolus* Chant, 1959: 84;

Amblyseius (*Amblyseius*) *herbicolus*, Muma 1961: 287;

Typhlodromus herbicolus, Hirschmann, 1962: 23;

Amblyseius herbicolus, Moraes *et al.*, 1986: 14; 1989: 79; 2004: 27; Chant & McMurtry, 2004: 209; 2007: 78;

Amblyseius deleoni Muma & Denmark, 1970: 68 (synonymy according to Daneshvar & Denmark, 1982; Denmark & Muma 1989);

Amblyseius giganticus Gupta, 1981: 33 (synonymy according to Gupta, 1986);

Amblyseius impactus Chaudhri, 1968, 553 (synonymy according to Daneshvar & Denmark, 1982; Denmark & Muma, 1989);

Amblyseius (*Amblyseialus*) *thermophilus* Karg, 1991: 12 (synonymy according to El-Banhawy & Knapp, 2011 and to Demite *et al.*, 2017);

Typhlodromus (*Amblyseius*) *amitae* Bhattacharyya, 1968: 677 (synonymy according to Denmark & Muma, 1989).

Amblyseius herbicolus (Chant) is the second most abundant phytoseiid mites on coffee plants (*Coffea arabica* L.) in Brazil, associated with *Brevipalpus phoenicis* (Geijskes) (Acari: Tenuipalpidae), vector of the coffee ring spot virus. *Amblyseius herbicolus* was found to be an efficient predator of the coffee ring spot mite (Reis *et al.* 2007).

Amblyseius herbicolus is also found associated with the broad mite, *Polyphagotarsonemus latus* Banks in crops such as chili pepper (*Capsicum annum* L.) in Brazil. This species has also a good potential for controlling *P. latus*. Rodríguez-Cruz *et al.* (2013) have studied biological, reproductive and life table parameters of *A. herbicolus* on three different diets: broad mites, castor bean pollen (*Ricinus communis* L.) and sunn hemp pollen (*Crotalaria juncea* L.). The predator was able to develop and reproduce on all these three diets. However, its intrinsic growth rate was higher on broad mites and castor bean pollen. Feeding on alternative food such as pollen can facilitate the predator's mass rearing and maintain its population on crops when prey is absent or scarce. Many polyphagous generalist phytoseiid mites are important natural enemies because they can feed on plant provided pollen and various prey species, and thus persist in crops even in the absence of target pests (McMurtry *et al.* 2013). Hence, populations

of these predators can be established in a crop by providing alternative food, thus increasing biological control. Alternative food affects *P. latus* control on chilli pepper plants by predatory mites (Duarte *et al.* 2015). *Amblyseius herbicolus* had high oviposition and population growth rates when fed with cattail pollen (*Typha latifolia* L.), chilli pepper pollen and bee-collected pollen, and a low rate on the alternative prey *Tetranychus urticae* Koch. Supplementing pepper plants with pollen resulted in better control of broad mite populations (Duarte *et al.* 2015). Release of *A. herbicolus* on young plants with weekly addition of honeybee pollen or cattail pollen until plants produce flowers seems a viable strategy to sustain populations of this predator (Duarte *et al.* 2015).

Specimens examined — **Moroni**, Adoudja (long. 11°41'S, lat. 43°15'E, alt. 100 m), 1 ♀ + 1 immature on *Alocasia macrorrhiza* (L.) G. Don (Araceae), 2-02-2017; 1 ♀ + 2 immatures on *Morinda citrifolia* L. (Rubiaceae), 2-02-2017.

Previous record — Large distribution worldwide. This species is mentioned from Mozambique, La Réunion Island, Kenya, Tanzania for the closest places.

Remarks (Table 1) — Measurements of the two females collected fit very well with the measurements reported literature, except for greater dimension of the ventrianal shield and a longer spermatheca for specimens from Grande Comore.

Tribe Euseiini Chant and McMurtry

Euseiini Chant & McMurtry, 2005b: 191.

Subtribe Euseiina Chant and McMurtry

Euseiina Chant & McMurtry, 2005b: 209.

Genus *Euseius* Wainstein

Amblyseius (*Amblyseius*) section *Euseius*, Wainstein, 1962: 15; *Euseius* De Leon, 1967: 86.

Euseius baetae (Meyer & Rodrigues)

Amblyseius baetae (Meyer & Rodrigues, 1966): 28;

Euseius baetae (Meyer & Rodrigues), Moraes *et al.*, 2001a: 11; Moraes *et al.*, 1986: 37; 2004: 62; Chant & McMurtry 2005b: 215; 2007: 120; El-Banhawy & Knapp, 2011: 36;

Euseius kangwanensis Ueckermann & Loots, 1988: 85 (synonym according to Ueckermann & Loots, 1988; Moraes *et al.*, 2001b; El-Banhawy & Knapp, 2011).

The 200 species of the genus *Euseius* are considered as Type IV species, polliniphagous generalists (McMurtry and Croft 1997; McMurtry *et al.* 2013) and *Euseius baetae* is supposed to belong to that Type IV. The biology of *E. baetae* is however totally unknown.

Specimens examined — **Mdé**, INRAPE (long. 11°41'S, lat. 43°14'E, alt. 50 m), 1 ♀ on *Manihot esculenta* Crantz (Euphorbiaceae), 2-02-2017; **Moroni**, Adoudja (long. 11°41'S, lat. 43°15'E, alt. 100 m), 2 ♀♀ on *Plectranthus scutellarioides* (L.) R.Br. (Lamiaceae), 2-02-2017; 1 ♀ on *Alocasia macrorrhizos* (L.) G. Don (Araceae), 2-02-2017.

Previous record — Congo, Kenya, Malawi, Mozambique, South Africa.

Remarks (Table 2) — Measurements of the 4 ♀♀ fit well with the measurements from the literature with slightly shorter dimensions in general.

Table 1 Comparisons of character measurements of female specimens of *Amblyseius herbicolus* collected in different locations (Localities followed by the number of specimens measured between brackets)

Characters	Grande Comore (2)	Kenya (10)	Various countries (8)	Senegal (2)	Turkey (3)	Holotype (1)
Dsl	385-428	335	353 (325-368)	360-365	313-352	369
Dsw	263-275	190	256 (240-274)	250-268	196-221	236
j1	40-43	32	37 (34-40)	35-38	33-38	38
j3	53-55	37	49 (38-58)	38-45	32-39	42
j4	6	4-6	6 (5-8)	6-8	8-9	9
j5	5	4-6	4 (3-5)	4	7-8	7
j6	5-6	4-6	7 (5-8)	8	6-7	11
J2	8-10	4-6	10 (8-11)	8-9	8-9	12
J5	10	4-6	8 (6-10)	8-10	9-10	9
z2	6-8	6	11 (8-16)	8-9	9-12	13
z4	8	6	8 (8-10)	10-11	8-12	9
z5	5	6	6 (5-6)	7-9	6-7	6
Z1	8	10	10 (8-13)	8-10	9-12	9
Z4	133-135	90	126 (101-152)	163-172	91-99	110
Z5	288-300	232	281 (251-306)	310-345	220-251	236
s4	120-125	92	113 (98-130)	123-135	86-96	100
S2	13	10	12 (8-14)	13-15	10-12	11
S4	13	10	11 (8-13)	8-10	9-11	13
S5	13	10	9 (8-10)	8-10	9-10	11
r3	8-15	10	14 (11-16)	10-11	10-16	15
R1	10-13	10	9 (8-10)	9-10	8-9	8
st1-st1	68	nr	not reported	nr	nr	nr
st1-st3	73-75	62	65 (58-70)	63-65	64-69	
st2-st2	73-75	69	71 (66-75)	75-78	70-73	
st3-st3	63-65	nr	not reported	nr	nr	nr
st4-st4	78-83	nr	not reported	nr	nr	nr
st5-st5	70-78	nr	71 (67-75)	65-70	58-64	
lisl	23	nr	not reported	nr	nr	nr
lsw	5	nr	not reported	nr	nr	nr
sisl	13	nr	not reported	nr	nr	nr
vsl	120-135	108	116 (112-118)	113-115	102-117	nr
vsw ZV2	63	48	57 (53-59)	65-69	44-48	nr
vsw anus	80	69	71 (66-77)	75-78	61-69	nr
JV5	80-85	52	not reported	nr	51-60	nr
SgeI	45-48	nr	42 (35-48)	43-48	40-48	nr
SgeII	38	52	38 (35-42)	38-40	34-39	nr
SgeIII	53-58	41	52 (45-59)	48-52	41-46	nr
StiIII	43	35	41 (34-48)	43-45	33-40	nr
SgeIV	135	110	124 (96-158)	160-162	98-128	112
StiIV	100	76	90 (67-109)	102-15	75-89	82
StIV	80	65	76 (66-86)	76-78	63-70	76
scl	38-40	nr	32 (32-35)	33-35	24-32	18
scw	2-3	nr	not reported	nr	nr	nr
Fdl	33	nr	31 (30-31)	nr	29-36	nr
teeth	11-12	8	10	nr	12	11-12
Mdl	30	nr	35 (34-35)	nr	31-33	nr
teeth	?	3	3	nr	4	4

Data from this study for Grande Comore, from El-Banhawy and Knapp (2011) for Kenya, from Zannou *et al.* (2007) for various countries of Africa (Benin 1, Burundi 1, Democratic Republic of Congo 1, Ghana 1, Kenya 3, Rwanda 1), from Kade *et al.* (2011) for Senegal, from Akyasi *et al.* (2016) for Turkey, from Denmark and Muma (1989) for the holotype intercepted in Massachusetts (Boston) but coming from Portugal. nr = not reported.

Table 2 Comparisons of character measurements of female specimens of *Euseius baetae* collected in different locations (Localities followed by the number of specimens measured between brackets)

Characters	Grande Comore (4)	Mozambique (4)	South Africa (7)	Congo (1)	Kenya (1)
Dsl	341 (310-375)	322-336	337 (343-359)	336	300
Dsw	329 (220-238)	201-243	252 (224-239)	230	210
j1	28	29-38	32 (28-32)	29	21
j3	8 (6-10)	11-12	9 (8-11)	16	25
j4	6 (5-8)	9-10	9 (8-11)	11	12
j5	7 (6-8)	9	9 (8-11)	11	12
j6	8 (6-8)	10-11	9 (8-11)	16	12
J2	9 (8-10)	12-13	11 (11-13)	13	13
J5	7 (5-8)	7-9	6	8	7
z2	9 (8-10)	11-12	9 (8-11)	14	14
z4	9 (8-10)	11-12	9 (8-11)	13	20
z5	7 (5-8)	9-10	9 (8-11)	11	12
Z1	9 (9-10)	10-12	9 (8-11)	11	12
Z4	10	11-14	11 (11-13)	16	15
Z5	59 (55-65)	59-65	55 (55-60)	53	50
s4	14 (13-14)	16-20	16 (16-19)	21	30
S2	9 (8-10)	12-14	11 (11-13)	14	16
S4	10 (9-10)	12-16	11 (11-13)	14	16
S5	9 (8-10)	11-13	11 (11-13)	14	14
r3	14 (13-15)	not reported	13 (13-16)	19	14
R1	9 (8-10)	not reported	9	11	14
St1-St1	57 (55-58)	not reported	not reported	nr	nr
St1-St3	58 (55-60)	77-86	88 (82-91)	58	58
St2-St2	67 (65-68)	not reported	not reported	70	70
St3-St3	74 (70-78)	71-78	95 (91-101)	nr	nr
St4-St4	79 (78-80)	not reported	not reported	nr	nr
St5-St5	72 (68-75)	not reported	88 (82-91)	74	80
Lisl	26 (25-28)	not reported	not reported	nr	nr
Lsiw	2	not reported	not reported	nr	nr
Sisl	10 (8-13)	not reported	not reported	nr	nr
Vsl	91 (78-100)	95-108	88 (82-91)	?	100
vsw ZV2	49 (45-50)	69	69 (69-76)	48	?
Vsw anus	71 (70-73)	not reported	not reported	67	70
JV5	39 (38-40)	not reported	38 (38-47)	nr	37
SgeI	22 (20-25)	not reported	28	nr	nr
SgeII	27 (25-28)	not reported	32 (32-35)	29	25
SgeIII	35	not reported	38 (41-44)	38	30
StiIII	27 (25-28)	not reported	32 (32-35)	29	nr
SgeIV	57 (55-58)	64-72	72 (63-72)	56	58
StiIV	38	40-47	41 (44-47)	38	51
StIV	67 (65-68)	71-80	76 (76-79)	70	95
scl	15	16-18	not reported	19	24
scw	5	not reported	not reported	nr	5
Fdl	25	not reported	nr	nr	nr
teeth	invisible	not reported	6	nr	4
Mdl	24 (23-25)	not reported	nr	nr	nr
teeth	invisible	not reported	2	nr	2

Data from this study for Grande Comore, from Meyer and Rodrigues (1966) for Mozambique holotype + paratype females, from Ueckermann and Loots (1988) for South Africa (measurements for holotype and between brackets for 6 paratype females), from Moraes *et al.* (2001b) for Congo and from El-Banhawy and Knapp (2011) for Kenya. nr = not reported.

Genus *Iphiseius* Berlese

Iphiseius Berlese, 1916: 33; Chant & McMurtry, 2005b: 217; 2007: 123.

Iphiseius degenerans (Berlese)

Seius degenerans (Berlese, 1889): 9;

Amblyseius (Iphiseius) degenerans, Muma, 1961: 288;

Typhlodromus degenerans, Hirschmann, 1962: 2;

Iphiseius (Iphiseius) degenerans, Pritchard & Baker 1962: 299;

Amblyseius degenerans, Zaher, 1986: 99; Northcraft, 1987: 521; Papadoulis & Emmanouel, 1991: 36;

Iphiseius degenerans, Berlese, 1921: 95; Evans, 1954: 518; Moraes *et al.*, 1986: 61; 2004: 92; Chant & McMurtry, 2005b: 215; 2007: 125;

Iphiseius martigellus El-Badry, 1968: 325 (synonymy according to Chant & McMurtry, 2005; El-Banhawy & Knapp, 2011).

The biological characteristics of this Ethiopian species have been well documented because of its use in controlling thrips on various cultivated plants in greenhouses. *Iphiseius degenerans* is a commercially available biological control agent of thrips and spider mites in greenhouse crops. It is able to feed on a variety of foods, but thrips' larvae and sweet pepper pollen are unfavourable food for immature development. This could compromise the establishment of this biological control agent when used against thrips in sweet pepper crops. According to the classification by McMurtry *et al.* (2013), *I. degenerans* is a type-III generalist predator. It is one of the most common native phytoseiid mite species on cassava in southern Africa (Zannou *et al.* 2005) and feeds on *Mononychellus tanajoa* (Bondar) (Nwilene and Nachman 1996), a widely distributed neotropical mite pest of cassava in Africa, insect larvae and pollen of many plants (Vantornhout *et al.* 2005).

Another study concluded that *I. degenerans* can be considered a suitable biological control candidate based on its preference for *Eutetranychus orientalis* nec (Klein) in the Mediterranean region (Fantinou *et al.* 2012).

Iphiseius degenerans preys on *Oligonychus perseae* Tuttle, Baker & Abbatiello outside the webbed nests. Although *I. degenerans* contribution to *O. perseae* biocontrol can be limited, it needs to be assessed, also taking into account the importance of alternative food source (e.g. Castor oil pollen) for predator population growth (Zappala *et al.* 2015).

Specimens examined — **MdÉ**, INRAPE (long. 11°41'S, lat. 43°14'E, alt. 50 m), 7 ♀♀ + 4 ♂♂ on *Ricinus communis* L. (Euphorbiaceae), 2-02-2017.

Previous record — Numerous countries in Northern and Southern Africa (Demite *et al.* 2017), in Mediterranean area (Cyprus, Greece, Italy, Portugal), in Near East or Middle East (Egypt, Israel, Lebanon, Saudi Arabia, Syria, Turkey, Yemen), in Europe (Georgia), in South America (Brazil) and in North America (USA in California, Florida, Georgia, New Hampshire).

Remarks (Tables 3 and 4) — Measurements of the 7 ♀♀ + 4 ♂♂ fit well with measurements of specimens reported in the literature for closest countries, except width of the ventral and the anal shields of the female and JV5 which is more than 40 % longer.

Table 3 Comparisons of character measurements of female specimens of *Iphiseius degenerans* collected in different locations (Localities followed by the number of specimens measured between brackets)

Characters	Grande Comore (7)	Kenya (66)	Various countries (12)
Dsl	420 (370-462)	360	373 (363-387)
Dsw	362 (350-375)	345	309 (290-317)
j1	24 (20-25)	23	24 (14-32)
j3	5	Not reported	5 (4-8)
j4	5	Not reported	4 (3-6)
j5	5	Not reported	4 (3-5)
j6	5	Not reported	5 (4-6)
J2	5	Not reported	6 (4-7)
J5	5	Not reported	6 (5-7)
z2	5	Not reported	5 (4-8)
z4	5	Not reported	5 (4-9)
z5	5	Not reported	4 (3-5)
Z1	5	Not reported	6 (5-8)
Z4	8 (6-9)	Not reported	7 (5-9)
Z5	14 (11-15)	12	14 (11-20)
s4	8	Not reported	9 (7-12)
S2	8	Not reported	8 (6-9)
S4	9 (8-9)	Not reported	9 (6-11)
S5	9 (8-9)	Not reported	9 (7-11)
r3	14 (13-15)	Not reported	12 (10-13)
R1	11 (9-13)	Not reported	9 (7-10)
St1-St1	57 (55-60)	Not reported	Not reported
St1-St3	54 (53-55)	46	49 (42-52)
St2-St2	71 (68-75)	64	71 (67-73)
St3-St3	86 (65-93)	Not reported	Not reported
St4-St4	96 (90-100)	Not reported	Not reported
St5-St5	96 (90-103)	100	96 (91-103)
Lisl	36 (30-38)	Not reported	Not reported
Lsiw	5 (4-8)	Not reported	Not reported
Sisl	14 (13-18)	Not reported	Not reported
Vsl	35 (33-38)	Not reported	Not reported
vsw ZV2	77 (70-83)	74	65 (58-73)
Asl	72 (68-75)	Not reported	Not reported
Vsw anus	77 (70-83)	Not reported	72 (68-77)
JV5	24 (20-25)	14	Not reported
SgeII	24 (21-25)	23	Not reported
SgeIII	30 (28-33)	35	Not reported
StiIII	25 (25-26)	Not reported	Not reported
SgeIV	43 (40-50)	30	Not reported
StiIV	29 (25-33)	25	Not reported
StIV	32 (30-33)	38	Not reported
Scl	Not reported	Not reported	Not reported
Scw	Not reported	Not reported	Not reported
Fdl	27 (25-30)	Not reported	26 (24-27)
teeth	7	Not reported	6-8
Mdl	25 (23-27)	Not reported	31 (27-35)
teeth	1	Not reported	1-2

Data from this study for Grande Comore, from El-Banhawy and Knapp (2011) for Kenya, from Moraes *et al.* (2007) for various countries (Burundi 1, Cameroon 1, Ghana 1, Kenya 2, Malawi 2, Rwanda 1, Sierra Leone 2, Uganda 1, Zambia 1) in Africa. nr = not reported.

Table 4 Comparisons of character measurements of male specimens of *Iphiseius degenerans* collected in different locations (Localities followed by the number of specimens measured between brackets)

Characters	Grande Comore (4)	Kenya (?)	Various countries (5)
Dsl	306 (295-312)	Not reported	318 (283-356)
Dsw	232 (238-250)	Not reported	255 (244-261)
j1	21 (18-23)	Not reported	20 (11-25)
j3	5	Not reported	5 (4-5)
j4	5	Not reported	3 (3-5)
j5	5	Not reported	4 (4-5)
j6	5	Not reported	4 (3-4)
J5	5	Not reported	5 (4-6)
z2	5	Not reported	4 (3-5)
z4	5	Not reported	4 (3-5)
z5	5	Not reported	3 (3-4)
Z1	5	Not reported	5
Z4	8	Not reported	5 (4-5)
Z5	14 (13-15)	Not reported	13 (9-16)
s4	5	Not reported	7 (6-9)
S2	7	Not reported	6 (5-7)
S4	7	Not reported	6 (5-8)
S5	7	Not reported	7 (5-10)
r3	12 (10-13)	Not reported	11 (10-12)
R1	6 (5-9)	Not reported	7 (6-9)
St1-St1	49 (48-50)	Not reported	Not reported
St1-St5	114 (110-118)	Not reported	Not reported
St2-St2	62 (60-63)	Not reported	Not reported
St3-St3	64 (60-68)	Not reported	Not reported
St4-St4	60 (58-63)	Not reported	Not reported
St5-St5	52 (48-58)	Not reported	Not reported
Vsl	71 (68-73)	Not reported	Not reported
vsw ZV2	171 (165-175)	210	176 (172-179)
Asl	51 (48-55)	Not reported	Not reported
Asw	64 (58-68)	75	66 (63-75)
JV5	22 (20-23)	Not reported	Not reported
SgeII	21 (18-25)	Not reported	21 (18-24)
SgeIII	29 (28-30)	Not reported	27 (21-32)
StiIII	24 (23-25)	Not reported	21 (20-24)
SgeIV	42 (40-43)	Not reported	36 (30-40)
StiIV	30 (28-30)	Not reported	28 (27-30)
StIV	31 (30-33)	Not reported	28 (26-32)
Shaft length	24 (23-25)	22	29 (22-45)
Fdl	23 (20-25)	Not reported	Not reported
teeth	4	Not reported	Not reported
Mdl	22 (20-23)	Not reported	Not reported
teeth	1	Not reported	Not reported

Data from this study for Grande Comore, from El-Banhawy and Knapp (2011) for Kenya, from Moraes *et al.* (2007) for various countries (Ghana 1, Kenya 1, Rwanda 1, Sierra Leone1, Uganda 1) in Africa. nr = not reported.

Tribe Neoseiulini Chant and McMurtry

Neoseiulini Chant & McMurtry, 2003a: 6.

Genus *Neoseiulus* Hughes

Neoseiulus Hughes, 1948: 141.

Neoseiulus longispinosus (Evans)

Typhlodromus longispinosus Evans, 1952: 413; Evans, 1953: 465; Womersley, 1954: 177; Ehara, 1958: 55;
Typhlodromus (Amblyseius) longispinosus, Chant, 1959: 74;
Amblyseius longispinosus, Corpuz and Rimando, 1966: 129; Schicha, 1975: 103;
Neoseiulus longispinosus, Moraes *et al.*, 1986: 85; 2000: 245; 2004: 129; Chant & McMurtry 2003a: 37; 2007: 29.

This species is distributed in many countries of the world, mainly in tropical areas, especially in Guadeloupe and other Islands of the French Antilles (Moraes *et al.* 2000; Mailloux *et al.* 2010; Kreiter *et al.* 2013; Kreiter *et al.* in press).

Neoseiulus longispinosus, a type II phytoseiid predatory mite (McMurtry *et al.* 2013), has received increasing attention in Asia for the control of spider mites since 2010 (Nusartlert *et al.* 2011). It can develop on different tetranychid species of the genera *Eutetranychus*, *Oligonychus*, and *Tetranychus* (Nusartlert *et al.* 2011). Several studies demonstrated the potential of the predatory mite to control spider mite outbreaks including *Oligonychus coffeae* (Nietner) on tea (Rahman *et al.* 2013), *Stigmaeopsis nanjingensis* (Ma & Yuan) on bamboo in China (Zhang *et al.* 1999) or *Eotetranychus cendanai* Rimando in greenhouse crops (Thongtab *et al.* 2001). In addition, *N. longispinosus* was also found to have a great potential for practical applications due to its resistance or tolerance to agricultural chemicals (Zhang *et al.* 1996).

Thus, the biology of this species has been studied mostly for pest control purposes including side effects of miticides (Bin Ibrahim and Tan 2000). The activity, feeding, development, predation, cannibalism, intra-guild predation and behaviour have been extensively studied by several authors (Croft *et al.* 1999a, b; Schausberger and Croft 1999a, b; 2000a, b; Blackwood *et al.* 2001). It was found very rarely in Mascareignes area except in a study on companion plants in citrus orchards in La Réunion (Le Bellec, unpublished data). This species seems to be more common on grasses of the lower vegetation, especially Fabaceae with populations of tetranychid mites. However, the recent results of Huyen *et al.* (2017) show that at least in controlled laboratory conditions the predatory mite *N. longispinosus* is a potential biological control agent against the citrus red spider mite *P. citri*.

Previous Records — Australia, China (Fujian, Guangdong, Guangxi, Hainan, Yunnan), Cuba, Dominican Republic, Guadeloupe, Egypt, Hawaii, Hong-Kong, India, Indonesia, Japan, Les Saintes, Malaysia, Marie-Galante, Martinique, New Zealand, Nicaragua, Pakistan, Papua New Guinea, Philippines, Russia, South Korea, Sri Lanka, Taiwan, Thailand, USA (Florida), Vietnam.

Specimens examined — **Moroni**, Adoudja (long. 11°41'S, lat. 43°15'E, alt. 100 m), 1 ♀ on *Plectranthus scutellarioides* (L.) R. Br. (Lamiaceae), 2-02-2017.

Remarks (Table 5) — Measurements of the single female collected show that most of the setae lengths are 7 to 15 % longer, except *j6*, *J2*, *J5*, and *Z4*. All ventral, spermathecal and cheliceral dimensions agree well except ventrianal shield length and width at the level of anus and *JV5* which are longer.

Table 5 Comparisons of character measurements of female specimens of *Neoseiulus longispinosus* collected in different locations (Localities followed by the number of specimens measured between brackets)

Characters	Grande Comore (8)	Martinique (8)	F.C.I. (7)	Sri Lanka (3)	Holotype
Dsl	380	321 (295-340)	332 (308-398)	321 (313-338)	332
Dsw	192	168 (150-183)	179 (154-200)	187 (175-208)	173
j1	20	16 (13-18)	18 (16-22)	18 (17-19)	14
j3	69	58 (48-63)	59 (52-64)	62 (61-64)	51
j4	70	56 (50-65)	59 (52-65)	58 (56-60)	49
j5	78	66 (60-73)	69 (64-75)	70 (69-71)	59
j6	78	68 (60-83)	72 (68-75)	70 (68-72)	64
J2	88	76 (68-88)	76 (73-78)	77 (75-79)	66
J5	10	9 (8-10)	9 (8-11)	8	10
z2	75	64 (58-70)	65 (62-68)	69 (68-70)	58
z4	78	70 (63-87)	69 (67-73)	73 (73-75)	58
z5	38	31 (28-35)	35 (32-40)	32 (32-38)	nr
Z1	83	74 (68-80)	75 (72-80)	77 (76-78)	67
Z4	78	69 (63-78)	71 (67-75)	72 (71-73)	68
Z5	85	77 (65-80)	80 (78-81)	80 (80-81)	72
s4	?	78 (73-88)	77 (73-80)	82 (80-83)	75
S2	88	69 (63-76)	72 (68-76)	73 (70-79)	67
S4	63	52 (45-58)	57 (48-76)	59 (57-62)	49
S5	18	14 (13-15)	16 (14-16)	21 (19-23)	15
r3	75	54 (45-63)	57 (49-62)	55 (55-56)	54
R1	70	57 (50-63)	61 (57-65)	60 (59-62)	58
St1-St1	48	46 (45-50)	Not reported	Not reported	nr
St1-St3	55	54 (53-55)	55 (49-57)	55 (53-56)	nr
St2-St2	60	58 (55-60)	60 (59-62)	53 (50-55)	nr
St3-St3	73	70 (68-73)	Not reported	Not reported	nr
St4-St4	85	72 (63-88)	Not reported	Not reported	nr
St5-St5	58	53 (50-63)	56 (52-60)	53 (51-54)	nr
Lisl	23	28 (23-33)	Not reported	Not reported	nr
Lsiw	3	3	Not reported	Not reported	nr
Sisl	18	13 (10-15)	Not reported	Not reported	nr
Vsl	125	111 (103-120)	115 (94-121)	106 (103-111)	97
vsw ZV2	90	84 (75-90)	86 (80-92)	91 (89-93)	87
vsw anus	78	70 (65-75)	75 (67-83)	75 (73-77)	nr
JV5	73	60 (55-63)	Not reported	Not reported	nr
StIV	75	81 (75-88)	80 (75-87)	68 (68-70)	80-87
Scl	25	20 (17-25)	28 (25-30)	21 (20-21)	30
Scw	5	5	Not reported	Not reported	4
Fdl	23	24 (23-25)	25 (22-27)	22 (21-22)	nr
teeth	?	4	4-5	5	2
Mdl	23	24 (23-25)	24 (22-25)	25 (23-25)	2
teeth	?	2	1-2	2	

Data from this study for Grande Comore, from Kreiter *et al.* (2018) for Martinique, from Moraes *et al.* (2000) for F.C.I (French Caribbean Islands, from various localities of five Islands), from Moraes *et al.* (2004) for Sri Lanka, and from Evans (1952) for the holotype from Indonesia, re-described by Schicha (1975). nr = not reported.

Subfamily Phytoseiinae Berlese

Phytoseiini Berlese, 1913: 3; Phytoseiinae, Vitzthum, 1941: 768.

Genus *Phytoseius* Ribaga

Phytoseius Ribaga, 1904: 177.

Phytoseius amba Pritchard & Baker

Phytoseius (*Pennaseius*) *amba* Pritchard & Baker 1962: 224; Blommers, 1976: 85;
Phytoseius (*Phytoseius*) *amba*, Denmark, 1966: 49;
Typhlodromus (*Pizytoseius*) *amba*, Van der Merwe, 1968: 101;
Phytoseius amba, Swirski & Ragusa, 1978: 408;
Pennaseius amba, Matthysse & Denmark, 1981: 352;
Phytoseius amba, Moraes *et al.*, 1986: 210; 2004: 232; Chant & McMurtry, 2007: 129.

Species of the genus *Phytoseius* are supposed to belong to the Type III species (McMurtry and Croft 1997; McMurtry *et al.* 2013), *i.e.* a polyphagous generalist predator. However, the biology of *Phytoseius amba* remains totally unknown.

Specimens examined — **Md **, INRAPE (long. 11 41'S, lat. 43 14'E, alt. 50 m), 1   on *Annona muricata* L. (Annonaceae), 2-02-2017.

Previous Records — Benin, Burundi, Cameroon, Cape Verde, DR Congo, Kenya, Madagascar Island, Malawi, Mozambique, Nigeria, Reunion Island, Rwanda, Senegal, South Africa, Zambia, Zimbabwe.

Remarks (Table 6) — The setae lengths seem very variable in this species following data of the literature. Measurements of the single adult female (Table 6) agree well with measurements of the literature, especially with those of Ueckermann *et al.* (2007) obtained with a great number of specimens (29) from various countries in Africa, with the exception of longer *j3* and *z3*, shorter *st5-st5*, ventrianal shield length and width at the level of anus and macrosetae of the basitarsus and telotarsus.

Conclusion

Only one species was known before 2017 from the Comoros Archipelago (Schicha 1984). After this preliminary survey conducted in only two locations, the number of species known from Comoros Islands is now of six; four Amblyseinae: *Amblyseius herbicolus*, *Euseius baetae*, *Iphiseius degenerans*, *Neoseiulus longispinosus*; and two Phytoseiinae: *Phytoseius amba* and *Phytoseius mayottae*. No Typhlodrominae were found until now. This is still a low number of species and from a tropical island, one could have expected a higher number, even from only two locations. The low number of species could be explained by the fact that samplings were made in crops more or less disturbed and not in wild areas.

Some of the species collected during this survey have interesting potential for biological control, especially *A. herbicolus*, *I. degenerans* and *N. longispinosus*. This must be underlined as new regulations on importation of macro-organisms are proposed in a lot of countries and specifically for over-sea territories for countries like France that have very far tropical territories. Therefore it is impossible to import and of course to sell and use exotic species if they are not indigenous in the territory. An importation permit must be requested, but it is expensive and chances to obtain are generally very low (Kreiter *et al.* 2016). The knowledge of the biodiversity, especially of efficient biological control agents from oversea territories,

Table 6 Comparisons of character measurements of female specimens of *Phytoseius amba* collected in different locations (Localities followed by the number of specimens measured between brackets)

Characters	Grande Comore (1)	South Africa (11)	Various Countries (29)	Madagascar (6)	Kenya (8)	Holotype
Dsl	270	264-284	282 (264-304)	Not reported	288 (271-312)	283
Dsw	155	130-145	141 (133-149)	Not reported	144 (139-149)	165
j1	23	24-26	22 (19-27)	19-25	24 (22-26)	24
j3	63	46-50	49 (40-56)	50-56	50 (41-60)	42
j4	5	7	4 (3-5)	nr	4 (2-5)	3
j5	5	7	4 (3-5)	nr	4 (2-5)	3
j6	6	7	5 (3-6)	nr	5 (2-7)	4
J2	9	9-11	6 (5-8)	nr	5	6
J5	8	11-14	10 (8-13)	nr	11 (10-12)	11
z2	13	9-11	7 (5-10)	nr	7 (5-10)	5
z3	43	26-29	23 (16-27)	19-31	24 (19-29)	17
z4	10	9-11	8 (5-11)	nr	8 (5-12)	5
z5	5	7	4 (3-5)	nr	5 (2-5)	5
Z4	63	60-80	62 (48-70)	59-65	62 (55-67)	64
Z5	70	57-80	76 (59-86)	64-77	75 (67-82)	83
s4	89	70-85	81 (53-102)	78-82	76 (70-86)	81
s6	75	60-80	77 (48-96)	71-78	78 (67-86)	79
r3	39	37-45	40 (34-46)	37-44	39 (34-43)	41
R1	15	12-16	13 (10-16)	Not reported	14 (12-17)	12
St1-St1	68	50-60	Not reported	Not reported	Not reported	nr
St1-St3	58	68-75	60 (56-64)	Not reported	59 (53-60)	60
St2-St2	63	Not reported	69 (64-72)	Not reported	70 (67-72)	79
St3-St3	68	Not reported	Not reported	Not reported	Not reported	Nr
St4-St4	73	Not reported	Not reported	Not reported	Not reported	Nr
St5-St5	55	67-72	67 (62-74)	Not reported	69 (67-72)	63
Lisl	18	Not reported	Not reported	Not reported	Not reported	nr
Lsiw	2	Not reported	Not reported	Not reported	Not reported	nr
Sisl	8	Not reported	Not reported	Not reported	Not reported	nr
Vsl	83	90-98	99 (86-106)	Not reported	99 (84-108)	100
vsw ZV2	48	48-55	55 (50-64)	Not reported	60 (55-67)	53
vsw anus	38	Not reported	49 (43-56)	Not reported	49 (48-53)	46
JV5	45	45-51	Not reported	54	Not reported	nr
SgeIV	25	24-28	25 (19-32)	23-27	26 (24-34)	21
StiIV	25	30-35	34 (26-40)	30-36	33 (29-38)	31
SbtIV	23	33-38	35 (27-43)	30-34	33 (26-43)	39
StuIV	23	26-30	32 (28-38)	28-32	Not reported	36
Scl	18	16-18	13 (8-19)	Not reported	20 (17-24)	12
Scw	10	4-7	Not reported	Not reported	Not reported	nr
fdl	23	24	25 (23-26)	Not reported	20 (17-24)	17
teeth	1	2	2-3	Not reported	nr	nr
mdl	23	24	25 (23-26)	Not reported	24	17
teeth	1	1	1-2	Not reported	nr	1

Data from this paper for Grande Comore, from Van der Merwe (1968) for South Africa, from Ueckermann *et al.* (2007) for various countries (Burundi 8, Cameroon 1, Ghana 2, Kenya 5, Rwanda 7, Sierra Leone 1, South Africa 4, Democratic Republic of Congo 1) in Africa, from Blommers (1976) for Madagascar, from Moraes *et al.* (1989) for Kenya and measurements of the holotype. nr = not reported.

not only for conversation purposes but also for agricultural and economical ones, is so of a considerable importance.

Acknowledgements

Thanks are due to Mr. Hadji MOUIGNI (from INRAPE) who has helped Jacques Fillâtre (Armefflor) to collect mites in Grande Comore Island and the two anonymous reviewers for improvement of the paper.

References

- Akyazi R., Ueckermann E.A., Soysal M. 2016. The new distribution of *Amblyseius herbicolus* in Turkey (Parasitiformes, Phytoseiidae) with a key of *Amblyseius* species found in Turkey. *Acarologia*, 56(2): 237-244. doi:10.1051/acarologia/20162241
- Athias-Henriot C. 1975. Nouvelles notes sur les Amblyseiini. II. Le relevé organotaxique de la face dorsale adulte (Gamasides, Phytoseiidae). *Acarologia*, 27: 20-29.
- Bhattacharyya S.K. 1968. Two new phytoseiid mites from eastern India (Acarina: Phytoseiidae). *J. Bombay Nat. Hist. Soc.*, 65(3): 677-680.
- Berlese A. 1889. Acari, Myriopoda et Scorpiones hucusque in Italia reperta. *Tipografia Del Seminario*, 6(54): 7-9.
- Berlese A. 1913. Systema Acarorum genera in familiis suis disposita. *Acaroteca Italica*, 1-2: 3-19.
- Berlese A. 1914. Acari nuovi. Manipulus IX. *Redia*, 10: 113-150.
- Berlese A. 1916. Centuria prima di Acari nuovi. *Redia*, 12: 19-66.
- Berlese A. 1921. Acari, Myriopoda et Pseudoscorpiones hucusque in Italia reperta. I. Índice sinonimico dei generi e delle specie illustrate nei fascicoli 1 a 101. *Redia*, 14(1/2): 77-105.
- Bin Ibrahim Y., Tan S.Y. 2000. Influence of sublethal exposure to abamectin on the biological performance of *Neoseiulus longispinosus* (Acari: Phytoseiidae). *J. Econ. Entomol.*, 93(4): 1085-1089. doi:10.1603/0022-0493-93.4.1085
- Blackwood J.S., Schausberger P., Croft B.A. 2001. Prey stage preferences in generalist and specialist phytoseiid mites (Acari: Phytoseiidae) when offered *Tetranychus urticae* (Acari: Tetranychidae) eggs and larvae. *Environ. Entomol.*, 30: 1103-1111. doi:10.1603/0046-225X-30.6.1103
- Blommers L. 1976. Some Phytoseiidae (Acarina: Mesostigmata) from Madagascar, with descriptions of eight new species and notes on their biology. *Bijdragen tot Dierkunde*, 46(1): 80-106.
- Chant D.A. 1959. Phytoseiid mites. Part I. Bionomics of seven species in southeastern England. Part II. A taxonomic review of the family Phytoseiidae, with descriptions of 38 new species. *Can. Entomol.*, 91, suppl. 12: 1-166.
- Chant D.A., McMurtry J.A. 1994. A review of the subfamilies Phytoseiinae and Typhlodrominae. *Intern. J. Acarol.*, 20: 223-310. doi:10.1080/01647959408684022
- Chant D.A., McMurtry J.A. 2003. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): Part I. Neoseiulini new tribe. *Intern. J. Acarol.*, 29(1): 3-46. doi:10.1080/01647950308684319
- Chant D.A., McMurtry J.A. 2004. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): Part III. The tribe Amblyseiini Wainstein, subtribe Amblyseiina n. subtribe. *Intern. J. Acarol.*, 30(3): 171-228. doi:10.1080/01647950408684388
- Chant D.A., McMurtry J.A. 2005. A review of the subfamily Amblyseiinae Muma (Acari: Phytoseiidae): Part V. The tribe Euseiini n. tribe, subtribe Typhlodromalina n. subtribe, Euseiina n. subtribe, and Ricoseiina n. subtribe. *Intern. J. Acarol.*, 31(3): 187-224. doi:10.1080/01647950508684424
- Chant D.A., McMurtry J.A. 2007. Illustrated keys and diagnoses for the genera and sub-genera of the Phytoseiidae of the World. *Indira Publishing House*, West Bloomfield, Michigan, USA, 220 pp.
- Chant D.A., Yoshida Shaul E. 1991. Adult ventral setal patterns in the family Phytoseiidae (Acari: Gamasina). *Intern. J. Acarol.*, 17: 187-199. doi:10.1080/01647959108683906
- Chaudhri W.M. 1968. Six new species of mites of the genus *Amblyseius* (Phytoseiidae) from Pakistan. *Acarologia*, 10: 550-562.
- Corpuz L.A., Rimando L. 1966. Some Philippine Amblyseiinae (Phytoseiidae: Acarina). *Philippine Agriculture*, 50: 114-136.
- Croft B.A., Luh H.-K., Schausberger P. 1999a. Larval size relative to larval feeding, cannibalism of larvae, egg, or adult female size and larval-adult setal patterns among thirteen phytoseiid mite species. *Exp. Appl. Acarol.*, 23: 599-610. doi:10.1023/A:1006236310613
- Croft B.A., McMurtry J.A., Luh H.-K. 1999b. Do literature citation frequencies for six prey-food groups reflect feeding specialization and preferences among for Phytoseiid predation types?. *Exp. Appl. Acarol.*, 23: 551-565. doi:10.1023/A:1006236310613
- Daneshvar H., Denmark H.A. 1982. Phytoseiids of Iran (Acarina: Phytoseiidae). *Intern. J. Acarol.*, 8(1): 3-14. doi:10.1080/01647958208683272
- De Leon D. 1967. Some mites of the Caribbean Area. Part I. Acarina on Plants in Trinidad, West Indies. *Allen Press Inc.*, Lawrence, Kansas, 66 pp.
- Demite P.R., Moraes G.J. de, McMurtry J.A., Denmark H.A., Castilho R.C. 2017. Phytoseiidae Database. Available from: www.lea.esalq.usp.br/phytoseiidae

- Denmark H.A. 1966. Revision of the genus *Phytoseius* Ribaga, 1904 (Acarina: Phytoseiidae). Fla Dep. Agric. Bull., 6: 1-105.
- Denmark H.A., Muma M.H. 1989. A revision of the genus *Amblyseius* Berlese, 1914 (Acari: Phytoseiidae). Occasional Papers of the Florida State Collection of Arthropods, USA, 4, 149 pp.
- Duarte M.V.A., Venzon M., Bittencourt M.C.de S., Rodriguez-Cruz F.A., Pallini A., Janssen A. 2015. Alternative food promotes broad mite control on chilli pepper plants. BioControl, 60: 817-825. doi:10.1007/s10526-015-9688-x
- Ehara S. 1958. Three predatory mites of the genus *Typhlodromus* from Japan (Phytoseiidae). Annot. Zool. Japonenses, 31: 53-57.
- El-Badry E.A. 1968. Three new species of phytoseiid mites from western Sudan. Rev. Zool. Bot. Afric., 77: 321-328.
- El-Banhawy E.M., Knapp M. 2011 Mites of the family Phytoseiidae Berlese from Kenya (Acari: Mesostigmata). Zootaxa, 2945: 1-176.
- Evans G.O. 1952. On a new predatory mite of economic importance. Bull. Entomol. Res., 43: 397-401. doi:10.1017/S0007485300040566
- Evans G.O. 1953. On some mites of the genus *Typhlodromus* Scheuten, 1857, from S. E. Asia. Ann. Mag. Nat. Hist., 6: 449-467. doi:10.1080/00222935308654444
- Evans G.O. 1954. The genus *Iphiseius* Berlese (Acarina: Laelaptidae). Proceed. Zool. Soc., 124: 517-526. doi:10.1111/j.1469-7998.1954.tb07793.x
- Fantinou A., Baxevas A., Drizou F., Labropoulos P., Perdakis D., Papadoulis G. 2012. Consumption rate, functional response and preference of the predaceous mite *Iphiseius degenerans* to *Tetranychus urticae* and *Eutetranychus orientalis*. Exp. Appl. Acarol., 58:133-144. doi:10.1007/s10493-012-9557-6
- Gupta S.K. 1981. On a collection of Phytoseiidae (Acari: Mesostigmata) from Himachal Pradesh (India), with descriptions of two new species. Ind. J. Acarol., 5: 32-36.
- Gupta S.K. 1986. Fauna of India (Acari: Mesostigmata) Family Phytoseiidae. Zoological Survey of India, Calcutta, India, 350 pp.
- Hirschmann W. 1962. Gangsystematik der Parasitiformes. Acarologie Schriftenreihe für Vergleichende Milbenkunde, Hirschmann-Verlag, Furth/Bay, 5(5-6), 80 pp.+ 32 plates.
- Hughes A.M. 1948. The mites associated with stored food products. Ministry of Agriculture and Fisheries, H. M. Stationary Office, London, 168 pp.
- Huyen L.T., Tung N.D., Lan D.H., Chi C.V., De Clercq P., Dinh N.V. 2017. Life table parameters and development of *Neoseiulus longispinosus* (Acari: Phytoseiidae) reared on citrus red mite, *Panonychus citri* (Acari: Tetranychidae) at different temperatures. Syst. Appl. Acarol., 22(9): 1316-1326. doi:10.11158/saa.22.9.3
- Kade N., Gueye-Ndiaye A., Duverney C., Moraes G.J. de 2011. Phytoseiid mites (Acari: Phytoseiidae) from Senegal. Acarologia, 51(1): 133-138. doi:10.1051/acarologia/20112001
- Karg W. 1991. Die Raubmilbenarten der Phytoseiidae Berlese (Acarina) Mitteleuropas sowie angrenzender Gebiete. Zoologische Jahrbucher Systematik, 118(1): 1-64.
- Kreiter S., Vicente V., Tixier M.-S., Fontaine O. 2016. An unexpected occurrence of *Amblyseius swirskii* Athias-Henriot in La Réunion Island (Acari: Phytoseiidae). Acarologia, 56(2): 175-181. doi:10.1051/acarologia/20162254
- Kreiter S., Mailloux J., Tixier M.-S., Le Bellec F., Douin M., Guichou S., Etienne J. 2013. New phytoseiid mites of the French West Indies, with description of a new species, and new records (Acari: Mesostigmata). Acarologia, 53(3): 285-303 doi:10.1051/acarologia/20132095
- Kreiter S., Zriki G., Ryckewaert P., Pancarte C., Douin M., Tixier M.-S. 2018. Phytoseiid mites of Martinique, with redescription of four species and new records (Acari: Mesostigmata). Acarologia, 58(2): 366-407. doi:10.24349/acarologia/20184248
- Lindquist E., Evans G.W. 1965. Taxonomic concepts in the Ascidae, with a modified setal nomenclature for the idiosoma of the Gamasina Acarina: Mesostigmata. Mem. Entomol. Soc. Can., 47: 1-64. doi:10.4039/entm9747fv
- Mailloux J., Le Bellec F., Kreiter S., Tixier M.-S., Dubois P. 2010. Influence of ground cover management on diversity and density of phytoseiid mites (Acari: Phytoseiidae) in Guadeloupean citrus orchards. Exp. Appl. Acarol., 52: 275-290. doi:10.1007/s10493-010-9367-7
- Matthysse J.G., Denmark H.A. 1981. Some phytoseiids of Nigeria (Acarina: Mesostigmata). The Fa Entomol., 64: 340-357. doi:10.2307/3494585
- Meyer M.K.P., Rodrigues M.C. 1966. Acari associated with cotton in southern Africa (with reference to other plants). Garcia de Orta, Revista Junta Investigacoes de Ultramar, 13(2): 195-226.
- McMurtry J.A., Croft B.A. 1997. Life-styles of phytoseiid mites and their roles in biological control. Ann. Rev. Entomol., 42: 291-321. doi:10.1146/annurev.ento.42.1.291
- McMurtry J.A., Moraes G.J. de, Sourasso N.F. 2013. Revision of the life styles of phytoseiid mites (Acari: Phytoseiidae) and implications for biological control strategies. Syst. Appl. Acarol., 18: 297-320. doi:10.11158/saa.18.4.1
- Moraes G.J. de, Kreiter S., Lofego A.C. 2000. Plant mites of the French Antilles. 3. Phytoseiidae. Acarologia, 40: 237-264.
- Moraes G.J. de, Lopes P.C., Fernando C.P. 2004a. Phytoseiid mite (Acari: Phytoseiidae) of coconut growing areas in Sri Lanka, with descriptions of three new species. J. Acarol. Soc. Jap., 13(2): 141-160. doi:10.2300/acari.13.141
- Moraes G.J. de, McMurtry J.A., Denmark H.A. 1986. A catalog of the mite family Phytoseiidae. References to taxonomy, synonymy, distribution and habitat. EMBRAPA – DDT, Brasília, Brazil, 353 pp.
- Moraes G.J. de, McMurtry J.A., Denmark H.A., Campos C.B. 2004b. A revised catalog of the mite family Phytoseiidae. Zootaxa, 434: 1-494. doi:10.11646/zootaxa.434.1.1

- Moraes G.J. de, McMurtry J.A., van den Berg H., Yaninek J.S. 1989. Phytoseiid mites (Acari: Phytoseiidae) of Kenya, with descriptions of five new species and complementary descriptions of eight species. *Intern. J. Acarol.*, 15(2): 79-93. doi:10.1080/01647958908683829
- Moraes G.J. de, Oliveira A.R., Zannou I.D. 2001a. New phytoseiid mites (Acari: Phytoseiidae) from tropical Africa. *Zootaxa*, 8: 1-10. doi:10.11646/zootaxa.8.1.1
- Moraes G.J. de, Ueckermann E.A., Oliveira A.R., Yaninek J.S. 2001b. Phytoseiidae mites of the genus *Euseius* (Acari: Phytoseiidae) from Sub-Saharan Africa. *Zootaxa*, 3: 1-70.
- Moraes G.J. de, Zannou I.D., Ueckermann E.A., Oliveira A.R., Yaninek J.S., Hanna R. 2007. Phytoseiid mites of the tribes Afroseiulini, Kampimodromini and Phytoseiulini, and complementary notes on mites of the tribes Euseiini and Neoseiulini (Acari: Phytoseiidae) from sub-Saharan Africa. *Zootaxa*, 1628: 1-22.
- Muma M.H. 1961. Mites associated with citrus in Florida. *Fla Agric. Exp. Stat. Bull.*, 640: 1-39.
- Muma M.H., Denmark H.A. 1970. Phytoseiidae of Florida. Arthropods of Florida and neighboring land areas, 6. Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville, USA, 150 pp.
- Myers N. 1988. Threatened biotas: hostspots in tropical forests. *Environmentalist*, 8: 187-208. doi:10.1007/BF02240252
- Myers N., Mittermeier R.A., Mittermeier C.G., Da Fonseca G.A., Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature*, 403: 853-858. doi:10.1038/35002501
- Northcraft P.D. 1987. First record of three indigenous predacious mites in Zimbabwe. *J. Entomol. Soc. S. Afric.*, 50(2): 521-522.
- Nusartlert N., Vichitbandha P., Baker G., Chandrapatya A. 2011. Pesticide-induced mortality and prey-dependent life history of the predatory mite *Neoseiulus longispinosus* (Acari: Phytoseiidae). *Trends in Acarology*. Netherlands, Springer: 495-498.
- Nwilene F.E., Nachman G. 1996. Functional responses of *Iphiseius degenerans* and *Neoseiulus teke* (Acari: Phytoseiidae) to changes in the density of the cassava green mite, *Mononychellus tanajoa* (Acari: Tetranychidae). *Exp. Appl. Acarol.*, 20: 259-271. doi:10.1007/BF00052876
- Papadoulis G.Th., Emmanouel N.G. 1991. The genus *Amblyseius* (Acari: Phytoseiidae) in Greece, with the description of a new species. *Entomologia Hellenica*, 9: 35-62. doi:10.12681/eh.13990
- Pritchard A.E., Baker E.W. 1962. Mites of the family Phytoseiidae from Central Africa, with remarks on genera of the world. *Hilgardia*, 33: 205-309. doi:10.3733/hilg.v33n07p205
- Rahman V.J., Babu A., Roobakkumar A., Perumalsamy K. 2013. Life table and predation of *Neoseiulus longispinosus* (Acari: Phytoseiidae) on *Oligonychus coffeae* (Acari: Tetranychidae) infesting tea. *Exp. Appl. Acarol.*, 60: 229-240. doi:10.1007/s10493-012-9649-3
- Reis P.R., Teodoro A.V., Pedro Neto M., Da Silva E.A. 2007. Life history of *Amblyseius herbicolus* (Chant) (Acari: Phytoseiidae) on coffee plants. *Neotrop. Entomol.*, 36(2): 282-287. doi:10.1590/S1519-566X2007000200016
- Ribaga C. 1904 (1902). Gamasidi planticoli. *Riv. Patol. Veg.*, 10: 175-178.
- Rodriguez-Cruz F.A., Venzon M., Pinto C.M.F. 2013. Performance of *Amblyseius herbicolus* on broad mites and on castor bean and sunnhemp pollen. *Exp. Appl. Acarol.*, 60: 497-507. doi:10.1007/s10493-013-9665-y
- Rowell H.J., Chant D.A., Hansell R.I.C. 1978. The determination of setal homologies and setal patterns of the dorsal shield in the family Phytoseiidae. *Can. Entomol.*, 110: 859-876. doi:10.4039/Ent110859-8
- Schausberger P., Croft B.A. 1999a. Predation on and discrimination between con- and heterospecific eggs among specialist and generalist phytoseiid mites (Acari: Phytoseiidae). *Environ. Entomol.*, 28: 523-528. doi:10.1093/ee/28.3.523
- Schausberger P., Croft B.A. 1999b. Activity, feeding, and development among larvae of specialist and generalist phytoseiid mite species (Acari: Phytoseiidae). *Biol. Control*, 28: 322-329. doi:10.1093/ee/28.2.322
- Schausberger P., Croft B.A. 2000a. Nutritional benefits of intraguild predation and cannibalism among generalist and specialist phytoseiid mites. *Ecol. Entomol.*, 25: 1-8. doi:10.1046/j.1365-2311.2000.00284.x
- Schausberger P., Croft B.A. 2000b. Cannibalism and intraguild predation among phytoseiid mites: are aggressiveness and prey preference related to diet specialization? *Exp. Appl. Acarol.*, 24: 709-725. doi:10.1023/A:1010747208519
- Schicha E. 1975. Predacious mites (Acarina: Phytoseiidae) on sprayed apple trees at Bathurst (N.S.W.). *J. Austral. Entomol. Soc.* 14: 217-219. doi:10.1111/j.1440-6055.1975.tb02029.x
- Schicha E. 1984. Contribution to the knowledge of the genus *Phytoseius* Ribaga in Australia, the South Pacific and Indian Ocean regions with four new species and records of known species (Acarina: Phytoseiidae). *Intern. J. Acarol.*, 10(2): 117-128. doi:10.1080/01647958408683361
- Swirski E., Ragusa S. 1978. Three new species of phytoseiid mites from Kenya (Mesostigmata: Phytoseiidae). *Zool. J. Linnean Soc.*, 63: 397-409. doi:10.1111/j.1096-3642.1978.tb02101.x
- Thongtab T., Chandrapatya A., Baker G. 2001. Biology and efficacy of the predatory mite, *Amblyseius longispinosus* (Evans) (Acari, Phytoseiidae) as a biological control agent of *Eotetranychus cendanai* Rimando (Acari, Tetranychidae). *J. Appl. Entomol.*, 125: 543-549. doi:10.1046/j.1439-0418.2001.00583.x
- Ueckermann E.A., Loots G.C. 1988. The African species of the subgenera *Anthoseius* De Leon and *Amblyseius* Berlese (Acari: Phytoseiidae). *Entomol. Mem. Depart. Agric. Water Supply, RSA*, 73: 1-168.
- Ueckermann E.A., Zannou I.D., Moraes G.J. de, Oliveira A.R. de, Hanna R., Yaninek J.S. 2007. Phytoseiid mites of the subfamily Phytoseiinae (Acari: Phytoseiidae) from sub-Saharan Africa. *Zootaxa*, 1658, 1-20.
- Van der Merwe G.G. 1968. A taxonomic study of the family Phytoseiidae (Acari) in South Africa with contributions to the biology of two species. *Entomol. Mem.*, South Africa Department of Agricultural Technical Services, 18: 1-198.

- Vantornhout I., Minnaert H.L., Tirry L., Clercq P. 2005. Influence of diet on life table parameters of *Iphiseius degenerans* (Acari: Phytoseiidae). *Exp. and Appl. Acarol.*, 35: 183-195. doi:10.1007/s10493-004-3940-x
- Vitzthum H. von 1941. Acarina. In: Bronns, H.G. (Ed.), *Klassen und Ordnungen des Tierreichs* 5, Akademischer Verlag, Leipzig, Germany, 764-767.
- Wainstein B.A. 1962. Révision du genre *Typhlodromus* Scheuten, 1857 et systématique de la famille des Phytoseiidae (Berlese 1916) (Acarina: Parasitiformes). *Acarologia*, 4: 5-30.
- Womersley H. 1954. Species of the subfamily Phytoseiinae (Acarina: Laelaptidae) from Australia. *Austral. J. Zool.*, 2: 169-191. doi:10.1071/ZO9540169
- Zaher M. A. 1986. Survey and ecological studies on phytophagous, predacious and soil mites in Egypt. 11 A: Predacious and non phytophagous mites (Nile Valley and Delta). *Fac. Agric., Cairo Univ.*, 567 pp.
- Zannou I.D., Hanna R., Moraes G.J. de, Kreiter S., Phiri G., Jone A. 2005. Mites of cassava (*Manihot esculenta* Crantz) habitats in Southern Africa. *Intern. J. Acarol.*, 31(2): 149-164. doi:10.1080/01647950508683667
- Zannou I.D., Moraes G.J. de, Ueckermann E.A., Oliveira A.R., Yaninek J.S., Hanna R. 2007. Phytoseiid mites of the subtribe Amblyseiina (Acari: Phytoseiidae: Amblyseiini) from sub-Saharan Africa. *Zootaxa*, 1550: 1-47.
- Zappalà L., Kreiter S., Russo A., Tropea Garzia G., Auger P. 2015. First record of the Persea Mite *Oligonychus perseae* (Acari: Tetranychidae) in Italy with a review of the literature. *Intern. J. Acarol.*, 41(2): 1-3. doi:10.1080/01647954.2015.1014415
- Zhang Y., Lin J., Wen Z., Mao D., Chi Y. 1996. Effect of temperature on life history and predatory functional response of *Amblyseius longispinosus* (Acari: Phytoseiidae). *Fujian J. Agric. Sci.*, 11: 41-44.
- Zhang Y., Zhang Z.-Q., Ji J., Lin J. 1999. Predation of *Amblyseius longispinosus* (Acari: Phytoseiidae) on *Schizotetranychus nanjingensis* (Acari: Tetranychidae), a spider mite injurious to bamboo in Fujian, China. *Syst. Appl. Acarol.*, 4: 63-68. doi:10.11158/saa.4.1.9