Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Supporting agricultural research
for sustainable development

**Acarologia is under free license and distributed under the terms of the
Creative Commons-BY**

Genus *Eryngiopus* Summers (Acari: Trombidiformes: Stigmaeidae) from Saudi Arabia; a new record and redescription of *E. discus* Meyer, with a key to the world species

Muneeb Ur Rehman^a, Muhammad Kamran^a, Fahad J. Alatawi^a

^a Acarology Laboratory, Department of Plant Protection, College of Food & Agriculture Sciences, King Saud University, 11451, P.O. Box 2460, Riyadh, Saudi Arabia.

ABSTRACT

Eryngiopus discus Meyer, 1969 (Acari: Trombidiformes: Stigmaeidae), the first species of the genus *Eryngiopus* Summers recorded from Saudi Arabia, is re-described and illustrated. The specimens of this species were collected from foxtail grass *Cenchrus* sp. (Poaceae) and *Euryops arabicus* (Asteraceae). Also, the species of the genus *Eryngiopus* are categorized into two species groups, *bakeri* and *gracilis*, based on three and four pairs of prodorsal setae, respectively. A key to the world species of the genus *Eryngiopus* is provided too.

Keywords *bakeri*, *gracilis*, predatory mites, species groups, stigmaeids

Zoobank <http://zoobank.org/F75FB617-F28C-4516-ACC9-9F212FD054DE>

Introduction

The family Stigmaeidae (Acari: Trombidiformes: Prostigmata) contains 32 genera (Fan *et al.*, 2016, Rehman *et al.*, 2016) of which only five genera including *Agistemus* Summers, *Cheyllostigmaeus* Willmann, *Eustigmaeus* Berlese, *Eryngiopus* Summers and *Ledermuelleriopsis* Willmann have been reported from Saudi Arabia so far (Alatawi, 2011a & b, Alatawi & Kamran, 2017).

The species of the genus *Eryngiopus* inhabit plant foliage, soil litter, lichens and mosses (Ueckermann & Meyer, 1987, Mاتيولي *et al.*, 2002). Members of this genus are considered as predators of soft-bodied insects and mite pests (Fan & Zhang, 2005, Khanjani *et al.*, 2014) and known from different geographical regions of the world (Vacante & Gerson, 1988, Fan & Zhang, 2005).

Up to now, 31 valid species from the genus are known worldwide (Doğan *et al.*, 2015, Fan *et al.*, 2016). Previously, few regional keys have been made for the afrotropical species of the genus (Meyer, 1969, Ueckermann & Meyer, 1987, Van Dis & Ueckermann, 1993). A key to the all known *Eryngiopus* species was provided by Vacante & Gerson (1988).

The objectives of this study were to re-describe and illustrate *E. discus* Meyer, since the original description lacks complete details of the species (Meyer, 1969) which is the requirement of modern taxonomy, and to provide a key to the world species of *Eryngiopus*.

Materials and methods

Mite specimens were collected by shaking the plant foliage on a white paper with a fine tip camel hairbrush and preserved in 70% ethanol. Samples of soil and leaf debris were

Received 27 November 2017

Accepted 12 March 2018

Published 01 June 2018

Corresponding author

Fahad J. Alatawi:

falatawi@ksu.edu.sa

Academic editor

Farid Faraji

DOI

10.24349/acarologia/20184262

© Copyright

Rehman M.U. *et al.*

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

How to cite this article Rehman M.U. *et al.* (2018), Genus *Eryngiopus* Summers (Acari: Trombidiformes: Stigmaeidae) from Saudi Arabia; a new record and redescription of *E. discus* Meyer, with a key to the world species. *Acarologia* 58(3): 655-664; DOI 10.24349/acarologia/20184262

also collected from the field and were processed through Tullgren's funnels. The preserved specimens were then mounted on glass slides in Hoyer's medium under a stereomicroscope (SZX10, Olympus). The mounted specimens were dried in an oven at 40°C for one week. The specimens were identified under a phase contrast microscope (BX51, Olympus) with the help of literature and diagnostic keys. Different body parts were pictured using an Auto-Montage system (Syncroscopy, Cambridge) and illustrated with Adobe illustrator (Adobe Systems Incorporated). All measurements are given in micrometers (µm). Body width was measured at the broadest area and setae were measured from the setal base to the tip. Terminology and setal nomenclature follow those of Kethley (1990) and Grandjean (1944). All specimens have been deposited in the Acarology Research Laboratory, Department of Plant Protection, College of Food and Agriculture Science, King Saud University, Riyadh, Kingdom of Saudi Arabia.

Results and discussion

Family Stigmaeidae Oudemans

Genus: *Eryngiopus* Summers

Eryngiopus Summers, 1964: 186.

Type species: *Eryngiopus gracilis* Summers, 1964.

Diagnosis based on Fan & Zhang (2005).

In the present study, the species of genus *Eryngiopus* are categorized into two species groups based on the number of prodorsal setae (Table 1); 1) species group *bakeri* having three pairs of prodorsal setae (*vi*, *ve* and *sci*) including nine species reported from Turkey, Egypt, Yemen, Thailand, China, South Africa and USA; 2) species group *gracilis* having four pairs of prodorsal setae (*vi*, *ve*, *sci* and *sce*) comprising 22 species mostly reported from USA, New Zealand, Italy, South Africa, China, Russia, Uzbekistan, Iran, Pakistan and India (Doğan *et al.*, 2015, Fan *et al.*, 2016).

Number of prodorsal setae, either 3 pairs (*vi*, *ve*, *sci*) or four pairs (*vi*, *ve*, *sci*, *sce*) (Table 1), is strong and constant morphological character to categorize the species of *Eryngiopus* into two species groups. Vacante & Gerson (1988) used number of prodorsal setae (three/four) pairs as an important and basic character to develop a key to 17 world species of *Eryngiopus*. Also, Van Dis & Ueckermann (1993) separated the Afrotropical *Eryngiopus* species and used number of prodorsal setae (three/four) pairs as the first couplet in the key.

Species group *bakeri*

Eryngiopus discus Meyer

Eryngiopus discus Meyer, 1969: 232.

Re-description of adult female (n= 8) (Figures 1-4)

Idiosoma oval — Length of body (excluding gnathosoma) 370 – 378; length of gnathosoma 75 – 80; width of idiosoma 170 – 175 at the level of setae *c2*.

Gnathosoma (Figures 1C, 3) — Chelicerae separated; palp five segmented; palptibial claw subequal to palptarsus; palptarsus with tridentate terminal eupathidion distally, one solenidion, one eupathidion and four simple setae; palptibia with one well developed claw and one seta-like accessory claw, and two simple setae; palpgenua with one serrate seta; palpfemur with one simple and 2 serrated setae. Subcapitulum with 2 pairs of subcapitular setae (*m*, *n*) and two pairs of adoral setae (*or1*, *or2*). Length of subcapitular setae as follows: *m* 55 – 57, *n* 29 – 31, *or1* 12 – 14, *or2* 14 – 17; distances: *or1-or1* 6 – 8, *or2-or2* 10 – 12, *or1-or2* 5 – 7, *m-m* 25 – 27, *n-n* 30 – 32, *or1-m* 37 – 40, *or2-n* 40 – 44, *or1-m* 35 – 38, *m-n* 6 – 7.

Table 1 Some morphological diagnostic characters of the species of the genus *Eryngiopus*.

Species	AG	TPS	PDS	Coxae	Troch	Femur	Genu	Tibiae	tarsus	1a	3a	4a	References
Species group <i>bakeri</i> (Three pairs of prodorsal setae; <i>vi</i> , <i>ve</i> , <i>sci</i>)													
<i>anatolicus</i>	3	-	U	2-1-2-1	1-1-1-1	6-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	22	24	17	Dogan <i>et al.</i> , 2015
<i>bakeri</i>	3	F	D	2-1-2-2	1-1-1-1	4-4-2-2	4-1-0-0	6-6-6-6	14-10-8-7	24	24	20	Vacante & Gerson, 1988
<i>custodis</i>	2	Tr	S	2-1-2-2	1-1-1-1	4-4-2-1	2-0-0-0	6-6-6-6	11-9-7-6	19	19	19	Van Dis & Ueckermann, 1993
<i>discus</i>	3	Tr	D	2-1-2-2	1-1-1-1	4-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	80	21	20	Meyer, 1969; Ueckermann & Meyer, 1987; present study
<i>jiangxiensis</i>	3	Sp	U	2-1-2-1	1-1-1-1	4-4-2-2	3-1-0-0	6-6-6-6	14-10-8-8	82	95	88	Hu & Chen, 1992
<i>nanchangensis</i>	3	-	U	2-1-2-1	1-1-1-1	4-4-2-2	3-1-0-0	6-6-6-6	14-9-8-7	80	60	17	Hu & Chen, 1994
<i>woodi</i>	3	F	D	2-1-2-2	1-1-1-1	4-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	103	37	30	Gomaa & El-Enany, 1985
<i>yasumatsui</i>	3	F	D	-	-	4-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	-	-	-	Ehara & Wongsiri, 1984
<i>yemenensis</i>	3	S	S	2-1-2-2	1-1-1-1	5-4-2-2	4-1-0-0	6-6-6-6	13-9-7-7	L	L	L	Van Dis & Ueckermann, 1993
Species group <i>gracilis</i> (Four pairs of prodorsal setae; <i>vi</i> , <i>ve</i> , <i>sci</i> , <i>sce</i>)													
<i>affinis</i>	3	Tr	U	2-1-2-1	1-1-1-1	6-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	25	28	21	Jafari <i>et al.</i> , 2014
<i>arboreus</i>	2	S	D	2-1-2-1	1-1-1-0	4-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	36	42	34	Vacante & Gerson, 1988; Fan & Zhang, 2005
<i>audreae</i>	3	S	D	2-1-2-1	1-1-1-0	4-4-2-2	3-1-0-0	6-6-6-6	14-9-7-7	59	62	17	Maake <i>et al.</i> , 2015
<i>bidens</i>	2	F	D	2-1-2-1	1-1-1-0	4-4-2-2	3-1-0-0	6-6-6-6	14-10-8-8	L	L	L	Meyer, 1969; Vacante & Gerson, 1988
<i>bifidus</i>	2	F	U	2-1-2-1	1-1-1-0	4-4-2-2	4-0-0-0	6-6-6-6	14-10-8-8	85	40	65	Vacante & Gerson, 1988; Fan & Zhang, 2005
<i>cirrus</i>	3	Tr	U	2-2-2-1	1-1-1-1	4-4-2-2	3-2-0-0	6-5-5-6	14-9-8-8	L	S	S	Chaudhri, 1979
<i>coimbatorensis</i>	3	S	D	-	-	3-3-2-2	1-1-1-1	-	-	-	-	-	Gupta & David, 1990
<i>dicrotrichus</i>	3	F	D	2-1-2-1	1-1-1-0	4-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	69	67	15	Fan <i>et al.</i> , 2000
<i>gracilis</i>	3	S	U	2-1-2-1	1-1-1-1	4-4-2-2	4-0-0-0	6-6-6-4	14-9-7-7	20	34	27	Summers, 1964; Vacante & Gerson, 1988
<i>hamedanicus</i>	3	S	D	2-1-2-1	1-1-1-1	6-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	28	28	22	Khanjani <i>et al.</i> , 2014
<i>hortus</i>	3	-	S	2-2-2-2	1-1-1-1	4-4-2-2	3-1-0-0	5-6-5-5	11-8-7-7	L	S	S	Chaudhri, 1979
<i>langroudiensis</i>	3	S	S	2-1-2-1	1-1-1-1	3-3-2-2	3-0-0-0	6-6-6-6	12-10-8-8	60	44	40	Khanjani <i>et al.</i> , 2014
<i>lindei</i>	3	F	D	2-1-2-1	-	4-4-2-2	3-1-0-0	6-6-6-6	14-10-8-8	L	L	L	Meyer, 1969; Vacante & Gerson, 1988
<i>microsetus</i>	2	S	D	2-1-2-1	1-1-1-0	4-4-2-2	4-0-0-0	6-6-6-6	14-10-8-8	72	40	45	Summers, 1964; Vacante & Gerson, 1988
<i>nelsonensis</i>	3	S	U	2-1-2-1	1-1-1-1	5-4-2-2	4-2-0-0	6-6-6-6	14-9-7-7	18	37	18	Vacante & Gerson, 1988; Fan & Zhang, 2005
<i>parsimilis</i>	2	F	S	2-1-2-1	1-1-1-0	4-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	25	25	32	Ueckermann & Smith-Meyer, 1987
<i>placidus</i>	3	F	D	2-1-2-1	1-1-1-0	4-4-2-2	4-0-0-0	6-6-6-6	14-10-8-8	70	60	40	Kuznetsov, 1977
<i>siculus</i>	3	F	D	2-1-2-1	1-1-1-0	4-4-2-2	4-0-0-0	6-6-6-6	14-10-8-8	86	80	45	Vacante & Gerson, 1988
<i>similis</i>	2	F	D	2-1-2-1	1-1-1-0	4-4-2-2	4-1-0-0	6-6-6-6	14-10-8-8	24	24	30	Vacante & Gerson, 1988; Fan & Zhang, 2005
<i>summersi</i>	3	F	U	2-1-2-1	1-1-1-1	4-4-2-2	4-1-0-0	6-6-6-6	14-9-8-8	75	70	80	Vacante & Gerson, 1988
<i>tauricus</i>	3	S	D	1-1-2-1	1-1-1-1	4-4-2-2	4-1-0-0	6-6-6-6	14-9-8-8	28	30	25	Vacante & Gerson, 1988; Kuznetsov, 1978
<i>vagantis</i>	3	S	U	1-1-2-1	1-1-1-1	5-4-3-2	4-1-0-0	6-6-6-6	14-9-8-8	30	80	64	Summers, 1964; Vacante & Gerson, 1988

AG, Aggenital setae; TPS, shape of terminal palpal sensillum either F (forked), S (simple), Tr (tri dentate) or Sp (spine like); PDS, Propodosomal dorsal shields either D (divided), U (undivided) or S (striated); PRS, Propodosoma with either three or four pairs of setae. Setal count on leg segments includes solenidia and micro setae.

Dorsum (Figures 1A, 2A-B) — Prodorsal shield divided medially by striations, bearing three pairs of setae (*vi*, *ve*, *sci*), setae *sce* absent, eyes present, 8 – 11 µm, post ocular bodies (*pob*) absent; hysterosomal striations with micro tubercles (Fig.1B), longitudinal between setae *cl* and *el*, and transverse posterior to *el*. Setae *fl* situated on small platelets. Suranal shield divided into three feebly developed plates bearing setae *h1* and *h2*; setae *h1* share larger median plate; setae *h2* on smaller lateral plates. Hysterosoma bears eight pairs of setae. Dorsal idiosomal setae slender, slightly barbed and tapering towards end. Length of dorsal setae as follows: *vi* 20 – 23, *ve* 28 – 32, *sci* 18 – 21, *cl* 12 – 14, *c2* 23 – 25, *d1* 11 – 14, *d2* 12 – 14, *el* 17 – 20, *e2* 14 – 17, *fl* 19 – 22, *h1* 27 – 31, *h2* 31 – 35. Distances between dorsal setae: *vi-vi* 25 – 27, *ve-ve* 48-53, *vi-ve* 19 – 21, *sci-sci* 77 – 83, *ve-sci* 31 – 34, *cl-cl* 62 – 66, *cl-c2* 68 – 71, *c2-c2* 124 – 127, *cl-d1* 73 – 75, *d1-d1* 93 – 97, *d1-d2* 41 – 43, *d1-el* 44 – 48, *d1-e2* 26 – 30, *d2-e2* 50 – 55, *d2-d2* 133 – 138, *el-el* 41 – 44, *e2-e2* 51 – 55, *el-e2* 45 – 48, *el-fl* 35 – 38, *fl-fl* 76 – 81, *f-h1* 41 – 46, *h1-h1* 31 – 25, *fl-h2* 42 – 45, *h2-h2* 75 – 80, *h1-h2* 29 – 32; ratios: *vi/vi* 0.76, *cl/cl-cl* 0.2, *d1/d1-d1* 0.12, *el/el-el* 0.41, *fl/fl-fl* 0.26, *h1/h1-h1* 1.27, *h2/h2-h2* 1.10, *h1/h2* 0.88.

Venter (Figure 1C, 2C) — The first pair of intercoxal setae *1a* ultra long and reaches to the bases of subcapitular setae *m*. The latter intercoxal setae *3a* and *4a* much shorter than first pair of intercoxals. Lengths of setae; *1a* 80 – 86, *1b* 16 – 18, *1c* 13 – 14, *2c* 15 – 17, *3a* 21 – 23,

Figure 1 *Eryngiopus discus* Meyer, adult female: A – Dorsal view of idiosoma; B – Micro tubercles on dorsal striations; C – Ventral view of idiosoma.

Figure 2 *Eryngiopus discus* Meyer, adult female: A – Prodorsum; B – Opisthosoma; C – Genital region.

3b 20 – 22, 3c 17 – 20, 4a 20 – 23, 4b 17 – 20 and 4c 15 – 17, aggenital setae three pairs: *ag1* 16 – 17, *ag2* 23 – 25, *ag3* 19 – 20, ano-genital area with one genital setae, *g* and three pairs of pseudanal setae (*ps1*-3): *g* 16 – 18, *ps1* 19 – 22, *ps2* 23 – 25, *ps3* 24 – 27. Distances between setae: 1a-1a 26 – 28, 3a-3a 56 – 58, 4a-4a 35 – 37, *ag1*-*ag1* 26 – 28, *ag2*-*ag2* 38 – 40, *ag3*-*ag3* 49 – 52, *ag1*-*ag2* 20 – 21, *ag2*-*ag3* 20 – 21, *ps1*-*ps1* 32 – 34, *ps2*-*ps2* 35 – 38, *ps3*-*ps3* 20 – 21; *g1*-*g1* 6 – 7.

Legs (Figure 4) — Length of legs I-IV, (tarsal claws to coxal bases): 188-192, 154 – 159, 148 – 155, 153 – 159 respectively. Empodial shaft branches up into 2 pairs of tenent hairs before pass beyond tips of claws. Number of setae on leg segments as follows: coxae 2-1-2-2; trochanters 1-1-1-1; femora 4-4-2-2; genua 3+1κ-1-0-0; tibiae 5+1φ-5+1φ-5+1φ-5+1φ; tarsi 13+1ω-9+1ω-7+1ω-7+1ω.

Male and immature stages — unknown

Material examined — Five females, *Euryops arabicus* (Asteraceae), Taif, 21.19385°N, 040.25704°E, October 11, 2016, coll. M. Ur Rehman, Muhammad Kamran; three females, foxtail grass *Cenchrus* sp. (Poaceae), Makkah to Taif road, Taif, 21.36528°N, 040.15897°E, 2015, coll. J.H. Mirza, M. Kamran.

Figure 3 *Eryngiopus discus* Meyer, adult female: A – Palp; B – Palptibial claw & palptarsus (not scaled).

Figure 4 *Eryngiopus discus* Meyer, adult female: A – Leg I; B – Leg II; C – Leg III; D – Leg IV.

Ecological note — The mite color was red-orange when collected. The species *Eryngiopus discus* was collected from foxtail grass *Cenchrus* sp. in association with date palm mite *Oligonychus afrasiaticus* (McGregor).

Key to the world species of the genus *Eryngiopus* Summers After Vacante & Gerson (1988)

Note — This key is based on the descriptions available in literature and few assumptions depending upon quality of illustrations.

1. Propodosoma with three pairs of setae (*vi*, *ve*, *sci*), prodorsal setae *sce* absent species group *bakeri* 2
— Propodosoma with four pairs of setae (*vi*, *ve*, *sci*, *sce*), prodorsal setae *sce* present species group *gracilis* 10
2. Aggenital setae two pairs *E. custodis* van Dis & Ueckermann
— Aggenital setae three pairs 3
3. Genu I with three setae 4
— Genu I with four setae 5
4. Tarsi II with 10 setae, ventral setae *3a* long = 88 *E. jiangxiensis* Hu & Chen
— Tarsi II with 9 setae, ventral setae *3a* short = 17 *E. nanchangensis* Hu & Chen
5. Femur I with five or six setae 6
— Femur I with four setae 7
6. Coxae IV with one seta, tarsi I with 14 setae *E. anatolicus* Doğan, Dilkaraoglu & Fan
— Coxae IV with two setae, tarsi I with 14 setae *E. yemenensis* van Dis & Ueckermann
7. Suranal shield entire (undivided) 8
— Suranal shield divided in to three feebly developed plates 9
8. Ventral setae *1a* very short (24), tarsi IV with seven setae *E. bakeri* (McGregor)
— Ventral setae *1a* ultra long (103), tarsi IV with eight setae *E. woodi* Gomaa & El-Enany
9. Setae *dl* less widely spaced $dl \neg dl = 80$, dorsal striations simple, aggenital setae (*ag2*) shorter 14 *E. yasumatsui* Ehara & Wongsiri
— Setae *dl* more widely spaced $dl \neg dl = 94-97$, dorsal striations with micro tubercles, aggenital setae (*ag2*) longer 23 *E. discus* Meyer
10. Aggenital setae two pairs 11
— Aggenital setae three pairs 16
11. Genu I with three setae *E. bidens* Meyer
— Genu I with four setae 12
12. Genu II without setae, intercoxal setae *1a* ≥ 70 13
— Genu II with one setae, intercoxal setae *1a* < 40 14
13. Prodorsal shield entire, terminal sensillum on palptarsus forked *E. bifidus* Wood
— Prodorsal shield divided, terminal sensillum on palptarsus simple .. *E. microsetus* Summers

14. Prodorsal shield completely striated	<i>E. parsimilis</i> Ueckermann & Meyer
— Prodorsal shield divided medially	15
15. Ventral setae 3a long = 42, terminal sensillum on palptarsus simple	<i>E. arboreus</i> Wood
— Ventral setae 3a short = 24, terminal sensillum on palptarsus forked	<i>E. similis</i> Wood
16. Femur I with 5 or 6 setae	17
— Femur I with 3 or 4 setae	20
17. Genu II with one seta, tarsi III and IV each with 8 setae	18
— Genu II with 2 setae, tarsi III and IV each with 7 setae	<i>E. nelsonensis</i> Wood
18. Coxae I with one seta, femur III with 3 setae	<i>E. vagantis</i> Summers
— Coxae I with 2 setae, femur III with 2 setae	19
19. Terminal sensillum on palptarsus simple, prodorsal shield divided medially	<i>E. hamedanicus</i> Khanjani, Mohammadi & Nazari
— Terminal palpal sensillum tridentate, prodorsal shield entire	<i>E. affinis</i> Barilo
20. Femur I with 3 setae	21
— Femur I with 4 setae	22
21. Genu setal formula 1-1-1-1, ratio $vi/vi-vi = 0.18$	<i>E. coimbatorensis</i> Gupta & David
— Genu setal formula 3-0-0-0, ratio $vi/vi-vi = 1.3$	<i>E. langroudiensis</i> Khanjani, Hazjizadeh & Ueckermann
22. Genu I with 3 setae	23
— Genu I with 4 setae	26
23. Coxae II with 2 setae	24
— Coxae II with 1 seta	25
24. Coxae IV with 2 setae, tarsi I with 11 setae	<i>E. hortus</i> (Chaudhri)
— Coxae IV with one seta, tarsi I with 14 setae	<i>E. cirrus</i> (Chaudhri)
25. Tarsi II with 9 setae, terminal palpal sensillum simple, first pair of intercoxal setae 1a large (>60)	<i>E. audreae</i> Maake, Ueckermann & Childers
— Tarsi II with 10 setae, terminal palpal sensillum forked, first pair of intercoxal setae 1a shorter	<i>E. lindei</i> Meyer
26. Trochanter IV with one seta	27
— Trochanter IV without setae	29
27. Coxae I with one seta, prodorsal shield divided medially	<i>E. tauricus</i> Kuznetsov
— Coxae I with 2 setae, prodorsal shield entire	28
28. Tibia IV with 4 setae, tarsi III and IV with 7 setae	<i>E. gracilis</i> Summers
— Tibia IV with 6 setae, tarsi III and IV with 8 setae	<i>E. summersi</i> Vacante & Gerson
29. Genu II with one seta	<i>E. dicotrichus</i> Fan, Zhang & Liu
— Genu II without setae	30

30. Suranal shield bearing two pairs of setae (*h1*, *h2*), propodosomal striations divergent beyond shields *E. placidus* Kuznetsov
 — Suranal shield with one pair of setae (*h1*), Propodosomal striations convergent beyond shields *E. siculus* Vacante & Gerson

Acknowledgement

The authors wish to thank the Deanship of Scientific Research at King Saud University, Riyadh for providing facilities and fund to complete this research work through the research project [RG-1438-055]. We are also grateful to Dr. Carlos H.W. Flechtmann (University of Sao Paulo, Departamento de Entomologia e Acarologia, Brazil) and Dr. Pholoshi A. Maake (Plant Protection Research Institute, Queenswood, South Africa) for providing literature.

References

- Alatawi F.J. 2011a. Phytophagous and predaceous mites associated with vegetable crops from Riyadh, Saudi Arabia. Saudi J. Biol. Sci., 18: 239-246. doi:10.1016/j.sjbs.2011.02.004
- Alatawi F.J. 2011b. Six new records of predaceous mites associated with some trees from Riyadh, Saudi Arabia. Acarines, 5: 37-399.
- Alatawi F.J., Kamran M. 2017. Predatory prostigmatid mite (Acari: Trombidiformes) fauna of the date palm agro-ecosystem in Saudi Arabia. Syst. Appl. Acarol., 22(9): 1444-1475. doi:10.11158/saa.22.9.11
- Chaudhri W.M., Akbar S., Rasool A. 1979. Studies on the predatory leaf inhabiting mites of Pakistan. US Department of Agriculture and Pakistan Agricultural Research Council. PL 480 Programme. Project No. PKARS, 30, pp. 139-229.
- Doğan S., Dilkaraoglu S., Fan Q.H., Erman O., Sevsay S., Adil S. 2015. Description of a new species of the genus *Eryngiopus* Summers (Acari: Stigmaeidae) from Turkey Syst. Appl. Acarol., 20(4): 431-440. doi:10.11158/saa.20.4.8
- Ehara S., Wongsiri T. 1984. Stigmaeid mites associated with plants in Thailand (Acarina, Stigmaeidae). Kontyu, 52(1), 110-118.
- Fan Q.-H., Zhang Y.-X., Liu Q.-Y. 2000. Raphignathoid mites on bamboo from Fujian, China (Acari: Prostigmata). Syst. Appl. Acarol., 4: 49-68. doi:10.11158/saasp.4.1.7
- Fan Q.-H., Flechtmann C.H. de Moraes G.J. 2016. Annotated catalogue of Stigmaeidae (Acari: Prostigmata), with a pictorial key to genera. Zootaxa, 4176(1): 1-199. doi:10.11646/zootaxa.4176.1.1
- Fan Q.-H., Zhang Z.-Q. 2005. Raphignathoidea (Acari: Prostigmata). Fauna N. Z. 52, pp. 1-400.
- Gomaa E.A., El-Enany M.A.M. 1985. Redescription of the genus *Eryngiopus* with a description of two new species from Egypt (Acari: Stigmaeidae). Bull. Zool. Soc. Egypt, 35: 92-97.
- Grandjean F. 1944. Observations sur les acariens de la famille des Stigmaeidae. Archives des Sciences Physiques et Naturelles, 26: 103-131.
- Gupta S.K. David H. 1990. A new *Eryngiopus* Summers (Acari: Stigmaeidae) from India. Entomon, 15(3-4): 281-282.
- Hu S.-Q., Chen X.-W. 1992. A new species of *Eryngiopus* from China (Acari: Stigmaeidae). Journal of Jiangxi University (Natural Science), 16(2): 108-110.
- Hu S.-Q., Chen X.-W. 1994. A new species of *Eryngiopus* from Jiangxi (Acari: Stigmaeidae). Journal of Nanchang University (Natural Science), 18(1): 93-96.
- Jafari S., Bagheri M., Saboori A. 2014. A new species of the genus *Favognathus* from Iran and redescription of *Eryngiopus affinis* Barilo (Acari: Cryptognathidae, Stigmaeidae). Syst. Appl. Acarol., 19: 353-362. doi:10.11158/saa.19.3.8
- Kethley J. 1990. Acarina: Prostigmata (Actinedida). Soil Biology Guide. John Wiley & Sons, New York, pp. 667-756.
- Khanjani M., Mohammadi L., Nazari A., Khanjani M. 2014. A new species of the genus *Eryngiopus* Summers (Acari: Stigmaeidae) from Hamedan Province, Iran. P. J. Acarol., 3: 129-135. doi:10.22073/pja.v3i2.10143
- Kuznetsov N.N. 1977. A contribution to the fauna of the family Stigmaeidae (Acariformes) in the Central-Chernozem zone. Zoologicheskii Zhurnal, 56(6): 953-956.
- Kuznetsov N.N. 1978. Revision of the genus *Stigmaeus* (Acariformes, Stigmaeidae). Zoologicheskii Zhurnal, 57(5): 682-694.
- Maake P.A., Ueckermann E.A., Childers C.C. 2015. A new species of the genus *Eryngiopus* Summers (Acari: Stigmaeidae) from citrus in Florida, USA. Int. J. Acarol., 41: 210-213. doi:10.1080/01647954.2015.1022218
- Matioli A.L., Ueckermann E.A., Oliveira C.A.L. 2002. Some stigmaeid and eupalopsellid mites from citrus orchards in Brazil (Acari: Stigmaeidae and Eupalopsellidae). Int. J. Acarol., 28(2): 99-120. doi:10.1080/01647950208684287
- Meyer M.K.P.S. 1969. Some stigmaeid mites from South Africa (Acari: Trombidiformes). Acarologia, 11(2): 227-271.

- Rehman M.U., Khan E.M., Alatawi F.J. 2016. A new species, a new record and a key to the reported stigmaeid species (Acariformes: Stigmaeidae) from Pakistan. Syst. Appl. Acarol., 21(9): 1267-1277. doi:[10.11158/saa.21.9.9](https://doi.org/10.11158/saa.21.9.9)
- Summers F.M. 1964. Three uncommon genera of the mite family Stigmaeidae (Acarina). Proceedings of the Entomological Society of Washington, 66: 184-192.
- Ueckermann E.A., Meyer M.K.P.S. 1987. Afrotropical Stigmaeidae (Acari: Prostigmata). Phytohyllactica, 19: 371-397.
- Vacante V., Gerson U. 1988. Three species of *Eryngiopus* (Acari: Stigmaeidae) from Italy, with key to species and summary of habitats. Redia, 70: 385-401.
- Van Dis J.C.S., Ueckermann E.A. 1993. New species of the genera *Stigmaeus* Koch, *Eryngiopus* Summers and *Eupalopsellus* Sellnick (Acari: Stigmaeidae, Eupalopsellidae) from the Afrotropical region. Phytohyllactica, 25: 117-135.