Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Supporting agricultural research
for sustainable development

**Acarologia is under free license and distributed under the terms of the
Creative Commons-BY**

Kuzinellus (Acari: Phytoseiidae) from China

Min Ma^a, Qing-Hai Fan^b, Sheng-Cai Li^a^a College of Agronomy, Shanxi Agriculture University, Taigu, China.^b Plant Health & Environment Laboratory, Ministry for Primary Industries, Auckland, New Zealand.

ABSTRACT

Two species of the genus *Kuzinellus* (Acari: Phytoseiidae) are known in China. We re-describe them based on types and fresh specimens and provide updated information and illustrations. It is the first time that *Kuzinellus trisetus* is recorded in Shanxi province.

Keywords Mesostigmata, Typhlodrominae, predatory mite**Zoobank** <http://zoobank.org/34DBCBD4-3436-4EA5-8E61-3B48BBCC793A>

Introduction

As a family of valuable beneficial predators Phytoseiidae has received an increasing attention in the last three decades (Helle and Sabelis, 1985; Lindquist *et al.*, 1996; McMurtry and Croft, 1997; Sabelis and Van Rijn, 1997; Gerson *et al.*, 2003; Moraes *et al.*, 2004). To date more than 2,750 species have been described (Demite *et al.*, 2017) and some species are commercially produced for controlling spider mites, thrips and whiteflies. The genus *Kuzinellus* Wainstein, 1976 is relatively small and consists of about 51 described species (Demite *et al.*, 2017; Kamran *et al.*, 2017). Only two species of this genus are known from China, an Oriental species, *K. cervix* (Wu and Li, 1984) from Hubei province and a Palearctic species, *K. trisetus* (Wu *et al.*, 1992) from Liaoning province. *Kuzinellus cervix* was subsequently found in Fujian, Hunan and Jiangxi (Wu *et al.*, 2009) but there was no record of *K. trisetus* from other provincial areas of the country. During a survey of phytoseiid mites of Shanxi province, *K. trisetus* was discovered from several localities. In this paper, we re-described both species based on the original types and fresh specimens of *K. trisetus*.

Materials and methods

The type specimens of *Kuzinellus cervix* and *K. trisetus* loaned from Guangdong Institute of Entomology, Guangzhou and fresh specimens of *K. trisetus* collected from Shanxi were used for this study. Illustrations were made using a drawing tube attached to a Nikon differential interference contrast (DIC) microscope, which was also used for measuring and imaging of specimens. Images and illustrations were edited with Photoshop CS4. Measuring method follows Ma *et al.* (2016). All measurements are given in micrometers (µm). The measurements for the holotype are followed into brackets by the range of measurements from paratypes or other specimens. The chaetotaxy of the idiosoma and legs follow Chant and McMurtry (2007) and Evans (1963), respectively, and the terminology of pore-like structures follows Beard (2001).

Results

Genus *Kuzinellus* Wainstein, 1976

Kuzinellus Wainstein, 1976: 699. Type species: *Paraseiulus kuzini* Wainstein, 1962: 139.

Received 06 January 2018
Accepted 19 June 2018
Published xx September 2018

Corresponding author
Qing-Hai Fan:
qinghai.fan@mpi.govt.nz

Academic editor
Marie-Stéphane Tixier

DOI
10.24349/acarologia/20184274

© Copyright
Ma M. *et al.*

Distributed under
Creative Commons CC-BY 4.0

How to cite this article Ma M. *et al.* (2018), *Kuzinellus* (Acari: Phytoseiidae) from China. *Acarologia* 58(4): 786-794; DOI 10.24349/acarologia/20184274

Diagnosis — Dorsal idiosomal setal patterns 13A:8A, with 19 pairs of setae on the dorsal shield. Setae *z6* and *JV2* present, *Z1* absent. Ventrianal shield with 4 pairs of preanal setae. *Kuzinellus* can be distinguished from its closely related *Paraseiulus* Muma by having *JV2* present and *Z1* absent. It differs from the other genus included in the tribe, Paraseiulini Wainstein (*Australiseius* Muma) by having *Z1* absent.

***Kuzinellus cervix* (Wu and Li, 1984)**

Typhlodromus (*Anthoseius*) *cervix* Wu and Li, 1984: 44; Moraes *et al.*, 2004: 317; Chant and McMurtry, 2007: 152.

Typhlodromus (*Paraseiulus*) *cervix*; Wu, 1985: 86.

Amblydromella cervix; Moraes *et al.*, 1986: 353.

Typhlodromus cervix; Wu and Lan, 1992: 1367; Hou, 1996: 13; Wu *et al.*, 1997: 172.

Kuzinellus cervix; Wu *et al.*, 2009: 330.

Redescription

ADULT FEMALE (n=2).

Dorsum (Figures 1A, 2A) — Dorsal shield nearly oval, incised at level of *R1*, reticulate throughout except area between *S4* and *S5*; 325 (336) long, 200 (203) wide; all setae smooth except *Z5* (barbed), bearing 11 pairs of discernible lyrifissures (*idl1a*, *id4*, *id6*, *idm3*, *idm5*, *idm6*, *is1*, *idl2*, *idl3*, *idl4* and *idx*) and 4 pairs of solenostomes (*gd2*, *gd6*, *gd8* and *gd9*). Muscle marks discernible, mainly on podosomal areas between *z2* and *j4*, *j4* and *j5*, *j5* and *z5*, *s4* and *z5*, *z5* and *j6*, *j6* and *J2*, *S2* and *S4*, and prior to *J5*. Lateral setae *r3* and *R1* smooth, on soft membranous cuticle lateral to dorsal shield, *r3* at level of *z4*, *R1* at level of shield incisions. Peritremes extending forward to bases of *j1*. Lengths of setae: *j1* 17 (17), *j3* 16 (17), *j4* 13 (12), *j5* 12 (12), *j6* 14 (14), *z2* 16 (16), *z3* 17 (16), *z4* 18 (19), *z5* 15 (16), *z6* 15 (15), *s4* 17 (19), *s6* 20 (20), *J2* 16 (16), *J5* 11 (12), *Z4* 19 (18), *Z5* 27 (29), *S2* 18 (20), *S4* 21 (21), *S5* 19 (21), *r3* 17 (21), *R1* 17 (18).

Ventral idiosoma (Figures 1B, 2B) — Sternal shield smooth (without reticulation), 56 (56–57) wide, anterior margin moderately convex, medial posterior margin vague, bearing 2 pairs of attenuate setae (*st1*, *st2*) and 2 pairs of lyrifissures (*iv1*, *iv2*), *iv1* between *st1* and *st2*, *iv2* at corners of the posterior margin; setae *st3* on soft cuticle, *st4* on platelets. Genital shield smooth, 113 (120) long, 58 (60) wide. Lengths of setae: *st1* 19 (20), *st2* 13 (12), *st3* 19 (19), *st4* 15 (13), *st5* 11 (18). A thin and long transversal sclerite between genital and ventrianal shields, a pair of small plates between *ZV1* and *ZV2*. Ventrianal shield approximately pentagonal, smooth, anterior margin convex, with prominent waist at *JV2* level, 116 (118) long, 75 (76) wide, with 4 pairs of pre-anal setae (*JV1*, *JV2*, *JV3* and *ZV2*) and a pair of pores *gv3* posteromedial to *JV2*; distance *gv3*–*gv3* 18 (18), 4 pairs of setae (*ZV1*, *ZV3*, *JV4* and *JV5*) and 3 pairs of lyrifissures on soft cuticle surrounding ventrianal shield. Primary metapodal plate 30 (32) long, 4 (4) wide, the secondary plate 8 (9) long, 2 (1) wide.

Calyx of insemination apparatus tubular (Figure 1D, 2D, 2E), 37 (39) long and 2 (2) wide, major duct long and slender, 34 long, 1 wide, minor duct thread-like, connecting to centre of atrium.

Gnathosoma — Chelicera (Figure 1C, 2C) with movable digit 25 (26) long, bearing 2 teeth, fixed digit 25 (27) long, bearing 3 discernible teeth, terminal two next to each other.

Leg lengths (I–IV): 299 (307), 232 (233), 230 (269) and 289 (344). Legs I, II and III without macrosetae. Basitarsus and telotarsus of leg IV (Figure 1E) each with a smooth macroseta, 20 (20) and 25 (31) long, respectively.

ADULT MALE. Unknown.

Material examined

Holotype female, Hubei: Shiyan, Shennongjia National Nature Reserve, *Pinus massoniana*, 15.VIII.1981, collected by Wei-nan Wu and Zhao-quan Li. Paratype: 1 female, same data

Figure 1 *Kuzinellus cervix* Wu and Li, 1984 (female). A – Dorsal idiosoma; B – Ventral idiosoma; C – Chelicera; D – Insemination apparatus; E – Legs.

Figure 2 *Kuzinellus cervix* Wu and Li, 1984 (female). A – Dorsal idiosoma; B – Ventral idiosoma; C – Chelicerae; D & E – Insemination apparatus.

as holotype. Both specimens are deposited at the Guangdong Institute of Applied Biological Resources, Guangdong, China.

Distribution

China: Fujian (Wu, 1985; Zhang and Lin, 1987), Hubei (Wu and Li, 1984), Hunan (Wu and Lan, 1992), Jiangxi (Hou, 1996).

Habitat

Fragaria × *ananassa* (Hou, 1996), *Hyllostachys heterocyclus* (Zhang and Lin, 1987), *Pinus massoniana* (Wu and Li, 1984; Wu, 1985).

Remarks

In their original description Wu and Li (1984) miscounted the number of dorsal setae, overlooked *z6* and incorrectly illustrated the sternal shield. Wu *et al.* (1997) corrected these mistakes. In 2009 Wu *et al.* transferred this species from *Typhlodromus* (*Anthoseius*) to *Kuzinellus*. We herein confirm their corrections and changes. Apart from these we have noted and illustrated the following characters which were not presented in the original and subsequent publications (Wu and Li, 1984; Wu *et al.*, 1997; 2009): lyrifissures, gland openings and muscle marks on idiosoma; atrium, major and minor ducts of insemination apparatus, and legs I–III. The second tooth of the fixed digit of chelicera is very small and situated next to the apical tooth. The margins of the platelets bearing *st4* and *iv2* were not discernible in the long preserved specimens. This species was originally described as *Typhlodromus* (*Anthoseius*) *cervix* and listed as such in the catalog of Moraes *et al.* (1986) and in the Phytoseiidae database of Demite *et al.* (2017), which would need to be updated.

Kuzinellus trisetus (Wu, Lan & Zhang, 1992)

Typhlodromus trisetus Wu *et al.*, 1992: 48; Wu *et al.*, 1997: 173.

Amblydromella (*Amblydromella*) *trisetus*; Denmark and Welbourn, 2002: 307.

Kuzinellus trisetus; Moraes *et al.*, 2004: 274; Wu *et al.*, 2009: 331.

Redescription

ADULT FEMALE (n=5).

Dorsum (Figures 3A, 4A) — Dorsal shield nearly oval, incised at level of *R1*; 399 (390–414) long, greatest width 221 (211–228), 208–218 wide, reticulate throughout except area between *Z4* and *S5*; all setae smooth except *Z5* (barbed). Dorsal shield with 14 pairs of discernible lyrifissures (*id1a*, *id2*, *id4*, *id6*, *idm2*, *idm3*, *idm4*, *idm5*, *idm6*, *idx*, *is1*, *idl2*, *idl3* and *idl4*) and 6 pairs of solenostomes (*gd2*, *gd4*, *gd6*, *gd8* and *gd9*). Muscle marks discernible, mainly on dorsal shield between dorsal setae and median setae, and anterior to *J5*. Lateral setae *r3* and *R1* smooth, on soft membranous cuticle laterad of dorsal shield, *r3* at level between *z4* and *s4*, *R1* at level of shield incisions. Peritremes extending forward to bases of *j1*. Lengths of setae: *j1* 18 (16–22), *j3* 16 (14–17), *j4* 14 (12–15), *j5* 13 (12–14), *j6* 14 (13–15), *z2* 15 (13–17), *z3* 16 (16–18), *z4* 17 (16–18), *z5* 15 (13–18), *z6* 15 (13–16), *s4* 18 (15–20), *s6* 21 (20–21), *J2* 19 (17–20), *J5* 12 (11–13), *Z4* 21 (20–23), *Z5* 27 (25–30), *S2* 20 (20–22), *S4* 21 (19–24), *S5* 19 (18–20), *r3* 19 (17–21), *R1* 18 (17–20).

Ventral idiosoma (Figures 3B, 4B, 4C) — Sternal shield approximately as long as wide, 60 (57–65) long, 59 (56–62) wide, smooth, anterior margin straight, posterior margin have 2 corners, posterior margin medially smooth or zigzagged, bearing 2 pairs of attenuate setae (*st1*, *st2*) and 2 pair of lyrifissures (*iv1*, *iv2*), *iv1* close to *st1*, *iv2* at corners of posterior margin; setae *st3* and *st4* on membranous cuticle. Genital shield smooth, 128 (124–132) long, 60 (54–62) wide. Lengths of setae: *st1* 23 (20–24), *st2* 21 (20–21), *st3* 21 (18–23), *st4* 20 (16–21), *st5* 22 (21–23). A series of slender transversal sclerites present between genital and ventrianal

Figure 3 *Kuzinellus trisetus* (Wu, Lan and Zhang, 1992) (female). A – Dorsal idiosoma; B – Ventral idiosoma; C – Chelicerae; D – Insemination apparatus; E – Legs.

Figure 4 *Kuzinellus trisetus* (Wu, Lan and Zhang, 1992) (female). A – Dorsal idiosoma; B – Sternal shield; C – Epigynal shield and ventrianal shield; D & E – Chelicerae; F, G & H – Insemination apparatus.

shields, a pair of small plates present between *ZV1* and *ZV2*. Ventrianal shield (Plate 1C) approximately pentagonal, smooth, anterior margin convex, with a prominent waist at level of *JV2*, 125 (122–130) long, 76 (64–80) wide, bearing 4 pairs of pre-anal setae (*JV1*, *JV2*, *JV3* and *ZV2*) and a pair of pores (*gv3*) posteromedial to *JV2*, distance *gv3*–*gv3* 21 (18–25), 4 pairs of setae (*ZV1*, *ZV3*, *JV4* and *JV5*) and 3 pairs of lyrifissures on soft cuticle surrounding ventrianal shield. Primary metapodal plate 39 (36–42) long, 5 (4–7) wide, the secondary plate 14 (13–15) long, 2 (2–3) wide.

Calyx of insemination apparatus (Figures 3D, 4F, 4G, 4H) U- or V-shaped with basal 1/2 to 2/3 membranous and thin, and apical 1/3 to 1/2 thick.

Gnathosoma — Chelicera (Figures 3C, 4D, 4E) with movable digit 25 (24–26) long, bearing 2 or 3 teeth, fixed digit 27 (25–28) long, bearing 3 discernible teeth, *pilus dentilis* thorn-shaped, 8 (6–9) long, opposing distal tooth of movable digit.

Leg lengths (I–IV): 356 (341–371), 290 (287–305), 298 (290–301), 390 (380–402). All legs without obvious macrosetae (Figure 3E).

ADULT MALE. Unknown.

Material examined

Holotype female, 10 paratype females, Qianshan, Liaoning, ex *Larix* sp.24.VII.1984, collected by Wei-nan Wu; 3 females, Ningwu, Luyashan National Nature Reserve, 38°44'34" N, 111°55'10" E, 2491 m, ex *Pinus* sp., 7.IX.2014, collected by Bing-Qian Su and Meng-Jiao Yin (accession no.: T14_0268); 1 female, Huguan, Taihang Mountains Grand Canyon, Purple Mass of Mountain, 35°54'47" N, 113°29'44" E, 1446 m, ex *Viburnum mongolicum*, 6.X.2014, collected by Min Ma (T14_0346); 1 female, same data as T14_0346 except: 35°54'47" N, 113°29'44" E, 1454 m, *Spiraea trilobata*, 7.X.2014, (T14_0347); 11 females, same data as T14_0268 except: Jiaocheng, Pangquangou National Nature Reserve, 37°49'35" N, 111°27'56" E, 1793 m, 28.VIII.2014, (T14_0191).

Holotype and paratypes are deposited in Guangdong Institute of Applied Biological Resources, Guangdong, China. Specimens collected from Shanxi are deposited in the Insect Ecology Laboratory, College of Agronomy, Shanxi Agriculture University, Taigu, China.

Distribution

China: Henan (Lin *et al.*, 2010), Liaoning (Wu *et al.*, 1992), Shanxi (present paper).

Habitat

Fern, *Dianthus* sp. (Lin *et al.*, 2010); *Larix* sp. (Wu *et al.*, 1992); *Pinus* sp.; *Spiraea trilobata*, *Viburnum mongolicum* (present paper).

Remarks

This is the first report of *Kuzienllus trisetus* in Shanxi province. We have noted and illustrated the following characters which were not presented in the original publication (Wu *et al.*, 1992): lyrifissures, gland openings and muscle marks on idiosoma, and legs I–III. The detailed structure of the insemination apparatus is given to show the unevenness of the calyx. The number of teeth on the movable digit is variable, from two to three. The apical tooth on the fixed digit is sometimes bifurcate. The areas around *st4* and *iv2* were not discernible in the long preserved specimens.

Key to species of *Kuzienllus* Wainstein in China (female)

1. Spermatheca with calyx elongate and tubular, uniformly sclerotized (Fig. 1D) ..... *Kuzienllus cervix* (Wu and Li, 1984)
- Spermatheca with calyx short, U- or V-shaped, with basal 1/2 to 2/3 membranous (Fig. 2D) ..... *Kuzienllus trisetus* (Wu, Lan and Zhang, 1992)

Acknowledgements

We would like to thank Dr Zhi-Qiang Zhang (Landcare Research, New Zealand) for his critical comments on a preliminary version of this manuscript. We appreciate Misses Bing-Qian Su and Meng-Jiao Yin for their assistance in the collection of specimens.

References

- Beard J.J. 2001. A review of Australian *Neoseiulus* Hughes and *Typhlodromips* De Leon (Acari: Phytoseiidae: Amblyseinae). *Invertebr. Taxon.*, 15: 73–158. doi:10.1071/IT99017
- Chant D.A., McMurtry J.A. 2007. Illustrated keys and diagnoses for the genera and subgenera of the Phytoseiidae of the world (Acari: Mesostigmata). West Bloomfield: Indira Publishing House, USA., 219 pp.
- Demite P.R., Moraes G.J. de, McMurtry J.A., Denmark H.A., Castilho R.C. 2017. Phytoseiidae Database. [accessed on 12 Nov. 2017] Available from: www.lea.esalq.usp.br/phytoseiidae
- Denmark H.A., Welbourn W.C. 2002. Revision of the genera *Amblydromella* Muma and *Anthoseius* De Leon (Acari: Phytoseiidae). *Int. J. Acarol.*, 28(4): 291–316. doi:10.1080/01647950208684308
- Evans G.O. 1963. Observations on the chaetotaxy of the legs in the free-living Gamasina (Acari: Mesostigmata). *Bull. Brit. Mus. (Nat. Hist.) Zool.*, 10(5): 277–303. doi:10.5962/bhl.part.20528
- Gerson U., Smiley R.L., Ochoa R. 2003. Mites (Acari) for pest control. Blackwell Science Ltd., UK, 539 pp. doi:10.1002/9780470750995
- Helle W., Sabelis M. W. 1985. Spider mites: Their Biology, Natural Enemies and Control Vol. 1B. Elsevier, Amsterdam, The Netherlands, 458 pp.
- Hou P.H. 1996. Investigation on species and vertical distribution of Phytoseiidae in Wuyishan Nature Reserve, Jiangxi Province. *Jiangxi Plant Protection*, 19(2): 12–14.
- Kamran M., Basahih J.S., Alatawi F.J. 2017. A new species of *Kuzinellus* Wainstein, 1976 (Acari: Mesostigmata: Phytoseiidae) from Saudi Arabia, with a key to the world species. *Int. J. Acarol.*, 43(7): 545–551. doi:10.1080/01647954.2017.1360937
- Lin J.Z., Ma L.M., Zhang Y.X., Ji J., Chen, X. 2010. Investigation of free living gamasid mite in Henan, China (IV) (Acari: Mesostigmata). *Wuyi Sci. J.*, 26: 1–10.
- Lindquist E.E., Sabelis M.W., Bruin J. 1996. Eriophyoid Mites - Their Biology, Natural Enemies and Control. World Crop Pest Series Vol. 6, Amsterdam: Elsevier Science Publishers, The Netherlands, 790 pp.
- Ma M., Fan Q.-H., Li S.C. 2016. *Typhlodromus* Scheuten (Acari: Phytoseiidae) from Shanxi province of China. *Syst. Appl. Acarol.*, 21(12): 1614–1630. doi:10.11158/saa.21.12.3
- McMurtry J.A., Croft B.A. 1997. Life-styles of phytoseiid mites and their roles in biological control. *Ann. Rev. Entom.*, 42: 291–321. doi:10.1146/annurev.ento.42.1.291
- Moraes G.J. de, McMurtry J.A., Denmark H.A. 1986. A catalog of the mite family Phytoseiidae. References to taxonomy, synonymy, distribution and habitat. Brasilia: EMBRAPA - DDT, Brazil, 353 pp.
- Moraes G.J. de, McMurtry J.A., Denmark H.A., Campos C.B. 2004. A revised catalog of the mite family Phytoseiidae. *Zootaxa*, 434: 1–494. doi:10.11646/zootaxa.434.1.1
- Sabelis M.W., Van Rijn P.C.J. 1997. Predation by insects and mites. In: Lewis T. (Ed.). *Thrips as Crop Pests*. London: CAB-International. p. 259–354.
- Wainstein B.A. 1962. Some new predatory mites of the family Phytoseiidae (Parasitiformes) of the USSR fauna. *Entomol. Obozr.*, 41: 230–240.
- Wainstein B.A. 1976. New tribe of the family Phytoseiidae (Parasitiformes). *Zool. Zh.*, 55: 696–700.
- Wu W.N., Lan W.M. 1992. Phytoseiidae. In: Peng J., Liu Y. (Eds.). *Iconography of Forest Insects in Hunan China*. Changsha: Hunan Science and Technology Press. p. 1473.
- Wu W.N., Li, Z.Q. 1984. Three new species of the genus *Phytoseius* from south China (Acarina: Phytoseiidae). *Acta Entomol. Sin.*, 27(4): 457–461.
- Wu W.N. 1985. A new species of the genus *Typhlodromus*, with notes on two other species from Fujian province. *Wuyi Sci. J.*, 5: 83–87.
- Wu W.N., Lan W.M., Zhang, S.Y. 1992. New species and new records of phytoseiid mites from northeast China III (Acari: Phytoseiidae). *Acta Zootaxon. Sin.*, 17(1): 48–56.
- Wu W.N., Liang L.R., Lan, W.M. 1997. Acari: Phytoseiidae. *Economic Insect Fauna of China* 53. Beijing: Science Press, 227 pp.
- Wu W.N., Ou J.F., Huang J.L. 2009. *Fauna Sinica, Invertebrata vol. 47. Arachnida Acari: Phytoseiidae*. Beijing: Science Press, 511 pp.
- Zhang Y., Lin J. 1987. Description of agricultural mites from Fujian province (I). *Fujian J. Agri. Sci.*, 5(1): 51–59.