Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

**Acarologia is under free license and distributed under the terms of the
Creative Commons-BY**

Two species of *Acaricis* (Acari: Tenuipalpidae) from New Zealand, moved from the genus *Tenuipalpus*, with a key to the known species

Elizeu B. Castro^a, Jennifer J. Beard^b, Ochoa, Ronald^c, Reinaldo J. F. Feres^{a,d}

^a São Paulo State University (UNESP), Institute of Biosciences, Humanities and Exact Sciences (Ibilce), Campus São José do Rio Preto, SP, 15054-000, Brazil.

^b Queensland Museum, South Brisbane, Queensland, 4101, Australia.

^c Systematic Entomology Laboratory (SEL), Agricultural Research Service (ARS), United States Department of Agriculture (USDA), Beltsville Agricultural Research Centre (BARC), Maryland, 20705, USA.

^d CNPq-Brazil researcher

ABSTRACT

Acaricis Beard & Gerson (Acari: Tenuipalpidae) is a small genus of flat mites with three species described from Australian and New Zealand sedges (Poales: Cyperaceae). In this article, we redescribe two species, *Tenuipalpus montanus* Collyer and *T. alpinus* Collyer, and move them to the genus *Acaricis*, so that these species become *Acaricis montanus* (Collyer) **comb. nov.** and *A. alpinus* (Collyer) **comb. nov.**. These two species share several characters with other *Acaricis* that species of *Tenuipalpus* do not. A key to the world species of *Acaricis* is also proposed.

Keywords flat mites, false spider mites, taxonomy, *Acaricis montanus*, *Acaricis alpinus*

Zoobank <http://zoobank.org/F90E71E2-A31B-4FEF-A1E3-FEF3C55055EA>

Introduction

Acaricis Beard & Gerson (Acari: Tenuipalpidae) is a small genus of flat mites with three species described from Australia and New Zealand sedges, family Cyperaceae (Order Poales) (Beard & Gerson 2009; Xu & Zhang 2013). Beard & Gerson (2009) erected the genus based on description of two species: *A. plana*, collected on *Scleria sphacelata* F. Muell., and *A. danutae*, collected on *Carex* sp., both from Australia. Later, Xu & Zhang (2013) described *A. urigersoni*, collected on *Gahnia lacera* (R. Lesson ex A. Rich.) Steud., from New Zealand.

In this article, we redescribe two species, *A. montanus* (Collyer, 1973) collected on *Pimelea sericeovillosa* Hook. (Thymelaeaceae; Order Malvales), and *A. alpinus* (Collyer, 1973), collected on “mat plants on rocks” (indeterminate family), moving them to the genus *Acaricis* from *Tenuipalpus* Donnadieu. These two species were described based on specimens collected in New Zealand.

Materials and methods

Measurements for the paratypes are given in micrometers (µm). The terminology used follows that of Lindquist (1985) and Mesa *et al.* (2009). Leg chaetotaxy is adapted from Lindquist (1985), Zhang & Fan (2004), Xu & Fan (2010) and Seeman & Beard (2011). Tarsal setae are presented as the total number followed by the number of solenidia in parentheses.

Received 10 May 2018
Accepted 25 September 2018
Published 12 October 2018

Corresponding author
Elizeu B. Castro:
elizeu_unesp@yahoo.com.br

Academic editor
Philippe Auger

DOI
10.24349/acarologia/20184290

© Copyright
Castro E. B. *et al.*

Distributed under
Creative Commons CC-BY 4.0

How to cite this article Castro E. B. *et al.* (2018), Two species of *Acaricis* (Acari: Tenuipalpidae) from New Zealand, moved from the genus *Tenuipalpus*, with a key to the known species. *Acarologia* 58(4): 855-867; DOI 10.24349/acarologia/20184290

The paratypes used in the redescription are deposited in the National Insect and Mite Collection, National Museum of Natural History, Smithsonian Institution (NMNH), located in the Systematic Entomology Laboratory (SEL), USDA, Beltsville, Maryland, USA, and at the Natural History Museum (BMNH), located in London, England, UK.

Taxonomy

Family Tenuipalpidae Berlese, 1913

Genus *Acaricis* Beard & Gerson, 2009

Acaricis Beard & Gerson, 2009: 31; Mesa *et al.* (2009): 113; Xu & Zhang 2013: 357
Type species *Acaricis plana* Beard & Gerson, 2009

Diagnosis (Based on Xu & Zhang 2013)

Female — Body elongate, usually more than twice as long as wide; palpus usually four segmented. Anterior margin of prodorsum with median forked projection forming a notch (elongate in Aust. species; short in NZ species); prodorsum with two or three pairs of setae (*sc1*, *sc2* present; *v2* present or absent); setae *sc2* lanceolate or acicular. Dorsal opisthosoma with nine pairs of setae: *c3*, *d1*, *d3*, *e1*, *e3*, *f2*, *f3*, *h1* present; setae *c1*, *c2*, *d2*, *e2* absent; setae *h2* elongate and filiform. Venter with two pairs of *4a* setae; setae *g1* inserted anterior to *g2* on membranous genital flap; setae *ps3* inserted far anterior to *ps2*. Leg chaetotaxy: tarsi I–II with seta *ft''* present, *ft'* absent; tarsi III–IV with seta *ft'* present, *ft''* absent; tarsi I–II with seta *ft''* and tarsi III–IV with seta *ft'* elongate, filiform, inserted on short tubercle; trochanter IV with one seta or nude (seta *v'* present or absent). Leg setal formulae (legs I–IV): femora 4-4-2-2 or 4-4-2-1, genua 2-2-1-0 or 2-2-0-0, tibiae 5-5-3-3 or 5-5-3-2; setae *d*, *l'*, *l''* on femora, genua and tibia I–II thickened, barbed, inserted in lateral position.

Male— tarsi I–II with two solenidia (all species); tarsi III–IV with one solenidium (except *A. danutae* without solenidia).

Remarks

All previous known species of *Acaricis* have a body more than twice as long as wide, while in *A. montanus* and *A. alpinus* the body is marginally less than twice as long as wide. However, the lower body lengths presented here for these two species may be due to the age and mounting media of the specimens, and it is possible that the lengths could be longer in freshly mounted specimens, as has been observed by Beard *et al.* (in press). In the original description, the female *A. alpinus* is 354 long and 170 wide (approx. 2.1 times longer than wide) (here 290–300, 150–160, 1.8–1.9, respectively) and the male is 292 long and 146 wide (exactly 2 times longer than wide) (here 230, 125, 1.8, respectively). The female *A. montanus* is 370 long and 216 wide (approx. 1.7 times longer than wide) (here 330–360, 190, 1.7–1.9, respectively). Collyer (1973) does not provide a range for the lengths and widths of these two species, and the single measurements provided are assumed to represent the holotype in each case. The delimiting marker points for our length and width measurements are the setae *v2–h1* and *sc2–sc2*, and although the exact delimiting points for body measurements were not described by Collyer (1973), based on the first description in her paper, the length was the full body including gnathosoma and the width was taken at the widest point on the specimen. With this in mind, a difference between the two sets of measurements would be expected, and our measurements would be expected to be lower than Collyer's as ours are based on distances between dorsal setae and not between the limits of body margins on flattened specimens. Our measurements are 54–64 shorter in length (=18–28%) and 10–20 shorter in width (=6–12%) than the original description for *A. alpinus* females, and 30–40 shorter in length (=8–11%) and 26 shorter in width (=12%) for female *A. montanus*. As a comparison, Beard *et al.* (in press) found that body measurements from older specimens mounted in PVA were approximately 14% shorter and 10% narrower than those from fresh specimens mounted in Hoyer's medium.

Results

***Acaricis montanus* (Collyer, 1973)**

***Tenuipalpus montanus* Collyer, 1973: 949**

(Figures 1–4)

Diagnosis

Female — Anterior margin of prodorsum with a short median forked projection forming a shallow notch. Prodorsum with oblique to transverse striations; prodorsal setae *v2* and *sc1* minute; *sc2* short lanceolate, thick; palpi four segmented, with setal formula 0, 0, 2, 2. Central region of opisthosoma with series of transverse striations and lateral region with few longitudinal striations; setae *c1* absent; setae *c3*, *d1*, *d3*, *e1* and *e3* minute; setae *f2*, *f3*, and *h1* short lanceolate, thick; *h2* elongate and filiform. Ventral integument completely striated; setae *1a*, *4a1*, *4a2* elongate, filiform; two pairs of *4a* setae; setae *g1* inserted marginally anterior to *g2* on membranous genital flap; setae *ag*, *g1*–*g2* and *ps2*–*ps3* smooth. Leg chaetotaxy (legs I–IV): femora 4-4-2-1, genua 2-2-1-0, tibiae 5-5-3-3, tarsi 8(1)-8(1)-5-5. Body 1.7–1.9 X longer than wide.

Figure 1 *Acaricis montanus* (Collyer) female. A – dorsum; B – venter.

Figure 2 Differential interference contrast (DIC) image of *Acaricis montanus* (Collyer) female – dorsum.

Type material examined

Paratypes — 1 female collected on *Pimelea sericeo-villosa* Hook. (Thymelaeaceae), with litter and soil from beneath this shrub, Mount Grey, 1250m elevation, Turret Range, Lake Manapouri area, New Zealand, 09.i.1970, coll. A.C.E. (NMNH); 1 female collected on seepage sward of *Astelia linearis* (Asteliaceae), *Dracophyllum* (Epacridaceae), *Forstera* (Stylidiaceae), *Phyllachne* (Stylidiaceae), Mt Barber, 1100m, Wilmot Pass, West of Lake Manapouri, New Zealand, 8.i.1970, coll. J.S.D. and J.M.H (BMNH).

Remarks — The details of the collection data as written on the paratype slide deposited in the NMNH are different to what was presented by Collyer in the original description.

Figure 3 *Acaricis montanus* (Collyer) female: ventral infracapitulum.

Redescription

Female (Figures 1–4, n=2).

Body measurements — distance between setae *v2–h1* 330–360, *sc2–sc2* 190; other measurements: *v2–v2* 40–42, *sc1–sc1* 90–92, *c3–c3* 185–195, *d1–d1* 47–48, *d3–d3* 160–165, *e1–e1* 18–20, *e3–e3* 150–160, *f2–f2* 140–150, *f3–f3* 125–130, *h1–h1* 63–70, *h2–h2* 100.

Dorsum (Figures 1A, 2) — Anterior margin of prodorsum with short median forked projection forming a shallow notch. Prodorsum with oblique striations; prodorsal setae *v2* and *sc1* minute, and *sc2* short, thick lanceolate and barbed. Central region of opisthosoma with transverse striations and lateral region with few longitudinal striations; setae *c1* absent; *c3*, *d1*, *d3*, *e1* and *e3* minute; setae *f2*, *f3*, and *h1* short, thick lanceolate and barbed; *h2* elongate and filiform. Setal lengths: *v2* 4, *sc1* 4–5, *sc2* 20–23, *c3* 3–4, *d1* 4–5, *d3* 3–4, *e1* 4, *e3* 4–5, *f2* 15–17, *f3* 19–21, *h1* 12–13, *h2* 100–105.

Venter (Figure 1B). Ventral integument mostly with fine transverse striations; setae *1a*, *4a1*, *4a2* elongate and filiform; two pairs of setae *4a*; setae *g1* inserted marginally anterior to *g2* on membranous genital flap; setae *ag*, *g1–g2* and *ps2–ps3* smooth; other ventral setae fine and smooth. Setal lengths: *1a* 120–135, *1b* 9–10, *1c* 10–12, *2b* 12–14, *2c* 10, *3a* 9–11, *3b* 12–14, *4a1* 85–95, *4a2* 90–105, *4b* 11–13, *ag* 15–17, *g1* 16–17, *g2* 18–20, *ps2* 12–13, *ps3* 11–12.

Gnathosoma (Figure 3) — Palpi four segmented, setae formula: 0, 0, 2, 2; tibia with two setae, *d'* 6, *d''* 6, tarsus with two eupathidia, 4–5, 5. Ventral setae *m* 6; distance between setae *m–m* 14–15.

Legs (Figure 4) — Chaetotaxy (from coxae to tarsi): I 2-1-4-2-5-8(1), II 2-1-4-2-5-8(1), III 1-2-2-1-3-5, IV 1-1-2-0-3-5. Tarsus I–II each with one abaxial solenidion ω'' 7 and 6, respectively, and two eupathidia, *pζ'–pζ''* (tarsi I 7, 7; tarsi II 6, 6).

Male — Unknown.

Remarks — According to Collyer (1973), this species was also collected on moss, *Celmisia* (Asteraceae), *Senecio* (Asteraceae), *Gaultheria* (Ericaceae), *Coprosma* (Rubiaceae), *Haastia sinclairii* (Asteraceae), grasses (Poaceae) in New Zealand. This collection data suggests that the true host association(s) are unknown.

Figure 4 *Acaricis montanus* (Collyer) female, dorsal aspect, right side, of: A – leg I; B – leg II; C – leg III; D – leg IV.

***Acaricis alpinus* (Collyer, 1973)**
***Tenuipalpus alpinus* Collyer, 1973: 946.**
(Figures 5–9)

Diagnosis

Female — Anterior margin of prodorsum with a short median forked projection forming a shallow notch. Prodorsum mostly smooth, with weak transverse striations posteriorly; prodorsal setae *v2*, *sc1* minute, smooth; setae *sc2* short, smooth, acicular; palps four segmented; central opisthosoma between setae *d1* and *e1* with few weak transverse striations, lateral opisthosoma with few weak longitudinal striations; dorsal opisthosomal setae short to minute, except setae *h2* elongate and filiform. Ventral integument mostly with fine transverse striations; setae *1a*, *4a1*, *4a2* elongate, filiform; setae *g1* inserted marginally anterior to *g2* on membranous genital flap; setae *ag*, *g1*–*g2* and *ps2*–*ps3* smooth. Leg chaetotaxy (legs I–IV): femora 4-4-2-1, genua 2-2-1-0, tibiae 5-5-3-3, tarsi 8(1)-8(1)-5-5. Body 1.8–1.9 X longer than wide. **Male**: Dorsal integument and setae like those of female; tarsi I–II each with two solenidia; tarsi III–IV with one solenidia.

Figure 5 *Acaricis alpinus* (Collyer) A – female: dorsum; B – male: dorsum.

Figure 6 Differential interference contrast (DIC) image of *Acaricis alpinus* (Collyer) (Female): dorsum.

Type material examined — Paratypes: 1 female and 1 male collected from mixed sample of mat plants on rocks, Mount Robert Ridge circa 1830 m elevation, Nelson Lakes National Park, New Zealand; October 1969, coll. G.W.R. (NMNH); 1 female same data (BMNH).

Redescription

Female (Figures 5–8, n=2)

Body measurements — Distance between setae *v2–h1* 290–300, *sc2–sc2* 150–160; other measurements: *v2–v2* 32–33, *sc1–sc1* 74–78, *c3–c3* 135–150, *d3–d3* 125–145, *e1–e1* 22–25, *e3–e3* 100–105, *f2–f2* 102–105, *f3–f3* 86–87, *h1–h1* 30–36, *h2–h2* 67–70.

Dorsum (Figures 5A, 6) — Anterior margin of prodorsum with short median forked projection forming a shallow notch. Prodorsum mostly smooth with weak transverse striations posteriorly; prodorsal setae *v2*, *sc1* minute; setae *sc2* short, smooth; central opisthosoma

Figure 7 *Acaricis alpinus* (Collyer) female. A – ventral opisthosoma; B – ventral infracapitulum.

between setae *d1* and *e1* with weak transverse striations; lateral opisthosoma with few weak longitudinal striations; dorsal opisthosomal setae short to minute, except setae *h2* elongate, filiform. Setal lengths: *v2* 3–4, *sc1* 4–5, *sc2* 7, *c3* 4, *d1* 3–4, *d3* 3–4, *e1* 4, *e3* 3–4, *f2* 5, *f3* 5, *h1* 6–7, *h2* 65–70.

Venter (Figure 7A) — Ventral integument with fine transverse striations; setae *1a*, *4a1*, *4a2* elongate, filiform; two pairs of setae *4a*; setae *g1* inserted marginally anterior to *g2* on membranous genital flap; setae *ag*, *g1*–*g2* and *ps2*–*ps3* smooth; other ventral setae fine and smooth. Setal lengths: *1a* 85–105, *1b* 9–11, *1c* 9–10, *2b* 11–12, *2c* 10–14, *3a* 10–12, *3b* 11–12, *4a2* 30–43, *4a2* 35–50, *4b* 8–10, *ag* 7–8, *g1* 8–9, *g2* 7–8, *ps2* 11–12, *ps3* 10–11.

Gnathosoma (Figure 7B) — Palpi four segmented, setae formula: 0, 0, 2, 2; tibia with two setae, *d'* 5, *d''* 5, tarsus with two eupathidia 3, 2. Ventral setae *m* 6–7; distance between setae *m*–*m* 12–13.

Legs (Figure 8) — Chaetotaxy (from coxae to tarsi): I 2-1-4-2-5-8(1), II 2-1-4-2-5-8(1), III 1-2-2-1-3-5, IV 1-1-2-0-3-5. Tarsi I–II each with one abaxial solenidion ω'' 8 and 7, respectively, and two eupathidia distally *pζ'*–*pζ''* (7, 7 for both tarsi).

Male (Figures 5B, 9, n=1)

Body measurements — Distance between setae *v2*–*h1* 230, *sc2*–*sc2* 125; other measurements: *v2*–*v2* 32, *sc1*–*sc1* 65, *c3*–*c3* 100, *d1*–*d1* 33, *d3*–*d3* 90, *e1*–*e1* 20, *e3*–*e3* 68, *f2*–*f2* 65, *f3*–*f3* 55, *h1*–*h1* 30, *h2*–*h2* 50.

Dorsum (Figure 5B) — Dorsal integument and setae similar to those of female. Setal lengths: *v2* 4, *sc1* 3, *sc2* 7, *c3* 4, *d1* 3, *d3* 3, *e1* 3, *e3* 3, *f2* 4, *f3* 5, *h1* 5, *h2* 55. **Venter.** Ventral integument and setae similar to those of female; setae *1a*, *4a1*, *4a2* elongate and filiform. Setal

Figure 8 *Acaricis alpinus* (Collyer) female: dorsal aspect, right side, of: A – leg I; B – leg II; C – leg III; D – leg IV.

Figure 9 *Acaricis alpinus* (Collyer) male: dorsal aspect, right side, of: A – leg I; B – leg II; C – leg III; D – leg IV.

lengths: *la* 100, *lb* 6, *lc* 8, *2b* 8, *2c* 7, *3a* 13, *3b* 11, *4a1* 38, *4a2* 40, *4b* 8, *ag* 7, *g1* 13, *g2* 13, *ps2* 8, *ps3* 6.

Gnathosoma — Similar to that of female; tibia with two setae, *d'* 5, *d''* 6, tarsus with two eupathidia 3, 2. Ventral setae *m* 4; distance between setae *m–m* 13.

Legs (Figure 9) — Chaetotaxy (from coxae to tarsi): I 2-1-4-2-5-9(2), II 2-1-4-2-5-9(2), III 1-2-2-1-3-6(1), IV 1-1-1-0-3-6(1). Setae *d*, *l'*, *l''* on femora, genu and tibia I–II slightly thickened, weakly barbed. Tarsi I–II each with two solenidion (one abaxial, one adaxial), tarsi I *ω'* 7, *ω''* 7, tarsi II *ω'* 7, *ω''* 7, and two eupathidia *pζ'*–*pζ''* (all 6–7); tarsus III–IV each with one abaxial solenidion, tarsi III 6, tarsi IV 7.

Remarks

According to Collyer (1973), this species was also collected from mixed mats of plants including: *Chionochloa crassiuscula* (Poaceae), *Astelia*, *Celmisia sessiliflora* (Asteraceae), *Cyathodes pumila* (Epacridaceae), *Gaultheria*, *Anisotome* (Apiaceae) and *Drosera* (Droseraceae) in New Zealand. This collection data suggests that the true host association(s) are unknown.

Discussion

All previously known species of *Acaricis* were collected on plants of the family Cyperaceae; whereas *A. montanus* and *A. alpinus* were collected in association with many species and families of alpine heath plants, none of which were identified as Cyperaceae (Collyer 1973). In discussing *A. montanus* and *A. alpinus*, Collyer (1973) states herself that there “can be no certainty of their host plant relationships” as the specimens were collected from samples of mixed plants. Despite the apparent difference in host relationships, *A. montanus* and *A. alpinus* both share with other *Acaricis* species: anterior margin of prodorsum with median forked process, two pairs of ventral setae *4a*; dorsal setae *c1* is absent; the palps are four segmented with setae formula 0, 0, 2, 2; genital setae *g1* are inserted (marginally) anterior to *g2*; setae *ft'* are suppressed on tarsus I–II and *ft''* suppressed on tarsi III–IV; setae *ft''* are filiform on tarsi I–II; and the males have two solenidia on tarsi I–II. *Acaricis alpinus* males also share one solenidion on tarsus III–IV with *A. plana* and *A. urigersoni* (male not known for *A. montanus*; *A. danutae* males without solenidion on tarsus III–IV). Additional solenidia on tarsus III are also commonly seen in species of *Tenuipalpus* sensu stricto, but are not commonly seen on tarsus IV (Castro *et al.* 2015). *Acaricis montanus* and *A. alpinus* differ to all other *Acaricis* in having a seta on trochanter IV, while the other three known species have a nude trochanter IV. The other New Zealand species, *A. urigersoni*, is the only *Acaricis* to have prodorsal setae *v2* absent, it also has different chaetotaxy on leg IV to all other species in the genus, with only one seta on femur IV (seta *d* present, *ev'* absent), only two setae on tibiae IV (*v'*, *v''* present; *d* absent); and genua III–IV are nude (only genua IV nude in other *Acaricis* species). All three New Zealand species also share a short median forked process on the anterior margin of the prodorsum, which is much longer in the two Australian species.

Key to species of *Acaricis* (based on adult females)

1. Prodorsal setae *v2* absent; prodorsal setae *sc1* minute, bifurcate; dorsal opisthosomal setae *e3* lanceolate; femora IV with one seta (*d* present; *ev'* absent); genua III–IV nude; tibiae IV with two setae (*v'*, *v''* present; *d* absent) *A. urigersoni* Xu & Zhang
— Prodorsal setae *v2* present; prodorsal setae *sc1* acicular; dorsal opisthosomal setae *e3* minute, acicular or barbed; femora IV with two setae (*d*, *ev'* present); genua III with one seta (*d* present), genua IV nude; tibiae IV with three setae (*d*, *v'*, *v''* present) 2
2. Body elongate, more than twice as long as wide; median forked projection on anterior margin of prodorsum long with deep notch; genital setae *g1* inserted well anterior to *g2* 3

- Body almost twice as long as wide (1.7–1.9 X longer than wide); median forked projection on anterior margin of prodorsum short with shallow notch; genital setae *g1* inserted marginally anterior to *g2* 4
3. Dorsal opisthosomal setae *e3* minute, fine; prodorsal setae *sc2* lanceolate, stout; dorsal seta *d* on femur III lanceolate, thickened *A. plana* Beard & Gerson
- Dorsal opisthosomal setae *e3* short, thickened, barbed; prodorsal setae *sc2* short, barbed; dorsal seta *d* on femur III short, fine *A. danutae* Beard & Gerson
4. Prodorsal setae *sc2* lanceolate, stout (20–23 µm) and distinctly longer than setae *v2* and *sc1* (both minute; 4–5 µm) *A. montanus* (Collyer)
- Prodorsal setae *sc2* acicular, short, fine (7 µm) and slightly longer than *v2* and *sc1* (both minute; 3–5 µm) *A. alpinus* (Collyer)

Acknowledgments

To Debra Creel and Andrew Ulsamer (SEL-USDA) for their help with references and technical support. To the Smithsonian Natural History Museum and National Agricultural Library (NAL-USDA), SEL-USDA for support and assistance with specimens and references. We thank Dr Janet Beccaloni, British Museum of Natural History (BMNH) for lending the types. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the USDA; USDA is an equal opportunity provider and employer. This work was supported by the "Fundação de Amparo à Pesquisa do Estado de São Paulo" (FAPESP) (Procs. 2016/01193-5 and 2017/00458-8) through a fellowship to the first author; by a research grant from the Australian Biological Resources Study, Department of Environment and Energy, Australian Government [grant RF214-23] to the second author; and by a research grant from the "Conselho Nacional de Desenvolvimento Científico e Tecnológico" (CNPq) (Proc. 303435/2013-5) to the fourth author.

References

- Beard J.J., Gerson U. 2009. A new flat mite genus, *Acaricis* (Prostigmata: Tenuipalpidae), from Australian sedges (Cyperaceae). *Zootaxa*, 2073: 31–44.
- Beard, J.J., Ochoa R., Baughan, G.R., Pooley, C., Dowling, A.P.G. In press. *Raoiella* of the world (Trombidiformes: Tetranychidae: Tenuipalpidae). *Zootaxa*.
- Castro E.B., Ramos F.E.M., Feres R.J.F., Ochoa R. 2015. A new species of *Tenuipalpus* Donnadieu (Acari: Tenuipalpidae) from Brazil, with ontogeny of chaetotaxy. *Syst. and Appl. Acarol.*, 20: 339–356. doi:10.11158/saa.20.3.11
- Collyer E. 1973. New species of the genus *Tenuipalpus* (Acari: Tenuipalpidae) from New Zealand, with a key to the world fauna. *N.Z. J. Sci.*, 16: 915–955.
- Lindquist E.E. 1985. External anatomy. In: Helle W., Sabelis M.W. (Eds), *Spider mites: their biology, natural enemies and control*. Vol. 1A. Chapter 1.1 Anatomy, Phylogeny and Systematics. Elsevier Science Publishers, Amsterdam, p. 3–29.
- Mesa N.C., Ochoa R., Welbourn W.C., Evans G.A., Moraes G.J.de. 2009. A catalog of the Tenuipalpidae (Acari) of the World with a key to genera. *Zootaxa*, 2098: 1–185.
- Seeman O.D., Beard J.J. 2011. A new species of *Aegyptobia* (Acari: Tenuipalpidae) from Myrtaceae in Australia. *Syst. and Appl. Acarol.*, 16: 73–89. doi:10.11158/saa.16.1.10
- Xu Y., Fan Q.-H. 2010. *Tenuipalpus orilloi* Rimando, a new record to the Chinese fauna (Acari: Tenuipalpidae). *Syst. and Appl. Acarol.*, 15: 135–138. doi:10.11158/saa.15.2.8
- Xu Y., Zhang Z.-Q. 2013. New Zealand Tenuipalpidae (Acari: Trombidiformes): A new species of *Acaricis* from Cyperaceae and its ontogenetic patterns in chaetotaxy. *Syst. and Appl. Acarol.*, 18: 357–388. doi:10.11158/saa.18.4.6
- Zhang Z.-Q., Fan Q.-H. 2004. Redescription of *Dolichotetranychus ancistrus* Baker & Pritchard (Acari: Tenuipalpidae) from New Zealand. *Syst. and Appl. Acarol.*, 9: 111–131. doi:10.11158/saa.9.1.17