Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Acarologia is under free license and distributed under the terms of the
Creative Commons-BY

New mite records (Acari: Mesostigmata, Trombidiformes) from soil and vegetation of some Syrian citrus agrosystems

Ziad Barbar^a

^a Department of Plant Protection, Faculty of Agriculture, Al-Baath University, P.O. Box 77, Al- Sham St., Homs, Syria.

ABSTRACT

This study gives the result of collections of mite fauna inhabiting soil, litter and plant species surrounding citrus orchards at three localities in Latakia governorate. Mites collected were identified and only new recorded species are presented. During this study, 21 species belonging to 12 families (4 Mesostigmata and 8 Trombidiformes) were reported as new to the Syrian fauna. Five species were phytophagous mites (belong to Tenuipalpidae and Tetranychidae), while the others are considered as generalist predators or feed on microorganisms in their habitats. The results could be of importance for integrated pest management applied in Syrian citrus orchards.

Keywords survey; taxonomy; phytophagous mites; predatory mites; biological control; Syria

Zoobank <http://zoobank.org/19DAAD5E-3742-4DF9-8E1C-CBEBEE00706C>

Introduction

Latakia is the main growing citrus governorate in Syria, produced about 0.90 million tons of citrus fruit in 2016 (Syrian Ministry of Agriculture and Agrarian Reform, 2016). In this governorate, recent surveys conducted in about 60 citrus orchards revealed the presence of 60 mite species belonging to 20 families. Of these, 22 mite species belonging to 11 families inhabit citrus leaves (nine Phytoseiidae, three Tetranychidae, two Cheyletidae, and one of each Acaridae, Camerobiidae, Cunaxidae, Iolinidae, Raphignathidae, Stigmaeidae, Tenuipalpidae and Tydeidae). Remaining taxa were collected from litter and (or) from vegetation adjacent to those orchards (Zriki *et al.*, 2015; Barbar, 2016, 2017; Barbar and Ueckermann, 2017; Barbar, 2018). These results showed the presence of a relatively high number of mite species, especially predators of phytophagous mites and other micro arthropods (Barbar, 2016). However, more investigations are needed to obtain a good vision about the diversity of this fauna in Latakian citrus agro ecosystems. The results of these investigations could be of a great interest to further studies about the use of predatory mites in the biological control programs.

The aim of the present study was to identify mite species inhabiting soil, litter and plant species surrounding citrus orchards at three localities in Latakia governorate, and to provide data (when available) about their trophic habits, geographic distribution and economic importance.

Materials and methods

Surveys of mite fauna were conducted in Latakia governorate in April and May, 2016, March and April, 2017, and March, 2018. Mite collected from: (1) leaves of common plant species surrounding citrus orchards located at Burj-Islam (35°40'12" N, 35° 7'24" E) and Attabiyyat

Received 23 April 2018
Accepted 21 May 2018
Published 09 November 2018

Corresponding author
Ziad Barbar:
ziadbarbar88@gmail.com

Academic editor
Serge Kreiter

DOI
10.24349/acarologia/20184298

© Copyright
Barbar Z.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

How to cite this article Barbar Z. (2018), New mite records (Acari: Mesostigmata, Trombidiformes) from soil and vegetation of some Syrian citrus agrosystems. *Acarologia* 58(4): 919-927; DOI 10.24349/acarologia/20184298

(35°30'24" N, 35°46'49" E); (2) ground litter of a pesticides free citrus orchard located at Al-ya'robiyah (35°30'24" N, 35°48'33" E). Mites were removed from leaves using the "dipping-checking-washing-filtering" method (Boller, 1984). For collecting mites from litter, materials were placed on a sieve (25 cm Ø x 10 cm; its screen with 5 mesh/cm) and shaken over a black plastic sheet (1.5 m²). Mites were picked off the sheet into a vial with 70 % alcohol. Mites were then mounted on slides in Hoyer's medium and dried in an oven at 40°C for four days.

Mite taxa were identified to family levels using the key of Krantz and Walter (2009). Identification to genera level was carried out mainly using the following works: **(1)** for Mesostigmata: Ameroseiidae, Ascidae and Laelapidae (Athias-Henriot, 1961; Evans and Till, 1979; Mašán, 2017) and Phytoseiidae (Chant and McMurtry, 2007); **(2)** for Trombidiformes: Anystidae (Meyer and Ueckermann, 1987), Cunaxidae (Skvarla *et al.*, 2014), Cryptognathidae (Khanjani and Ueckermann, 2008), Eupalopsellidae (Meyer and Ueckermann, 1984), Raphignathidae (Fan and Zhang, 2005), Stigmaeidae (Fan *et al.*, 2016), Tenuipalpidae (Meyer, 1979; Mesa *et al.*, 2009) and Tetranychidae (Bolland *et al.*, 1998). The specimens were deposited in the Arthropod Collection of the Department of Plant Protection, Faculty of Agriculture, Al-Baath University, Homs, Syria. Mite taxa in the text are arranged alphabetically by order, as are families within orders, genera within families and species within genera.

All mite species that have already been recorded from Syria were excluded from results.

Results and discussion

Twenty-one mite species belonging to 12 families (4 Mesostigmata and 8 Trombidiformes) are recorded for the first time from Syria in the present study. Five species were phytophagous (three species of Tenuipalpidae and two species of Tetranychidae), while the others are considered as generalist predators or feeding on microorganisms in their habitats (Swift and Goff, 2001; Gerson *et al.*, 2003). Eleven species were found in soil and litter: 1 Ameroseiidae, 1 Ascidae, 2 Laelapidae, 2 Anystidae, 1 Cryptognathidae, 2 Raphignathidae and 2 Stigmaeidae. Remaining species were collected from vegetation surrounding citrus orchards: 1 Cunaxidae, 1 Eupalopsellidae, 1 Phytoseiidae, 2 Stigmaeidae, 3 Tenuipalpidae and 2 Tetranychidae.

Order Mesostigmata

Family Ameroseiidae

Epicroopsis horridus (Kramer, 1876)

Specimens collected — Al-ya'robiyah: Five females from litter and soil under *Eucalyptus* sp., 7 and 9 April, 2017. This is the second ameroseiid species recorded from Syria (i.e., *Ameroseius lidiae* Bregetova; Barbar 2016). *Epicroopsis horridus* occurs mostly in leaf litter and soil detritus and known from Europe and Asia (Hajizadeh *et al.*, 2013; Mašán, 2017). The biology of mite species belonging to the genus *Epicroopsis* largely unknown (Narita and Moraes, 2016).

Family Ascidae

Gamasellodes vulgator Athias-Henriot, 1961

Specimen collected — Al-ya'robiyah: One female from litter and soil under *Eucalyptus* sp., 30 March, 2017. This is the second ascid species from Syria (i.e., *Gamasellodes americanus* (Garman); Barbar, 2016). It was previously recorded from Algeria (Athias-Henriot, 1961) and India (Bhattacharyya and Sanyal, 2002). The biology of this species is unknown. However, several *Gamasellodes* species can feed and reproduce on several nematode groups in soil (Moraes *et al.*, 2015).

Family Laelapidae

This is the first record of this family from Syria. Two species were collected:

***Androlaelaps aegypticus* Hafez, El-Badry and Nasr, 1982**

Specimen collected — Al-ya'robiyah: One female from litter and soil under *Eucalyptus* sp., 30 March, 2017. This is the third international record of this species. It was previously recorded from Egypt (Hafez *et al.*, 1982) and Iran (Soleimani *et al.*, 2010). *Androlaelaps aegypticus* is a very active predator and can feed on immature stages of *Tyrophagus putrescentiae* (Schränk), *Rhizoglyphus echinopus* (Fumouze and Robin) (Acaridae) and *Lepidoglyphus destructor* (Schränk) (Glycyphagidae). Differences in biological characteristics (incubation period, life cycle and female fecundity) of this species were observed according to prey and temperatures (Mostafa *et al.*, 2017).

***Stratiolaelaps miles* (Berlese, 1892)**

Specimens collected — Al-ya'robiyah: Two females from litter and soil under *Eucalyptus* sp., 4 and 9 April, 2017. This species is widely distributed in the northern hemisphere and being used in the control of some glasshouse pests namely larvae of sciarid flies (Diptera: Sciaridae) and western flower thrips [*Frankliniella occidentalis* (Pergande)] (Gerson *et al.*, 2003; Walter and Campbell, 2003).

Family Phytoseiidae

This is the largest known family from Syria with 23 reported species (Barbar, 2017; Demite *et al.*, 2018). One additional phytoseiid species is recorded in the present study:

***Typhlodromus (Typhlodromus) ernesti* Ragusa and Swirski, 1978**

Specimens collected — Borj-Islam: Nineteen females and two males on *Pinus halepensis* Mill., 25 May, 2016. It was associated with the tenuipalpid species *Cenopalpus wainsteini* (Livschitz and Mitrofanov) in this study. This species is known from several countries in Europe and also from Tunisia but nothing is known about its biology (Demitte *et al.*, 2018).

Order Trombidiformes

Family Anystidae

This is the first record of this family from Syria. Two species were collected:

***Anystis wallacei* Otto, 1992**

Specimens collected — Al-ya'robiyah: One tritonymph and one deutonymph from litter under *Acacia cyanophylla* Lindley, 21 and 30 March, 2017, respectively. This species is described from pastures in France, Spain, Morocco and Australia and could be a natural enemy of penthaleid mite species and Collembola (Otto, 1992; Gerson *et al.*, 2003).

***Erythracarus pyrrholeucus* (Hermann, 1804)**

Specimens collected — Al-ya'robiyah: Four females from litter and soil under *A. cyanophylla*, 28 and 30 March, 2017. This species is recorded from Australia, Iran, Ukraine and the United States (Otto 1999). Small mite species and insects (i.e., Psocoptera) appear to be a possible food-source for species of *Erythracarus* including *E. pyrrholeucus* (Otto, 1999).

Family Cunaxidae

***Cunaxoides lootsi* Den Heyer, 2013**

Specimens collected — Attabiyyat: Six females on *Sarcopoterium spinosum* (L.), 15 March, 2018. This predatory mite is the fourth cunaxid species reported from Syria (Barbar, 2016). It

was only recorded from Iran on grass (Den Heyer *et al.*, 2013).

Family Cryptognathidae

This is the first record of this family from Syria. One species was collected:

***Favgognathus cordylus* Luxton, 1993**

Specimens collected — Al-ya'robiyah: One female from litter and soil under *A. cyanophylla*, 7 April, 2017. Nothing is known about the biology of this species. However, members of this family probably feed on the content of plant cells or on microorganisms in their habitats (Luxton, 1993; Swift and Goff, 2001).

Family Eupalopsellidae

This is the first record of this family from Syria. One species was collected:

***Eupalopsis maseriensis* (Canestrini and Fanzago, 1876)**

Specimen collected — Borj-Islam: One female on unidentified plant species, 25 May, 2016. This species preys on the citrus chaff scale *Parloria pergandii* Comstock. It was recorded from wild plant species and litter in several countries (i.e., Italy, USSR and Turkey) (Doğan, 2008; Meyer and Ueckermann, 1984).

Family Raphignathidae

Only one species of this family is known from Syria (i.e., *Raphignathus gracilis* (Rack); Barbar, 2016). Two additional species were collected in this study:

***Raphignathus hecmatanaensis* Khanjani and Ueckermann, 2003**

Specimens collected — Al-ya'robiyah: Two females from litter and soil under *A. cyanophylla*, 16 April, 2016 and 30 March, 2017. This species is known from Iran (Khanjani and Ueckermann, 2003) and Turkey (Bingül *et al.*, 2018).

***Raphignathus zhaoui* Hu, Jing and Liang, 1995**

Specimens collected — Al-ya'robiyah: Seven females from litter and soil under *A. cyanophylla*, 16 and 5 April, 2016 and 2017, respectively. This species was reported in China and Iran from cultivated and wild plant species, soil and moss (Fan and Yin, 2000; Khanjani and Ueckermann, 2003).

Family Stigmaeidae

Only one species of this family is known from Syria (i.e., *Agistemus exsertus* Gonzalez-Rodriguez; Barbar, 2016). Four additional species were collected in this study:

***Agistemus duzgunesae* Koç, Cobanoğlu and Madanlar, 2005**

Specimens collected — Borj-Islam: One female on *Prunus amygdalus* L. and one female on *Olea europaea* L., 25 May, 2016. It was associated with *Bryobia rubrioculus* (Scheuten), *Tetranychus urticae* Koch and *Brevipalpus oleae* Baker in this study. It was only reported from Turkey and Greece on several cultivated plant species and natural vegetation (Koç *et al.*, 2005; Stathakis *et al.*, 2014).

***Stigmaeus additicus* Dönel and Doğan, 2011**

Specimen collected — Al-ya'robiyah: One female from litter and soil under *A. cyanophylla*, 7 April, 2017. This is the second international record of this predatory mite. It was described from Turkey from litter under *Acacia* sp. and from moss and lichen on soil (Dönel and Doğan, 2011).

***Stigmaeus ayyildizi* Dönel and Doğan, 2011**

Specimens collected — Al-ya'robiyah: Three females from litter and soil under *A. cyanophylla*, 7 April, 2017. This is the second international record of this mite species. It was described from Turkey from grassy soil (Dönel and Doğan, 2011).

***Zetzellia graeciana* González-Rodríguez, 1965**

Specimens collected — Borj-Islam: Three females on *Punica granatum* L., 25 May, 2016. It was associated with *Brevipalpus obovatus* Donnadieu and *Tenuipalpus punicae* Pritchard and Baker in this study. This species is known from several European countries and was collected from apples, citrus and other plants (Fan *et al.* 2016). It preys on *Panonychus ulmi* Koch and *T. urticae* (Inserra, 1970 [1973]).

Family Tenuipalpidae

Five species of this family are known from Syria (Barbar, 2016). Three additional species were collected in this study:

***Brevipalpus oleae* Baker, 1949**

Specimens collected — Borj-Islam: Two females on *O. europaea*, 25 May, 2016. This species was associated with the phytoseiid, *Typhlodromus* (*Typhlodromus*) *athiasae* Porath and Swirski in this study. It was recorded from Morocco, Greece, Italy, Portugal and Tunisia (Castagnoli and Pegazzano, 1986; Chatti *et al.*, 2017).

***Cenopalpus lanceolatisetae* (Attiah, 1956)**

Specimens collected — Borj-Islam: Four females on *Crataegus monogyna* L., 25 May, 2016. This species was associated with the predator *T. (T.) athiasae* in this study. It was described from Egypt on several Rosaceae (i.e., *Prunus domestica* L., *Prunus armeniaca* L., *Pyrus malus* L. and *Pyrus communis* L.) and distributed in Western Palearctic (Mesa *et al.*, 2009; Khanjani *et al.*, 2012).

***Tenuipalpus caudatus* (Dugés, 1834)**

Specimens collected — Borj-Islam: Three females on *Cupressus sempervirens* L. and one female on *Pistacia lentiscus* L., 25 May, 2016. This species was associated with the predator *T. (T.) athiasae* in this study. It was described from France on *Viburnum tinus* L. and found also on other plant species in several countries (Mesa *et al.*, 2009).

Family Tetranychidae

Fifteen species of this family are known from Syria (Migeon and Dorkeld, 2006-2017; Zeity, 2017). Two additional species were collected in this study:

***Bryobia kissophila* Eyndhoven, 1955**

Specimens collected — Al-ya'robiyah: Six females and one larva on *Malva sylvestris* L., 16 April, 2016. This species was collected from three plant species particularly from *Hedera helix* L. in Europe and was also collected from other countries as South Africa, Australia and United States (Migeon and Dorkeld, 2006-2017).

***Pseudobryobia nikitensis* (?) (Livshitz and Mitrofanov, 1969)**

Specimens collected — Borj-Islam and Attabiyyat: Nine females on *S. spinosum*, 25 May, 2016 and 15 March, 2018.

This species was also previously recorded on *S. spinosum* in Ukraine (Livshits and Mitrofanov, 1969) and Italy (Vacante, 1983) and on *Silene dioica* (L.) (unknown locality, Migeon and Dorkeld, 2006-2017). The Syrian specimens are very close to those described by

Figure 1 Variations in the shape of dorsal body setae in the Syrian specimens of *Pseudobryobia nikitensis*: a – prodorsal setae; b – *sc1* seta; c – *c1* seta; d – *d1* seta.

Livshits and Mitrofanov (1969) and similar to those reported in the redescription of Vacante (1983), except for a few small morphological differences namely: **(1)** in Syrian specimens, setae *v1*–*v2* elongated, slender and serrate (Figure 1a); setae *sc1*–*sc2* and *c1*–*c3* subspatulate (Figure 1b, c); remaining dorsal body setae palmate serrate and almost rounded distally (Figure 1d), but dorsohysterosomal setae are subspatulate, lanceolate and more or less acute in the

description of Livshits and Mitrofanov (1969), and setae *d*, *e*, and *f* are subspatulate in the redescription of Vacante (1983) (according to Figure presented); (2) femur II with 9 (10) setae in Syrian specimens oppose to 8 in the description of Livshits and Mitrofanov (1969); genu III with 6 (7) setae in Syrian specimens oppose to 5 setae in the description of Livshits and Mitrofanov (1969).

Furthermore, the assignment of this species to the genus *Pseudobryobia* may be uncertain. Indeed, in the diagnoses proposed by the authors who worked on this genus (e.g. Meyer, 1987; Baker and Tuttle, 1994; Smiley and Baker, 1995), in addition to the absence of prodorsal lobes, it is mentioned that members of the fourth pair of dorsocentral setae (*fl* setae) are located in normal dorsal position (not marginal). However, this condition is neither met by the specimens of *P. nikitensis* described by Livshits and Mitrofanov (1969), Vacante (1983) nor by those reported in this study.

Acknowledgements

I would like to thank Edward A. Ueckermann, Gilberto J. de Moraes, Salih Doğan, Mohamed W. Negm, Grzegorz Gabryś, Qing-Hai Fan, Carlo Duso, Lucia Zappalà, Adel M. Mostafa and Hany M. El-Kawas for supplying papers concerning the description and biology of several species in this research. Thanks are due to Philippe Auger for valuable comments and suggestions during preparation of this paper.

References

- Athias-Henriot C. 1961. Mésostigmates (Urop. excl.) édaphiques Méditerranéens (Acaromorpha, Anactinotrichida) (collect. Prof. H. Franz et C. Athias-Henriot). Première Série. *Acarologia*, 3: 381-509.
- Attiah H.H. 1956. The genus *Brevipalpus* in Egypt (Acarina: Tenuipalpidae). *Bull. Soc. Entomol. Egypte*, 40: 433-477.
- Baker E.W. 1949. The genus *Brevipalpus* (Acarina: Pseudoleptidae). *Am. Midl. Nat.*, 42(2): 350-402. doi:10.2307/2422013
- Baker E.W., Tuttle D.M. 1994. A Guide to the Spider Mites (Tetranychidae) of the United States. West Bloomfield: Indira Publishing House. pp. 347.
- Barbar Z. 2016. The mite fauna (Acari) of two Syrian citrus orchards, with notes on their morphology and economic importance. *Syst. Appl. Acarol.*, 21 (8): 991-1008. doi:10.11158/saa.21.8.1
- Barbar Z. 2017. Evaluation of three pesticides against phytophagous mites and their impact on phytoseiid predator in an eggplant open-field. *Acarologia*, 57 (3): 529-539. doi:10.24349/acarologia/20174170
- Barbar Z. 2018. New records of three larval Erythraeidae (Acari: Trombidiformes) from Latakia, Syria. *Syrian J. Agric. Res.*, 5 (1): 183-190.
- Barbar Z., Ueckermann E.A. 2017. Two new species and a new record of Bdellidae (Acari: Trombidiformes) from Syria. *Acarologia*, 57 (4): 1089-1102. doi:10.24349/acarologia/20174221
- Berlese A. 1892. Acari, Myriopoda et Scorpiones hucusque in Italia reperta 63, 17 text pages+Plates 1-10 (reprint by Junk, The Hague, 1979).
- Bhattacharyya A.K.M., Sanyal A.K. 2002. Three new species and some new records of the genus *Gamasellodes* Athias-Henriot (Acarina: Ascidae) from India. *Acarologia*, 42: 229-238.
- Bingül M., Doğan S., Doğan S. 2018. Asymmetric variations in some species of the genus *Raphignathus* Duges (Acari: Raphignathidae). *Trakya Univ. J. Nat. Sci.*, 19 (1): 55-58. doi:10.23902/trkjnat.334190
- Bolland H.R., Gutierrez J., Flechtman C.H.W. 1998. World Catalogue of the Spider Mite Family (Acari: Tetranychidae). Leiden: Brill Academic Publishers. pp. 392.
- Boller E.F. 1984. Eine anfache Ausschwemm-Methode zur schellen Erfassung von Raumilben, Trips und anderen Kleinarthropoden im Weinbau. *Schweiz Zeitschrift für Obst-und Weinbau*, 120: 249-255.
- Canestrini G., Fanzago F. 1876. Nuovi Acari Italiani. *Acad. Sci. Veneto-Trent. Instr. Padua Atti.*, 5: 130-142.
- Castagnoli M., Pegazzano F. 1986. Acariens. In: *Traité d'Entomologie Oléicole*. Y. Arambourg (Ed.) Cons. Oleicole Int., Madrid: 303-336.
- Chant D.A., McMurtry J.A. 2007. Illustrated Keys and Diagnoses for the genera and subgenera of the Phytoseiidae of the World (Acari: Mesostigmata). West Bloomfield: Indira Publishing House. pp. 220.
- Chatti A., Kreiter S., Lebdi-Grissa K., Ksantini M. 2017. Phytophagous and predatory mites on olive trees in Tunisia. Catalogue, description of one new species and key for identification (Acari, Eriophyidae, Tetranychidae, Tenuipalpidae and Phytoseiidae). *Acarologia*, 57 (2): 233-254. doi.org/10.1051/acarologia/20164152
- Demite P.R., Moraes G.J. de, McMurtry J.A., Denmark H.A., Castilho R. de. C. 2018. Phytoseiidae Database. Available from <http://www.lea.esalq.usp.br/phytoseiidae> (accessed 7 April, 2018).

- Den Heyer J., Ueckermann E.A., Khanjani M. 2013. Iranian Cunaxidae (Acari: Prostigmata: Bdelloidea). Part III. Subfamily Cunaxoidinae. J. Nat. Hist., 47 (31-32): 2049-2070. doi:10.1080/00222933.2012.763060
- Doğan S. 2008. The first report of *Eupalopsis* (Acari: Eupalopsellidae) in Turkey: *Eupalopsis maseriensis* (Canestrini & Fanzago). Syst. Appl. Acarol., 13: 155-158. doi:10.11158/saa.13.2.10
- Dönel G., Doğan S. 2011. The stigmatid mites (Acari: Stigmatidae) of Kelkit Valley (Turkey). Zootaxa, 2942: 1-56.
- Dugés A. 1834. Recherches pour servir à l'ordre des Acariens en général et la famille des Trombidies en particulier. Ann. Sci. Nat. Paris (ser. 2), 1: 5-46.
- Evans G.O., Till W.M. 1979. Mesostigmatic mites of Britain and Ireland (Chelicerata: Acari-Parasitiformes). An introduction to their external morphology and classification. Trans. Zool. Soc. Lond., 35: 139-270. doi:10.1111/j.1096-3642.1979.tb00059.x
- Fan Q.-H., Yin X.-M. 2000. The genus *Raphignathus* (Acari: Raphignathidae) from China. Syst. Appl. Acarol., 5: 83-98. doi: 10.11158/saa.5.1.11 doi:10.11158/saa.5.1.11
- Fan Q.-H., Zhang Z.-Q. 2005. Raphignathoidea (Acari: Prostigmata). Fauna N. Z., 52: 1-400.
- Fan Q.-H., Flechtman C.H.W., Moraes G.J.de. 2016. Annotated catalogue of Stigmatidae (Acari: Prostigmata), with a pictorial key to genera. Zootaxa, 4176 (1): 001-199. doi:10.11646/zootaxa.4176.1.1
- Gerson U.L., Smiley R., Ochoa R. 2003. Mites (Acari) for Pest Control. Malden: Blackwell Science Ltd. 539 pp. doi:10.1002/9780470750995
- González-Rodríguez R.H. 1965. A taxonomic study of the genera *Mediolata*, *Zetzellia* and *Agistemus* (Acarina: Stigmatidae). Univ. Calif. Publ. Entomol., 41: 1-64.
- Hafez S.M., El-Badry E.A., Nasr A.K. 1982. Soil mites of the family Laelapidae from Egypt. Res. Bull., Fac. Agric., Ain Shams Univ., 1759: 1-15.
- Hajizadeh J., Tajmiri P., Maśán P. 2013. Redescription of *Ameroseius lanceosetis* Livshitz & Mitrofanov, 1975 (Acari: Mesostigmata), with a checklist and a key to the ameroseiid mites of Iran. Int. J. Acarol., 39: 146-152. doi:10.1080/01647954.2012.747566
- Hermann J.F. 1804. Mémoire Aptérologique (Strasbourg: F. -L. Hammer).
- Hu C. Y., Jing Z. Q., Liang L. R. 1995. Two new species and one new record of the genus *Raphignathus* Dugés (Acari: Raphignathidae). J. Suzhou Railw. Teach. Coll., 12(3): 21-26. [in Chinese with English abstract].
- Insera R. 1970 [1973]. Morphological observation and biological notes on *Zetzellia graeciana* Gonzalez (Acarina: Stigmatidae). Boll. Zool. Agrar. Bachic., 10: 85-119.
- Khanjani M., Ueckermann E.A. 2003. Two new species of the genus *Raphignathus* Dugés (Acari: Raphignathidae) from Iran. Acarologia, 43: 299-306.
- Khanjani M., Ueckermann E.A. 2008. New species of the genus *Favognathus* Luxton (Acari: Cryptognathidae). Acarologia, 48 (3-4): 177-186.
- Khanjani M., Khanjani M., Saboori A., Seeman O. 2012. The false spider mites of the genus *Cenopalpus* Pritchard & Baker (Acari: Tenuipalpidae) from Iran. Zootaxa, 3433: 1-59.
- Koç K., Cobanoğlu S., Madanlar N. 2005. *Agistemus duzgunesae* sp. n. (Acari, Stigmatidae) from Turkey. Biol. Bratisl., 60 (2): 121-124.
- Kramer P. 1876. Zur Naturgeschichte einiger Gattungen aus der Familie der Gamasiden. Archiv für Naturgeschichte, 42: 46-105.
- Krantz G.W., Walter D.E. 2009. A Manual of Acarology. Third Edition. Lubbock: Texas Tech University Press. pp. 807.
- Livshits I.Z., Mitrofanov V.I. 1969. New species of the genus *Bryobia* from the Crimea. Bjul. Gos Nikit. Bot. Sada, 4: 59.
- Luxton M. 1993. New species of cryptognathid mites from Israel (Acari: Prostigmata: Cryptognathidae). J. Nat. Hist., 27: 1213-1217. doi:10.1080/00222939300770711
- Maśán P. 2017. A revision of the family Ameroseiidae (Acari, Mesostigmata), with some data on Slovak fauna. ZooKeys, 704: 1-228. doi:10.3897/zookeys.704.13304
- Mesa N.C., Ochoa R., Welbourn W.C., Evans G.A., Moraes G.J. de 2009. A catalog of the Tenuipalpidae (Acari) of the world with a key to genera. Zootaxa, 2098: 1-185.
- Meyer M.K.P. 1979. The Tenuipalpidae (Acari) of Africa with keys to the world fauna. Entomol. Mem. Dep. Agric. Repub. S. Afr., 50: 1-133.
- Meyer M.K.P.S. 1987. African Tetranychidae (Acari: Prostigmata) with reference to the world genera. Entomol. Mem. Dep. Agric. Water Supply Repub. S. Afr., 69: 1-175.
- Meyer M.K.P., Ueckermann E.A. 1984. The family Eupalopsellidae (Acari: Prostigmata) with descriptions of new species from South Africa. Phytophylactica, 16: 121-142.
- Meyer M.K.P., Ueckermann E.A. 1987. A taxonomic study of some Anystidae (Acari: Prostigmata). Entomol. Mem. Dep. Agric. Water supply, Plant Prot. Res. Inst., Repub. S. Afr., 68: 1-37.
- Migeon A., Dorkeld F. 2006-2017. Spider Mites Web: a comprehensive database for the Tetranychidae. Available from: <http://www.montpellier.inra.fr/CBGP/spmweb> (accessed September 15, 2017).
- Moraes G.J. de, Venancio R., Santos V.L.V. dos, Paschoal A.D. 2015. Potential of Ascidae, Blattisociidae and Melicharidae (Acari: Mesostigmata) as biological control agents of pest organisms. In: Carrillo D., Moraes G.J. de, Pea J.E. (Eds.). Prospects for Biological Control of Plant Feeding Mites and Other Harmful Organisms —Heidelberg: Springer International Publishing, pp. 33-75. doi:10.1007/978-3-319-15042-0_2
- Mostafa A.M., Sakr H.H., Yassin E.M.A., Abd El-Khalik A.R. 2017. Laboratory studies on the mesostigmatid mites *Androlaelaps aegypticus* (Laelapidae) and *Proctolaelaps gizanensis* (Ascidae) on three mite pests at different conditions. Egypt. J. Acad. Boil. Sci., 10 (4): 63-70. doi: 10.21608/eajbsa.2017.12485

- Narita J.P.Z., Moraes G.J. de 2016. A new species of *Epicriopsis* Berlese (Acari: Mesostigmata: Ameroseiidae) from Brazil, with a key to the world species of the genus. *Zootaxa*, 4114: 477-484. doi:10.11646/zootaxa.4114.4.7
- Otto J.C. 1992. A new species of *Anystis* Von Heyden compared with *Anystis salicinus* (Linnaeus) (Acarina: Anystidae). *Int. J. Acarol.*, 18 (1): 25-35. doi:10.1080/01647959208683925
- Otto J.C. 1999. Revision of the genus *Erythracarus* Berlese (Acarina: Anystidae Erythracarinae). *J. Nat. Hist.*, 33: 825-909. doi:10.1080/002229399300146
- Ragusa S., Swirski E. 1978. Description of three new species of *Typhlodromus* Scheuten from Italy with redescription of *Typhlodromus baccettii* Lombardini (Acari: Phytoseiidae). *Int. J. Acarol.*, 4: 211-220. doi:10.1080/01647957808683118
- Skvarla M.J., Fisher J.R., Dowling A.P.G. 2014. A review of Cunaxidae (Acariformes, Trombidiformes): Histories and diagnoses of subfamilies and genera, keys to world species, and some new locality records. *ZooKeys*, 418: 1-103. doi:10.3897/zookeys.418.7629
- Smiley R.L., Baker E.W. 1995. A report on some tetranychid mites (Acari: Prostigmata) from Yemen. *Int. J. Acarol.*, 21: 135-164. doi:10.1080/01647959508684055
- Soleimani M., Ostovan H., Joharchi O. 2010. First report of the mite species *Androlaelaps aegypticus* (Mesostigmata: Laelapidae) from Iran. *Plant Prot. J.*, 2 (2): 157-160.
- Stathakis T.I., Kapaxidi E.V., Papadoulis G.Th. 2014. A new record and a new species of the genus *Agistemus* Summers (Acari: Stigmaeidae) from Greece. *Zootaxa*, 3780 (1): 153-170. doi:10.11646/zootaxa.3780.1.6
- Swift S.F., Goff M.L. 2001. Mite (Acari) communities associated with 'Ohi'a, *Metrosideros polymorpha* (Myrtaceae), at Hono O NalPali and Kui'a Natural Area Reserves on Kaua'I Island, Hawaiian Islands. *Pac. Sci.*, 55: 23-40. doi:10.1353/psc.2001.0008
- Syrian Ministry of Agriculture and Agrarian Reform, 2016. www.moaar.gov.sy (accessed March 15, 2018).
- Vacante V. 1983. Prima raccolta di AcariTetranichidi in Sicilia. *Phytophaga*, 1: 41-114.
- Walter D.E., Campbell N.J.H. 2003. Exotic vs endemic biocontrol agents: would the real *Stratiolaelaps* miles (Berlese) (Acari: Mesostigmata: Laelapidae), please stand up?. *Biol. Control*, 26: 253-269. doi:10.1016/S1049-9644(02)00171-8
- Zeity M. 2017. Some new records of spider mites (Acari, Tetranychidae) from Syria. *Acarologia*, 57(3): 651-654. doi:10.24349/acarologia/20174184
- Zriki G., Shaabo A., Boubou A. 2015. A preliminary survey of the spider mites (Acari: Tetranychidae) in Latakia Governorate of Syria. *Acarologia*, 55: 303-309. doi.org/10.1051/acarologia/20142173