Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Supporting agricultural research
for sustainable development

Acarologia is under **free license** and distributed under the terms of the
Creative Commons-BY

Feather mites (Acari: Astigmata) of the Zoobotanical State Park in Teresina, Brazil

Maria Marta Lima Alves Moro^a, Eric Carvalho Waquim^b, Luanna Soares de Melo Evangelista^c, Fabio Akashi Hernandez^d

^a Department of Biology, Center for Nature Sciences, Federal University of Piauí (UFPI), Teresina, Piauí, Brazil.

^b Academic Veterinary Medicine, Center of Agrarian Sciences, Federal University of Piauí (UFPI), Teresina, Piauí, Brazil.

^c Department of Parasitology and Microbiology, Health Sciences Center, Federal University of Piauí (UFPI), Teresina, Piauí, Brazil.

^d Department of Ecology and Zoology, Center of Biological Sciences, Federal University of Santa Catarina, Florianópolis (UFSC), Santa Catarina, Brazil.

Original research

ABSTRACT

Feather mites are the most diverse avian ectoparasites. They mainly live on the plumage of birds, feeding on uropygial oil, and may occasionally cause skin irritation, especially on caged birds. Here we report the results of a survey of feather mites on wild birds of the Zoobotanical State Park of Teresina, Piauí, Brazil. The following mites were collected from seven bird specimens kept in the park from January to July 2017: *Pandionacarus fuscus* (Nitzsch, 1818) on the osprey (*Pandion haliaetus*); *Freyana dendrocygni* Dubinin, 1950 and *Heterobrepheosceles megathrix* Peterson & Atyeo, 1977 on the white-faced whistling-duck (*Dendrocygna viduata*); *Nyctibiolichus megamerus* Atyeo, 1979 on the common potoo (*Nyctibius griseus*); *Hieracolicus hirundo* (Mégnin & Trouessart, 1884) on the harpy eagle (*Harpia harpyja*) and *Hieracolicus* sp. on the southern caracara (*Caracara plancus*). These findings increase the known ranges of the mite species, with four of them reported for the first time in Brazil.

Keywords birds; ectoparasitism; feather mites; record; captivity; *Psoroptidia*

Introduction

Feather mites (Arthropoda: Chelicerata: Arachnida: Acariformes: Astigmata) are the most diverse and abundant permanent avian ectosymbionts, associated with nearly all bird orders (Gaud and Atyeo, 1996). Most feather mites exhibit morphological adaptations to inhabit specific microhabitats on their hosts' body – flight feathers of the wings, tail, down, and contour feathers, quills and skin; some members of the Pyroglyphidae (Analgoidea) also inhabit nests (Dabert and Mironov, 1999).

While a number of feather mite lineages quite certainly evolved by coevolution with their hosts, others have successfully colonized new hosts by horizontal transfer (Doña et al., 2018). Therefore, the understanding of the origin of mite-bird associations is intrinsically tied to the knowledge about bird phylogeny, ecology and behavior (Proctor and Owen, 2000). Also, mites can affect bird reproductive success and exposure them to diseases, especially in birds held in captivity (Atyeo and Gaud, 1987; Mironov, 2013; pers. obs. by FAH).

Given the exuberant diversity of birds found in the Neotropical region, it is not surprising that the study of feather mites on birds of Brazil is still incomplete, which renders a huge potential for taxonomic discoveries in this country. Researches involving these organisms,

Received 13 August 2019
Accepted 17 September 2019
Published 22 October 2019

Corresponding author
Fabio Akashi Hernandez:
abakashi@gmail.com

Academic editor
Auger, Philippe

DOI
10.24349/acarologia/20194345

ISSN 0044-586X (print)
ISSN 2107-7207 (electronic)

© Copyright
Lima Alves Moro M. M. et al.

Distributed under
Creative Commons CC-BY 4.0

as well as training of students, ornithologists and parasitologists, have been improving the understanding of the ecological and biological associations of these mites with their hosts in Brazil (Roda and Farias, 2007; Pedroso and Hernandes, 2016).

The aim of this work was to investigate the feather mites associated with some wild birds of the Zoobotanical State Park of Teresina, Piauí State, Brazil.

Materials and Methods

This work was evaluated and approved by the Ethics Committee in Animal Experimentation of the Federal University of Piauí (UFPI), (protocol 242/2016). The material was collected at the Zoobotanical State Park, in an urban area of the municipality of Teresina, Piauí. The park was created in 1973 and has approximately 137 hectares, being an area designated for the protection, conservation, and researches of fauna and flora, governed by the public administration of the State Secretariat for the Environment and Water Resources of Piauí.

Feather mites were collected from wild birds held in captivity, from January to July of 2017. Mites were collected manually from the feathers using fine brushes and tweezers; a few feathers were partially cut and only one *post-mortem* bird – an osprey, *Pandion haliaetus* (Linnaeus, 1758) – was defrosted and washed for ectosymbionts. This was done by washing the bird with water and detergent (Mironov & Galloway 2002), and the resulting liquid was filtered in paper filters; the mites were collected with a fine brush from under a dissecting scope. The number of examined feathers for each bird species is shown in Table 1.

Table 1 Birds examined for feather mites at the Zoobotanical State Park, Teresina, PI, Brazil, 2017.

Birds	EB	IB	EF	Feather mites
<i>Pandion haliaetus</i> (Accipitriformes: Pandionidae)	1	1	12	<i>Pandionacarus fuscus</i> * (Avenzoariidae)
<i>Harpia harpyja</i> (Accipitriformes: Accipitridae)	1	1	2	<i>Hieracolicus hirundo</i> ** (Gabuciniidae)
<i>Caracara plancus</i> (Falconiformes: Falconidae)	2	2	8	<i>Hieracolicus</i> sp.** (Gabuciniidae)
<i>Dendrocygna viduata</i> (Anseriformes: Anatidae)	5	2	17	<i>Freyana dendrocygni</i> * (Freyanidae) <i>Heterobrephosceles megathrix</i> * (Alloptidae)
<i>Nyctibius griseus</i> (Nyctibiiformes: Nyctibiidae)	2	1	5	<i>Nyctibiolichus megamerus</i> * (Pterolichidae)
Total	11	7	44	

Legend: EB – number of examined birds; IB – number of infested birds; EF – number of examined feathers for each bird specimen; *First record of the mite species in Brazil; **species recorded in a new locality in Brazil.

After collecting, the mites were counted and stored in vials with 70% ethanol, labeled with locality, date and host. The sorting was done in the Laboratory of Parasitology of the Department of Parasitology and Microbiology of UFPI, and the material was sent to FAH for mounting and identification. Mites were mounted in slides with Hoyer's medium, and subsequently analyzed and photographed on a Leica DM3000 microscope equipped with a DFC295 digital camera. Debris surrounding the specimens were digitally removed using Adobe Photoshop CS6. Voucher specimens were deposited at the Acari collection of the Universidade Federal de Santa Catarina. The classification of birds follows BirdLife International (2018).

Results

All examined bird species were shown to harbor feather mites. We found feather mites on 7 of the 11 bird specimens that we examined. The following mite species were recovered from their respective hosts: *Pandionacarus fuscus* (Nitzsch, 1818) (Avenzoariidae) from the osprey *Pandion haliaetus* (Pandionidae); *Freyana dendrocygni* Dubinin, 1950 (Freyanidae) and *Heterobrephosceles megathrix* Peterson & Atyeo, 1977 (Alloptidae) on the white-faced whistling duck (*Dendrocygna viduata*) (Linnaeus, 1766) (Anatidae); *Nyctibiolichus megamerus* Atyeo, 1979 (Pterolichidae) on the common potoo (*Nyctibius griseus*) (Gmelin, 1789) (Nyctibiidae); *Hieracolichus hirundo* (Méglin & Trouessart, 1884) on the harpy eagle (*Harpia harpyja*) (Linnaeus, 1758) (Accipitridae), and *Hieracolichus* sp. (Gabuciniidae) on the southern caracara (*Caracara plancus*) (Miller, 1777) (Falconidae) (Table 1).

Discussion

Besides reporting feather mites for the first time from the Northern Brazilian state of Piauí, this study also reports four species for the first time in Brazil, and two mites constitute new locality records. *Pandionacarus fuscus* (Figure 1), one of the largest feather mites at about 1.2–1.3 mm in length, and commonly associated with the osprey (Miller *et al.*, 1997; Philips, 2000), was recently reported on a toucan – *Ramphastos citreolaemus* Gould, 1844 (Piciformes: Ramphastidae) – in Colombia (Parra-Henao *et al.*, 2011), that is definitely a contamination. The osprey is a worldwide distributed bird, including in several parts of Brazil (BirdLife International, 2016), and we herein report *Pandionacarus fuscus* for the first time in this country.

Dendrocygna viduata was reported as a host of *Freyana dendrocygni* (Figure 2), along with *D. javanica*, *D. arcuata*, *D. eytoni*, and *D. bicolor* (McDaniel *et al.*, 1966). Mites of this genus are normally associated with aquatic birds of the family Anatidae (ducks, geese, swans) (Dubinin, 1950; Aksin, 2007). There are records of the mite *Heterobrephosceles megathrix* (Figures 3A–C) from lesser whistling ducks in Malaysia and Thailand, and white-faced whistling ducks in Uganda (Peterson and Atyeo, 1977). Gaud (1952: 87) reported undetermined alloptid females on the latter host, but he asserted that the absence of males could not allow a reliable identification of the genus. The females of *H. megathrix* remained unknown until this date, and were found herein for the first time. *Freyana dendrocygni* and *Heterobrephosceles megathrix*, both found on *Dendrocygna viduata*, are recorded for the first time in Brazil.

Nyctibiolichus megamerus (Figure 3D), one of the two known species of the genus *Nyctibiolichus* Atyeo, was described from *Nyctibius griseus* in Panama and Jamaica (Atyeo, 1979). Up to this date, this mite had not been reported from hosts or countries other than those in the original publication. In our study, only a few females and immatures, and no males of this mite were recovered, that can be due to the low number of feathers collected. Despite about 25 caprimulgid species occurring in Brazil, feather mites had been previously reported from only three host species in that country, which suggests the existence of a number of undescribed taxa (Mironov and Fain, 2003; Enout *et al.*, 2012; Hernandez, 2014; Pedroso and Hernandez, 2016).

Figure 1 *Pandionacarus fuscus* found on the osprey (*Pandion haliaetus*): A – male; B – female. Scalebar A, B = 100 μ m.

Figure 2 *Freyana dendrocygni* found on the white-faced whistling duck (*Dendrocygna viduata*): A – male, B – female. Scalebar A–B = 100 μm .

The genus *Hieracolicus* includes nine described species primarily associated with birds of the family Accipitridae (Accipitriformes) (Gaud and Atyeo, 1974; Mironov *et al.*, 2007). Nevertheless, a few studies (Barreto *et al.*, 2012; Pereira, 2015, present study) indicate that this genus is also associated with falcons (Falconiformes), which are presently considered to be distant relatives to hawks and eagles (Prum *et al.*, 2015).

Feather mites studied from birds held in captivity can constitute an important resource for understanding patterns of bird-mite associations. For instance, they can help to confirm poorly known associations. Until recently, *Hieracolicus hirundo* (Figure 4) was represented only by a few specimens in a single slide at the Museum National d'Histoire Naturelle, Paris. Based on those and also on freshly collected specimens from a harpy eagle in captivity, this species was recently redescribed (Hernandes, 2017a).

However, a caveat must be made to unsuspecting students or researchers studying feather mites from birds held in captivity: it is not uncommon to recover unnatural mite-bird associations, because birds can acquire mites from other birds held in the same or neighboring cages. Thus, rheas – *Rhea americana* (Linnaeus, 1758) (Rheiformes: Rheidae) – were once reported as hosts of the same mite species commonly associated with ostriches – *Struthio camelus* (Linnaeus, 1758) (Struthioniformes: Struthionidae) (Méglin and Trouessart, 1884; André, 1960), and flamingos (Phoenicopteriformes: Phoenicopteridae) may eventually show mites commonly found on ducks (Anseriformes: Anatidae) (pers. obs. by FAH). These are obvious contaminations.

Figure 3 *Heterobrepheosceles megathrix* (A–C) found on the white-faced whistling duck (*Dendrocygna viduata*): A – male, B – tritonymph (A and B attached in a precopulatory guarding position), C – female; *Nyctibiolichus megamerus* (D) found on the common potoo *Nyctibius griseus*: D – female. Scalebar A-D = 100 μm.

Figure 4 *Hieracolicus hirundo* found on the harpy eagle (*Harpia harpyja*): A – male; B – female. Scalebar A-B = 100 μm.

All mites reported herein, however, are natural associates of their respective hosts, for they have been originally reported from the same hosts in other localities. This fact may help to reinforce the knowledge about these mite-bird associations. Mironov *et al.* (2018) demonstrated that feather mites may persist on a host in captivity for more than 40 years after being captured from the wild, even after undergoing through antiparasitic treatments. Likewise, cockatiels, *Nymphicus hollandicus* (Kerr, 1792) (Psittaciformes: Cacatuidae), bred and kept in captivity for decades in Brazil – and often treated for parasites in pet stores, were also shown to contain feather mites from its place of origin, Australia (Hernandes, 2017b).

The studies on feather mites collected from birds kept in captivity are still very scarce in Brazil (Valim *et al.*, 2006; Pereira *et al.*, 2018). We hope to stimulate further studies since valuable results may be obtained from these birds, including confirmation of parasite-host associations, discovery of new species, recollecting of rare species, new records and hosts, information on microhabitats, and also provide a better understanding on epidemiological case studies.

Acknowledgements

The authors are thankful to the administration staff of the Zoobotanical State Park of Teresina, Piauí, for granting the opportunity to perform this study.

References

- Aksin N. 2007. *Freyana anatina* (Koch, 1844) Feather Mites (Acarina, Freyanoidea) Recorded for the First Time on Wild Ducks (Subfamily, Anatinae) in Turkey. *Türkiye Parazit Derg.* 31(4): 302-305.
- André M. 1960. Sarcopitides plumicoles parasites des autruches. *Acarologia*, 2(4): 556-567.
- Atyeo W.T. 1979. New feather mite taxa (Acarina: Pterolichoidea) from the avian order caprimulgiformes). *J. Med. Entomol.* 16(6): 528-535. doi:10.1093/jmedent/16.6.528
- Atyeo W.T., Gaud J. 1987. Feather mites (Acarina) of the parakeet, *Melopsittacus undulatus* (Shaw) (Aves: Psittacidae). *J. Parasitol.*, 73: 203-6. doi:10.2307/3282367
- Barreto M., Burbano M.E., Proctor H.C., Mironov S.V., Wauthy G. 2012. Feather mites (Acariformes: Psoroptidia) from Colombia: preliminary list with new records. *Zootaxa*, 3516: 1-68. doi:10.11646/zootaxa.3516.1.1
- BirdLife International 2016. *Pandion haliaetus*. The IUCN Red List of Threatened Species 2016: e.T22694938A93478747. Accessed 23 July 2019 doi:10.2305/IUCN.UK.2016-3.RLTS.T22694938A93478747.en
- BirdLife International 2018. digital checklist of the birds of the world. Version 3. Available at: http://datazone.birdlife.org/userfiles/file/Species/Taxonomy/HBW-BirdLife_Checklist_v3_Nov18.zip. Accessed 22 July 2019.
- Dabert J., Mironov S.V. 1999. Origin and evolution of feather mites (Astigmata). *Exp. Appl. Acarol.*, 23: 437-454. doi:10.1023/A:1006180705101
- Doña J., Proctor H., Mironov S., Serrano D., Jovani R. 2018. Host specificity, infrequent major host switching and the diversification of highly host-specific symbionts: The case of vane-dwelling feather mites. *Global Ecol. Biogeogr.*, 27: 188-198. doi:10.1111/geb.12680
- Dubinin V.B. 1950. Systematical analysis of species of feather mites (Sarcopitiformes, Analgesoidea) parasitic on Anseriformes. *Parasitol. Sb.*, 12: 17-72.
- Enout A.M.J., Lobato D.N.C., Diniz F.C., Antonini, Y. 2012. Chewing lice (Insecta, Phthiraptera) and feather mites (Acari, Astigmata) associated with birds of the Cerrado in Central Brazil. *Parasitol. Res.*, 111: 1731-1742. doi:10.1007/s00436-012-3016-5
- Gaud J. 1952. Sarcopitides plumicoles des oiseaux de Madagascar. *Mem. Inst. sci. Madagascar, Ser. A*, 7(1): 81-107.
- Gaud J., Atyeo W.T. 1974. Gabuciniidae, famille nouvelle de Sarcopitiformes plumicoles. *Acarologia*, 16(3): 522-561.
- Gaud J., Atyeo W.T. 1996. Feather mites of the World (Acarina, Astigmata): the supraspecific taxa. *Ann. Mus. R. Afr. Centr. Sci. Zool.*, 277: 1-193 [text], 1-436 [illustrations].
- Hernandes F.A. 2014. The feather mites of nightjars (Aves: Caprimulgidae), with descriptions of two new species from Brazil (Acari: Xolalgidae, Gabuciniidae). *Folia Parasit.*, 61(2): 173-181. doi:10.14411/fp.2014.024
- Hernandes F.A. 2017a. Rediscovery and redescription of *Hieracolicus hirundo* (Mégnin & Trouesart, 1884) (Astigmata: Gabuciniidae) from the Harpy Eagle *Harpya harpyja* (Linnaeus, 1758) (Accipitriformes: Accipitridae). *Syst. Appl. Acarol.*, 22(4): 509-517. doi:10.11158/saa.22.4.7
- Hernandes F.A. 2017b. *Psittophagus hollandicus* n. sp., a new feather mite species (Acariformes: Pterolichidae) from the cockatiel *Nymphicus hollandicus* (Kerr, 1792) (Psittaciformes: Cacatuidae) in Brazil. *Acarologia*, 57(4): 893-900. doi:10.24349/acarologia/20174215

- McDaniel B., Tuff D., Bolen E. 1966. External parasites of the black-bellied tree duck and other Dendrocygnids. *Wildlife Found.*, 78(4): 462-468.
- Mégnin P., Trouessart E.L. 1884. Les sarcoptides plumicoles. Révision du groupe des Analgesinae, et description des espèces et genres nouveaux de la collection du Musée d'Angers (Suite 1). *Jl. Micrographie*, 8: 257-266; 331-338.
- Miller M.J., Ewins P.J., Galloway T.D. 1997. Records of ectoparasites collected on ospreys from Ontario. *J. Wildl. Dis.*, 33:373-376. doi:10.7589/0090-3558-33.2.373
- Mironov S.V. 2013. *Allopsoroptoides galli* n. g., n. sp., a new genus and species of feather mites (Acari: Analgoidea: Psoroptoididae) causing mange in commercially raised domestic chicken in Brazil. *Syst. Parasitol.*, 85: 201-212. doi:10.1007/s11230-013-9422-y
- Mironov S.V., Fain, A. 2003. New feather mites of the family Ascouracaridae (Astigmata: Pterolichoidea) from some parrots and nightjars. *Acarologia*, 43(1): 99-111.
- Mironov S.V., Galloway, T.D. 2002. New feather mite taxa (Acari: Analgoidea) and mites collected from native and introduced birds of New Zealand. *Acarologia*, 42(1): 185-201.
- Mironov S.V., Proctor H.C., Barreto M., Zimmerman G. 2007. New genera and species of feather mites of the family Gabuciniidae (Astigmata: Pterolichoidea) from New World raptors (Aves: Falconiformes). *Can. Entomol.*, 139: 757-777. doi:10.4039/n07-001
- Mironov S.V., Efeykin B.D., Ibanez J.C., Sumayae A.M., Tolstenkov O.O. 2018. Captive individuals of endangered Philippine raptors maintain native feather mites (Acariformes: Pterolichoidea) species. *Int. J. Parasitol.-Par.*, 7: 116-133. doi:10.1016/j.jippaw.2018.03.002
- Parra-Henao G., Pineda E.P.A., Valencia G.L., Monroy D.M.R., Crespo G.J. 2011. Detection of ectoparasites in wild birds evaluated in Medellín (Colombia). *Rev. Col. Cienc. Pec.*, 24: 29-37.
- Pedroso L.G.A., Hernandez F.A. 2016. New records of feather mites (Acariformes: Astigmata) from nonpasserine birds (Aves) in Brazil. *Checklist*, 12(6): 1-25. doi: 10.15560/12.6.2000 doi:10.15560/12.6.2000
- Pereira D.M. 2015. Ácaros plumícolas de aves silvestres encaminhadas ao Centro de triagem de animais silvestres do Maranhão (CETAS-MA). 60p. Masters Science Dissertation - Universidade Estadual do Maranhão, São Luís, Maranhão.
- Pereira D.M., Hernandez F.A., Santos A.C.G., Nogueira R.M.S. 2018. Feather mites (Acari: Astigmata) of captive Psittaciformes in Brazil. *Arq. Bras. Med. Vet.*, 70(3): 843-849. doi:10.1590/1678-4162-9815
- Peterson P.C., Atyeo W.T. 1977. The feather mite family Alloptidae Gaud IV - A new genus of the Alloptinae (Acarina, Analgoidea). *Acarologia*, 19(2): 318-330.
- Philips J.R. 2000. A review and checklist of the parasitic mites (Acarina) of the Falconiformes and Strigiformes. *J. Raptor Res.*, 34(3): 210-231.
- Proctor H., Owens I. 2000. Mites and birds: diversity, parasitism and coevolution. *Trends Ecol. Evol.*, 15: 358-364. doi:10.1016/S0169-5347(00)01924-8
- Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Lemmon E.M., Lemmon A.R. 2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing. *Nature*, 526: 569-573. doi:10.1038/nature15697
- Roda A.S., Farias A.M.I. 2007. Ácaros plumícolas em beija-flores no município de Vicência, Pernambuco, Brasil. *Lundiana*, 8(1): 13-16.
- Valim M.P., Teixeira R.H.F., Gazeta G.S., Serra-Freire N.M. 2006. Duas espécies de cisnes (Aves: Anatidae) de cativeiro como novos hospedeiros para *Brephosceles discidicus* Peterson, 1971 (Acaridida: Pterolichidae). *Lundiana*, 7(2): 141-143.