Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Supporting agricultural research
for sustainable development

**Acarologia is under free license and distributed under the terms of the
Creative Commons-BY**

Occigamasus, a new genus of pergamasine mites, with description of two new species from the west coast of North America (Parasitiformes: Gamasina: Parasitidae)

Ilinca Juvara-Balş^a

^a Muséum d'histoire naturelle, CP 6434, Genève 6. Switzerland.

Original research

ABSTRACT

Occigamasus n. gen., *O. lindquisti* n. sp. and *O. makarovae* n. sp. from Vancouver Island (Canada) and Oregon (U.S.A.) are described. Five other new species and their sites are noted but not enough specimens were available for adequate descriptions. *Cycetogamasus californicus* (Banks) sensu Hennessey and Farrier, 1988 is transferred to *Occigamasus* as a new combination.

Keywords Gamasina; new genus; Parasitidae; Pergamasinae

Zoobank <http://zoobank.org/D9D1B3C7-98C9-4284-9F53-8D451D4746E5>

Introduction

Predatory mites of the Family Parasitidae Oudemans, 1901 are important soil and litter inhabitants in the Holarctic region. The family was divided into two subfamilies Parasitinae Oudemans, 1901 and Pergamasinae Juvara-Balş, 1972. Various studies of parasitid mites have been done in Europe (Berlese 1906; Bhattacharyya 1963; Athias-Henriot 1967a, 1967b, 1967c, 1971; Micherdzinsky 1969, Hyatt 1980). However, the diversity of these mites in North America has been scarcely investigated with the most important works being those of Banks (1904, 1910) and Ewing (1909).

Hennessey and Farrier (1988, 1989) re-examined and further described and illustrated the species of Parasitidae described by Banks and Ewing. Initially a systematic revision of thirty species in ten families of gamasine mites was published (1988), including seven species in six genera of Parasitidae, followed by a paper concerning only the family Parasitidae (1989) from North and South Carolina. In the latter paper, fourteen species, including one new genus and four new species were described, and new synonymies, new combination and a lectotype designated.

Scant information is available about parasitid mites in Canada. Richards (1976) published a paper concerning Parasitidae mites associated with bumblebees in Alberta. An estimated 35 species representing 6 genera of Parasitidae were recorded, without attributed names, as occurring in Canada (Lindquist *et al.* 1979); and that number of species was increased to 51, with almost twice as many species estimated to await discovery (Beaulieu *et al.* 2019). This paper deals with Pergamasinae mites from North America: Vancouver Island, British Columbia and Alberta (Canada) as well as Oregon (USA). Here, I describe an unusual new genus, *Occigamasus* and two new species, *O. lindquisti* and *O. makarovae*. The sites of other undescribed species belonging to the new genus, which need more specimens to be described adequately, are indicated.

Received 25 March 2019
Accepted 10 December 2019
Published 13 December 2019

Corresponding author
Ilinca Juvara-Balş: ibals@bluewin.ch

Academic editor
Faraji, Farid

DOI
10.24349/acarologia/20194354

ISSN 0044-586X (print)
ISSN 2107-7207 (electronic)

© Copyright
Juvara-Balş I.

Distributed under
Creative Commons CC-BY 4.0

Material and Methods

The mites studied (52 slides) are from the Canadian National Collection (CNC) deposited in the Research and Development center, Agriculture and Agri-Food, Ottawa. E. E. Lindquist collected most of the mites from Vancouver Island and Oregon. L. Russel collected one sample from Oregon and A. Fjellberg another one from Comox Glacier – Vancouver Island.

The specimens were mounted in Hoyer's medium and became flattened, so that not all the characters are easily visible. The morphological terminology and the system of setal notation for the chelicerae and palpi follows Evans and Till (1979). Terminology for the leg chaetotaxy is based on Evans (1963). The setal notation for the idiosomal dorsum follows Lindquist and Evans (1965) and that for the opisthogaster follows Lindquist (1994). The notation for adenotaxy is based on the system presented by Johnston and Moraza (1991) which is slightly different from that established by Athias-Henriot (1969a). For comparison with other papers the nomenclature of Athias-Henriot (1969a) is given in parentheses: *gdj2* (*gd1*); *gdj4* (*gd2*); *gds3* (*gd3*); *gds4* (*gd4*); *gdz5* (*gd5*); *gdz6* (*gd6*); *gdZ2* (*gd7*); *gdZ3* (*gd8*); *gdZ4* (*gd9*). Measurements of female structures: epigynium shield height (*h*) is the length from the tip of the shield to its posterior margin, epigynium basal width (*b*) represents the length of the posterior margin; setal distance *st-st* is a transverse measurement between insertions of the two setae of the pair *st* on sternal shield. Leg length is taken from the base of the coxa to the apex of the tarsus; tarsus I and IV length were taken from the base of the segment to its apex, without the pretarsus. Measurements in micrometers are specified after each structure. The following abbreviations are used: *hyp.* hypostomatal setae, *pcx.* palpcoxal setae. The definition of cingulum follows Athias-Henriot 1977, as a sclerotized strip which fuses the peritrematal shield with the anterolateral corner of the opisthogastric shield at the level of coxa IV.

The material is deposited in the Canadian National Collection (CNC) of Insects and Arachnids, Ottawa and the Museum d'Histoire Naturelle (MHNG), Geneva, Switzerland (one male, two females of *O. lindquisti* and one female paratype of *O. makarovae*).

Taxonomy

The new genus has a peculiar combination of features which make it unique in the subfamily Pergamasinae (Juvara-Balş 2002; Hruzová and Fend'a (2018). The main characteristics to define a new genus in this subfamily are:

Both sexes: The position of peritrematal shield in connection with the dorsal and ventral one; the presence or absence of the cingulum (Athias-Henriot 1977); the chaetotaxy of dorsal and ventral idiosoma, and the number of gland-pores, especially the presence or absence of gland-pore *gv1*; the shape of the gnathotectum and the form of the palpfemoral and palpgenual setae *al*. Female: the number of teeth on movable digit of chelicera and the characteristics of the epi- and endogynium.

The new genus shares with *Paragamasus* Hull 1918 sensu lato (Athias-Henriot 1971, Karg 1993, Hruzová and Fend'a 2018) the following characteristics: the number of dorsal setae and the leg chaetotaxy (Evans 1963), the peritrematal shield, which is free posteriorly and attached anteriorly to dorsal shield, the trifid gnathotectum and the spatulate setae *al1*, *al2* inserted on palpgenu. The form of dorsal setae *s1*, *s2* (very tiny) and those of male's genital lamina, rounded with a concavity at its anterior margin, resemble those described (Micherdzinski 1969, Juvara-Balş 1977) in *Aclerogamasus* Athias-Henriot, 1971, a former subgenus of *Paragamasus* sensu lato.

The new genus *Occigamasus* differs from the other genera of Pergamasinae in having four teeth on movable digit of female chelicera and the peritrematal shield free posteriorly. This characteristic is found only in *Heteroparasitus* (*Medioparasitus*) *athiasae* (Juvara-Balş 2002) but this genus is different in many other features. Witalinski (1972) noted a variation of three or four teeth on the movable digit in *Heteroparasitus quadratus* Witalinski, 1972. The fact that some species have four teeth seems to be an allopatric characteristic. Other characteristics which characterize *Occigamasus* are the pilose seta inserted on dorsal face of fixed digit as

well as the palpfemoral seta *al* pectinate with four tines (*O. lindquisti* **n. sp.**) or more in the other related species. These features are otherwise specific to some genera of the subfamily Parasitinae.

On podonotum all the setae of the dorsal hexagon (*z5*, *j5*, and *j6*) are similar in length and width. In *Occigamasus* **n. gen.** seta *pd2* is very long on tarsus IV. This characteristic is homoplastic, being described also in some species of *Paragamasus*, *Thenargamasus*, *Leptogamasus* and even in the subfamily Parasitinae (Athias-Henriot 1969a, Micherdzinski 1969, Hruzová and Fend'a 2018).

Descriptions

Occigamasus new genus

Zoobank: [06A18EF2-963A-4E10-AE3C-AF4E8C3E808E](https://zoobank.org/06A18EF2-963A-4E10-AE3C-AF4E8C3E808E)

Figs. 1–9

Type species — *Occigamasus lindquisti* **n. sp.**

Diagnosis — Dorsal shield of adult entire in female, with a median transverse suture between the podonotum and opisthonotum in male. Podonotum with 22 pairs of acicular, simple setae, slightly curved; setae *s1* and *s2* tiny (6–14). Opisthonotal region with 20–21 acicular, simple setae, their tips reaching the following row of setae. Dorsal adenotaxy with five gland pores: *gdj2*, *gdz5*, *gds4*, *gdz6*, *gdZ3*. Ventral adenotaxy with three pairs of gland pores: *gv1* and *gv3* with one solenostome, *gv2* with two solenostomes; *gv1* located closed to medial line of posterior margin of sternal shield or in vicinity of seta *st3*. Epigynium subapical structure spindle-like. Endogynium formed as a big sack which has various structures inside. Ventrianal shield with eight pairs of opisthogastric setae, *ZV1* very short (6–7), *JV5* inserted on soft cuticle. Peritrematal shield free from ventrianal shield posteriorly and attached to podonotal region of holodorsal shield anteriorly.

Gnathotectum simply trifold. Hypognathal groove with 10–11 denticulate rows. Palpgenu with setae *all*, *al2* spatulate; palpfemoral seta *al* pectinate with four or more tines. Movable digit of female chelicera with four teeth, movable digit of male with two teeth; dorsal seta on fixed digit pilose. Trochanter IV, dorsally, with a small protuberance bearing a tiny spine at its apex (Fig. 5E).

Etymology — The name of the genus combines the Latin suffix “occi”-, from, “occidens – occido” meaning ‘west’, with - “gamasus”, referring to gamasine mites. It means the gamasus from the west.

Occigamasus lindquisti **n. sp.**

Zoobank: [C72F41AF-161A-48F9-8413-EB6A549EE64F](https://zoobank.org/C72F41AF-161A-48F9-8413-EB6A549EE64F)

Figs. 1–6

Type material — Canada. British Columbia (B. C.), Vancouver Island: Holotype: 1 female, Cowichan Lake, Experimental Station, mixed Douglas fir, alder, fern litter, 25 July 1975. Paratypes: 2 females, 1 male, same data as holotype; Cowichan Lake Experimental Station, Mesachie Lake: 1 male, 2 females, Douglas fir litter, 24 July 1975; 1 female, from moss, herbs on log; 3 females, moss on rock; 1 male, 2 females, from mixed cedar deciduous shrub litter, 25 July 1975; 1 male, from alder, crab apple litter, 8 July 1975.

Other material examined. Canada. Vancouver Island (B. C.): 2 males, 2 females, Miracle Beach Provincial Park, from Douglas fir shrub litter, 10 July 1975. U.S.A., Oregon Curry County: 3 males, 3 females, Loeb State Park, 8 miles NE, from tan oak, willow, fern litter, 12 July 1985.

Diagnosis — Length of idiosoma: 936–940 female, 768–816 male. Both sexes: gnathotectum with three simple prongs, the central one slightly shorter than the lateral ones. Female: epigynial spindle-like structure 72 long; endogynium a double sack with two curved prongs on its posterior margin and one or two denticles on lateral walls; sternal shield with gland pores *gv1*

Figure 1 *Occigamasus lindquisti* n. sp., female: A–C, F. male: D, E. A, D – Podonotum; B, E – Opisthonotum; C – Submarginal setae UR on lateral cuticle; F – Opisthogaster.

Figure 2 *Occigamasus lindquisti* n. sp., female: A – Sternal shield; B – Tritosternum; C – Paragynia and epigynum; D – Subapical structure of epigynum; Endogynium, E – dorsal view; F – ventral view.

Figure 3 *Occigamasus lindquisti* n. sp., female: A–E. male: F. A – Gnathosoma, ventral view; B – Palpfemur and palpgenu; C, F – Gnathotectum; D – Chelicera; E – Dorsal seta of fixed digit.

located medially on posterior margin; on tarsus IV seta *pd2*, 240 long. Male: armature of leg II with simple rounded protuberance on genu, triangular spur on tibia and a basal protuberance on femoral apophysis; on tarsus IV seta *pd2*, 190–216 long.

Etymology — The species is dedicated to the eminent acarologist Evert E. Lindquist who collected the mites and gave us the opportunity to study the material.

Female (n = 16)

Figs. 1 A–C F, 2–5

Dorsal idiosoma (Figs. 1A, B) — Colour brownish-yellow, holodorsal shield ornamentation finely reticulate on podonotum, scutellate on opisthonotum. Length of idiosoma 936–940. Peritrematal shield united anteriorly with dorsal shield and free posteriorly; length of peritreme ca. 360. All dorsal setae acicular and simple.

Podonotum with 22 pairs of setae, their length: *j1* = 36; *j2* = 84–96; *j3, j4, j5* = 108–114; *j6* = 78; *z1* = 24; *z2* = 36; *z3* = 42; *z4* = 84; *z5* = 60; *z6* = 96–108; *s1* = 12, *s2* = 6–7; *s3* = 96–100; *s4* = 60; *s5* = 120; *s6* = 40; *r2* = 36; *r3* = 168–180; *r4* = 30; *r5* = 36.

Opisthonotum with 20–21 pairs of simple setae, slightly curved, their tips reaching alveoli of following row of setae; their length from 84 to 108. On lateral soft integument five pairs of *UR* setae with lengths 6, 8–9, 18, 36 and 54, respectively. A tiny, fine seta, *x*, (6–12), probably a conversion of a poroid, located on soft cuticle posteriorly to peritrematal shield. Dorsal adenotaxy with five gland-pores: *gdj2*, *gdz5*, *gds4*, *gdz6*, *gdZ3*. Poroidiotaxy with five pores on podonotum and seven on opisthonotum.

Ventral idiosoma — Tritosternum located between presternal plates, its length = 54, basal width = 30; laciniae pilose, 120–132 long (Fig. 2B). Presternal plates formed by two pairs of sclerites, inner one long ellipsoidal and outer one shorter and round; sternal shield reticulated, its mid-length 132–144; gland pores *gv1* located centrally on posterior margin; distance between *gv1*–*gv1* = 36 (Fig. 2A). Distance between pairs of sternal setae: *st1*–*st1* = 120; *st2*–*st2* = 120–144; *st3*–*st3* = 132–144. Length of sternal setae: *st1* = 60; *st2* = 66–72; *st3* = 78. Paragynia weakly reticulated with bean-like, sclerotized formations on their anteromedial margin, *st4* = 66; metagynial sclerite ellipsoidal; epigynum weakly reticulated, its subapical structure spindle-like with a posteriorly rounded membrane (Figs. 2C, D). Epigynum length (h) = 192–216, basal width (b) = 228–240; h/b = 0.8–0.9; length of *st5* = 54, genital pores *iv5* near posterior margin on soft cuticle. Endogynium (Figs. 2E, F) a big sack extending dorsally with a pocket-like structure; on its lateral margin with one or two teeth; posterior margin with two curved prongs (12 long) and distance between them from 18 to 36; this variation depends on different localities from which females were sampled. Ventrianal shield (Fig. 1F) with eight pairs of opisthogastric setae, their lengths: *ZV1* = 7, *ZV2* = 96, *ZV3* = 48, *ZV4* = 72, *JV1* = 48, *JV2*, *JV4* = 66; *JV3* = 77. Setae *JV5* = 72 on soft cuticle. Gland pore *gv2* with two solenostomes, gland pore *gv3* simple near anal opening. Length of simple paranal setae 24 and of anal seta 12.

Gnathosoma — Gnathotectum simple trifid, central prong slightly shorter than lateral ones (Fig. 3C). Hypognathal groove with 10 rows of fine denticles; palpcoxal setae pilose, hypostomatic setae simple, their lengths: *hyp.1*, *pcx* = 72; *hyp.2*, *hyp.3* = 65. Corniculi triangular; palptrochanter with simple setae *v1* (42) and pilose setae *v2* (48) (Fig. 3A). Palpfemur with seta *al* quadripectinate and palpgenu with setae *al1*, *al2* entire, spatulate (Fig. 3B). Chelicera (Figs. 3D, E): movable digit with four teeth; fixed digit with two teeth distal to pilus dentilis and a pilose seta on dorsal face.

Legs (Figs. 4, 5) — Chaetotaxy, specific to Parasitidae as established by Evans (1963) for genus *Pergamasus*. Leg lengths: I = 1056 (tarsus = 250–252); II = 770–787; III = 630; IV = 1128 (tarsus = 360–372).

Chaetome : *coxae* I–IV: 2-2-2-1; *trochanter* I–IV: 6-5-5-5; *femur*: I = 13 (2-5/4-2); II = 11 (2-2/2 3/1-1); III = 6 (1-4/1-0); IV = 6 (1-4/1-0); *genu*: I = 13 (2-3/2 3/1-2); II = 11 (2- 3/1 2/1-2); III = 9 (2 2/1 2/1 1); IV = 10 (2-2/1 3/1-1); *tibiae*: I = 14 (2-3/2 3/2-2); II = 10 (2-2/1 2/1-2); III = 8 (2-1/1 2/1-1); IV = 10 (2-1/1 3/1-2); *tarsi* II–IV: 18.

Figure 4 *Occigamasus lindquisti* n. sp., female: Leg II: A – Tarsus; B – Tibia and genu; C – Femur. Leg III: D – Tarsus; E – Tibia; F – Genu and femur; G – Trochanter.

Figure 5 *Occigamasus lindquisti* n. sp., female: Leg IV: A – Tarsus; B – Tibia; C – Genu; D – Femur; E – Trochanter.

On tarsus (leg II–IV) setae *av1*, *pv1* short (18) and stout; *all*, *pl1* setae fine (48 long), other setae long and simple or slightly pilose.

Setae on leg I mostly simple, thin, their lengths around 66–72, except *av* (genu) short and pilose, *pd3* (tibia) simple, 36 long. Tarsal dorso-distal sensory field, located in the distal third of this segment, consists of a capsule with an arched phanere 48 (sensilla *s* Lindquist and Moraza 2016) and three solenidia (24–30), followed by 10 long sensilla inserted subapically; a single solenidion located ventrally. Leg II (Figs. 4A, C) most of setae simple, their lengths from 72 to 84; on tarsus *md* simple (96), *mv* stout and short; *av1*, *pv1* short (12), *all*, *pl1* (48); setae *al2*, *al3*, *av2*, *pl3*, *pd3* slightly pilose. Tarsus III (Fig. 4D) with long (102) setae *al3*, *pl3*, *mv* stout and short (36), other setae simple or slightly pilose (*pl2*, *al2*, *ad3*). Tarsus IV (Fig. 5A) elongated, with long simple setae; some slightly pilose (*al3*, *pl3*, *ad3*, *pd3*). Seta *pd2* very long (240) almost as long as telotarsus. Lengths of some other setae: *pl3* = 84; *al3*, *ad3* = 120; *mv* = 96; *al2* = 115; *pl2*, *av2* = 130; *pv2* = 72; *all* = 48; *ad1*, *pd1* = 18. A strip of desclerotization present subapically at level of setal insertion *av2* on telotarsus (Fig. 5A). Tibia, genu, femur (Figs. 5B–D) with long fine setae. Trochanter IV with a dorsal protuberance bearing a tiny spine apically (Fig. 5E).

Male (n = 9)

Figs. 1 D–E, 3 F, 6

Idiosoma 768–816 long (Figs. 1D–E). Holodorsal shield provided with a transverse suture, which delineates slightly reticulate podonotum from scutellate opisthonotum. Peritrematal shield 330–324 long. Chaetotaxy, adeno- and poroidotaxy as in female. Lengths of some simple podonotal setae: *j1* = 54, *j2* = 60, *j3* = 96–108, *j5* = 100, *j6* = 78; *z1* = 24–30, *z2* = 24, *z4* = 84, *z5* = 60, *z6* = 84; *s1-s2* = 6, *s4* = 90, *s5* = 84–90, *s6* = 72; *r2* = 36–48, *r3* = 144–168, *r4* = 18, *r5* = 78. Opisthonotum with simple, acicular, curved setae, their tips reaching the following row of setae; their lengths 60–72.

Ventral idiosoma — Slightly sclerotized, cuticular ornamentation reticulate on sternogenital shield and scutellate on opisthogaster. Tritosternum as in female, base length 60, width 36; laciniae 78–84 pilose. Presternal platelets ovoid, each with a tiny anterior fragment. Genital lamina rounded, anterior margin slightly concave; subgenital microsclerite rectangular with rounded anterolateral corners (Fig. 6B). Setae in sterno-genital part simple, their length: *st1*, *st3* = 72; *st2* = 66; *st4* = 66; *st5* = 54. Gland pore *gv1* simple, distance between *gv1*–*gv1* = 50; gland pore *gv2* double with two solenostomes.

Opisthogaster with eight pairs of ventral setae: *ZV1* = 6, others from 36 (*ZV2*) to 84 (*JV5*). Length of simple paranal setae 24 and of postanal seta 18. Gland pore *gv3* near paranal setae.

Gnathosoma — Gnathotectum simple trifid, all prongs of same length (Fig. 3F). Hypognathum and corniculi as in female. Hypognathum with 10 rows of denticles; length of setae: *hyp.1* = 54, *hyp.2* = 48, *hyp.3* = 36–42; slightly pilose *pcx.* = 60. Palptrochanter with setae *v1* simple and *v2* pilose. Palpfemur and palpgenu as in female. Chelicera (Fig. 6A): fixed digit slightly hooked, with subapical denticle distal to pilus dentilis and a row of tiny denticles behind it; dorsal seta pilose. Movable digit with two teeth; arm of spermatotreme with inner protuberance in proximal third; arthrodial membrane fan shaped as a range of setule.

Legs — Lengths of legs: I = 1024 (tarsus = 247–252); II = 756; III = 688; IV = 1044–1128 (tarsus = 336–360). Chaetotaxy of legs as in female. Armature of leg II illustrated in Figs. 6C, D: pyramidal femoral apophysis with small protuberance at its base and short thumb-like axillary process; rounded protuberance on genual distal margin; triangular apophysis located medially on tibia. Leg IV: seta *pd2*, on tarsus, from 190 (site Miracle Beach) to 216 (site Cowichan); trochanter, dorso-laterally, with a protuberance with a spine apically.

See remarks under *O. makarovae* n. sp.

Figure 6 *Occigamasus lindquisti* n. sp., male: A – Chelicera; B – Sternogenital shield. Leg II: C – Tibia; D – Genu and femur.

***Occigamasus makarovae* n. sp.**

Zoobank: 638F2793-AC16-4702-8BBE-A758AC1B6020

Figs. 7 C–D, 8

Type material — U.S.A., Oregon, Deschutes County: 1 female holotype; 1 female, 2 males, 2 deutonymphs, paratypes. Proxy Falls, St. NWY, 28 miles SW Sisters, ca. 1000m alt., from wet moss and deep damp mixed maple-hemlock litter, 19 July 1985.

Diagnosis — Female endogynium a double sack with a median finger-like protrusion and a membrane with two anterolateral denticles; spindle-like epigynial structure long (66–72). Male femoral apophysis thumb-like.

Etymology — The species is dedicated to Olga Makarova for her major contributions to systematics of Gamasina families, including Parasitidae.

Description — Only the characteristics specific to *O. makarovae* are described. Chaetotaxy, adenotaxy and poroidotaxy as in the new genus.

Female (n = 2)

Figs. 7 C–D, 8 B–C, E

Dorsal idiosoma — Length of idiosoma 864; length of peritrematal shield 336. Lengths of some simple acicular, setae: on podonotum $j1 = 48$, $j2 = 84$, $j3-j5 = 108$; $z1 = 18-24$, $z2 = 42$; $s1 = 18$, $s2 = 12$; $r2 = 24$, $r3 = 168$; others from 96 to 108; on opisthonotum length of slightly curved setae from 96 to 108.

Ventral idiosoma — Characteristics of sternal shield, paragynia, metagynial sclerite, epigynium as well as their dimensions as in *O. lindquisti*. Gland pore *gv1* located medially, distance *gv1-gv1* = 24. Endogynium (Fig. 7C) a big sack with a median finger-like protrusion on its posterior margin and a membrane with two anterolateral denticles; epigynium with a long (66–72) subapical spindle-like structure surrounded by a triangular membrane with its posterior margin sclerotized (Figs. 7C, D). Ventrianal shield with 8 pairs of setae, their length: *ZV1* = 8; *ZV2-ZV4* = 60; *JV1*, *JV2*, *JV4* = 72; *JV3* = 66. Setae *JV5* = 78 on soft cuticle.

Gnathosoma — Gnathotectum (Fig. 8C) with three smooth prongs their tips at same level. Hypostomatic and palpcoxal setae simple, their length *hyp.1*, *hyp.2*, *pcx* = 72; *hyp.3* = 48. Seta *al* on palpfemur pentapectinate (Fig. 8E). Chelicera (Fig. 8B): movable digit with four teeth; fixed digit with two small teeth between apex and pilus dentilis and behind it a denticulate ridge ended with a tooth; dorsal seta pilose.

Legs — Chaetotaxy typical of *Occigamasus* n. gen. Setae on legs slender and simple; *pd2* on tarsus IV = 228–230. Trochanter IV with dorsal protuberance with a tiny spine apically (Fig. 8G). Measurements: tarsus I = 240; tarsus IV = 372–384.

Male (n = 2)

Figs. 8 A, D, F–G

Lengths of some dorsal setae: $j1 = 60$, $j2, j3 = 100$; $z1 = 18$; $s1, s2 = 14$; $r2 = 60$, $r3 = 168$, $r4 = 24$; on opisthonotum simple, slightly curved setae, their lengths from 72 to 96.

Ventral idiosoma — Genital lamina rounded with subgenital microsclerite rectangular with anterolateral corners rounded. Other characteristics as in *O. lindquisti*.

Gnathosoma — Gnathotectum smoothly trifid, all prongs of same length (Fig. 8D). Chelicera (Fig. 8A): fixed digit straight with slightly truncate apex, inner margin with a ridge of small denticles behind pilus dentilis; movable digit as in *O. lindquisti*.

Legs — Chaetotaxy as in female. Armature of leg II illustrated in Fig. 8F: femoral apophysis thumb-like, axillary process ellipsoidal and straight. Spurs on genu and tibia as in *O. lindquisti*. Measurements: tarsus I = 240; tarsus IV = 336. Length of seta *pd2* (tarsus IV) = 204.

Figure 7 *Occigamasus* spp., female endogynia: A – *Occigamasus* sp. 1; C – *Occigamasus makarovae* n. sp.; E – *Occigamasus* sp. 2; F – *Occigamasus* sp. 3; G – *Occigamasus* sp. 4; I, J – *Occigamasus* sp. 5. Subapical structure of epigynum: B – *Occigamasus* sp. 1; D – *O. makarovae* n. sp.; H – *Occigamasus* sp. 4; K – *Occigamasus* sp. 5.

Figure 8 *Occigamasus makarovae* n. sp., female: B, C, E. male: A, D, F, G. A, B – Chelicera; C, D – Gnathotectum ; E – Palpfemur; F – Leg II femur, genu, tibia; G – Trochanter IV.

Deutonymph (n = 2)

Fig. 9

Idiosoma. Weakly sclerotized, yellowish colored. Podonotal shield reticulated only laterally while opisthonotal, sternal, and anal shields well reticulated. All setae simple. Peritreme as in adult.

Dorsal idiosoma (Fig. 9A) — Podonotal shield: mid-length 360, width 380 (measured at its widest point). Podosoma with 20 pairs on podonotal shield and two pairs (*s2* and *r4*) inserted, off the shield, on soft cuticle. Short setae: *z1* = 14–15; *s1* = 6–10, *s2* = 12; *r2* = 30, *r4* = 18. Other setae from 60 to 96. Setae *r3* very long 145–156. Adenotaxie: gland pores *gdj2*, *gdj5*, *gds4* weakly developed. Gland pore *gdz6* shifted on anterior margin of opisthonotal shield. Opisthonotal shield with 14 pairs of setae, their length from 72 to 96, seven pairs of pores and gland pores *gdZ3*. Only setae *R1* inserted on shield, other setae of *R* series and submarginal setae (*UR*) located on soft cuticle.

Ventral idiosoma (Fig. 9B) — Tritosternum flanked by a pair of small platelets. Sternal shield with four pairs of setae (*st1*–*st4*) and with three pairs of of lyrifissures *iv1*–*iv3*. Setae *st5* simple, lyrifissures *iv5* near coxae IV. Gland pore *gv2* not developed. Opisthogastric region with eight pairs of setae. Setae *ZV1* very small (2–3). Only pores *ivo4* and *ivp* visible.

Gnathosoma — All the characteristics as in female.

Legs — Chaetotaxy as in adult. Leg I has, apically, a strip of desclerotization located ventrally behind the sensory field (Fig. 9C). Measurements: tarsus I = 240; tarsus IV = 300; setae *pd2* on tarsus IV = 204–216.

Remarks — I could not observe all the characteristics of the deutonymph. The two

Figure 9 *Occigamasus makarovae* n. sp., deutonymph: Idiosoma: A – dorsal view; B – ventral view. C – Tarsus I apically, sensory field, dorsal view.

specimens were squashed so that the posterior part of the idiosoma was in bad condition and the chelicerae were hidden into the gnathosoma. However, I remarked two peculiar characters. On the podosoma the setae *s2* and *r4* were inserted outside the shield like in Parasitinae (Hyatt, 1980). The seta *R1* is located on the opisthonotal shield as in Pergamasinae (Bhattacharyya, 1963). The leg I has apically a ventral fissure which was not mentioned in Pergamasinae. Probably this strip of desclerotization helps the movement of tarsus I which is a sensory segment.

The most important feature distinguishing the female of *O. makarovae* from *O. lindquisti* is the shape of the endogynium (Figs. 2E, 7C). The posterior margin of the endogynium has in *O. lindquisti* two curved prongs about 12 long while in *O. makarovae* only a finger-like protrusion. The fixed digit of the chelicera of these two species is different. Both species have two denticles between pilus dentilis and apex. The inner margin of *O. lindquisti*, which follows the denticles, is smooth whereas in *O. makarovae*, behind the pilus dentilis, there is a denticulate ridge ended by a tooth. The male of *O. makarovae* is similar to those of *O. lindquisti* but it is recognizable by the characteristics of femoral apophysis on leg II and by those of the fixed chelicera. In *O. makarovae* the femoral apophysis is thumb-like whereas *O. lindquisti* has a protuberance on its base. The fixed digit of chelicera in *O. makarovae* has a truncate apex and those of *O. lindquisti* a slightly hooked one.

Undescribed species of *Occigamasus* new genus and their sites

In the Canadian National Collection of Acari (CNC) I identified material of other undescribed species belonging to the new genus. Unfortunately, there was not enough material to describe the new taxa properly. The specimens are mounted dorsally or ventrally so that not all characteristics could be observed. The males are absent in some of the samples. To facilitate future systematic studies, I will mention the sites from where they were sampled, the characteristics of the female endogynium which separate these species (Fig. 7) and some morphological features. Unless otherwise indicated, the samples were collected by E. E. Lindquist.

Occigamasus sp. 1

Material examined — U.S.A. Oregon, Benton County: 1 male, 1 female Mc Glynn Corvallis, ravine, moss on bank, 23 Jan. 1977, coll. L. Russel.

Morphological characteristics — Female (Figs. 7A, B): endogynium a sack with a denticulate plate; epigynum dorsally, with long, sclerotized subapical structure; *gv1* posterad *st3*. Male: leg II with a short apophysis on genu and an elongate spur on tibia; trochanter IV without protuberance. Measurements. Female: tarsus I = 204; tarsus IV = 300, seta *pd2* = 120.

Occigamasus sp. 2

Material examined — Canada. British Columbia, Vancouver Island: 1 female, NE intersection Loc Hwy 4 – 45° lat., cedar litter nearby bog, 29 July 1975; idem, 2 females, Heather Mt. trail, 10 miles NW Caycuse, 1000 m alt., moss in creek, 4 Aug. 1975.

Morphological characteristics — Female (Fig. 7E): endogynium with two lateral processes and a denticulate medial plate; epigynum with the sclerotized spindle-like structure (60 long); *gv1* located medially on posterior margin of sternal shield. Trochanter IV, dorso-laterally, with a spine apically on a protuberance. Measurements: tarsus I = 264; tarsus IV = 384–420; seta *pd2* = 252.

Occigamasus sp. 3

Material examined — Canada, Alberta: 1 female, 1 male, Waterton Lakes National Park, edge of Rowe Lake, from litter, alpine exposure, 2500 m alt., 2–5 Sept. 1980; 1 female, rocky slopes, damp moss; 1 female, 1 male, from grass and substrate, subalpine, by Lake Rowe, 1950 m alt.

Morphological characteristics — Female (Fig. 7F): endogynium a sack with medial spine, lateral protuberances and four anterolateral denticles on the upper margin; epigynum with spindle-like structure; *gv1* located medially on posterior margin of sternal shield. Trochanter IV, dorso-laterally, with triangular protuberance. Male: armature on leg II with a small rounded protuberance on genu and trapezoidal spur on tibia. Measurements. Female: tarsus I = 240, tarsus IV = 360; seta *pd2* = 216. Male: tarsus I = 228, tarsus IV = 288; seta *pd2* = 192.

***Occigamasus* sp. 4**

Material examined — Canada. British Columbia, Vancouver Island: 2 females, Heather Mt. trail, 10 miles NW Caycuse, from hemlock-huckleberry litter, 1000 m, 4 Aug. 1975; 1 female, 1 deutonymph, Comox Glacier, mature hemlock, fir, spruce forest, 176 m alt., moist litter (AF 384), 27 Aug. 1983, coll. A. Fjellberg.

Morphological characteristics — Female (Fig. 7G): endogynium a big sack ventrally with a bifid structure having denticles between the branches, on the anterior margin with a denticulate club surrounded by big denticles; epigynum, dorsally, with a sclerotized ellipsoidal structure with a central spine; *gv1* located lateral to seta *st3*; distance *gv1*–*gv1* = 144. Trochanter IV, dorso-laterally, a small protuberance with a tiny spine apically. Measurements. Female: tarsus I = 252–288, tarsus IV = 336–384; seta *pd2* = 156 (Heather Mt Trail) to 240 (Comox Glacier).

***Occigamasus* sp. 5**

Material examined — Canada. British Columbia, Vancouver Island: 4 females, SE end Long Beach, from salal-kinni-kinnick (*Arctostaphylos uva-ursi* L.) litter on sand dunes, 29 July 1975.

Morphological characteristics — Big mite (Idiosoma length = 1104–1152), well sclerotized, colour brown-reddish. Female (Figs. 7I, J): on sternal shield *gv1* in vicinity of *st3*; endogynium a sack with a club structure anteriorly and many large spines on lateral walls; on posterior margin with a sclerotized, ellipsoidal structure. Epigynum with a bifid, slightly sclerotized subapical structure (Fig. 7K). Trochanter IV with small triangular protuberance on its dorso-lateral face. Measurements. Female: tarsus I = 288, tarsus IV = 396–420; seta *pd2* = 288.

Discussion

The geographical distribution of *Occigamasus* is, until now, limited to the west coast of North America i.e., Vancouver Island, Oregon and California which are included in the Oregonian biotic province and to the Waterton Lakes National Park, Alberta which belong to the Montanian biotic province (Dice 1943 p.128 in Scudder 1979). A preliminary examination of additional material containing the undescribed species 1–5 suggests that in the future we will be able to divide *Occigamasus* into several species groups. Therefore, for a better understanding of this genus and its species diversity, to find and examine additional specimens from more localities in northwestern North America.

The form of setae on the legs, especially those of tarsus IV and tarsus I, the latter dorso-distal sensory field, has been thoroughly examined only in some genera of Parasitidae (Athias-Henriot 1969a, 1980; Micherdzinsky 1969; Hennessey and Farrier 1989; Al-Atawi *et al.* 2002). On tarsus I the number and form of solenidia differ among the genera of Pergamasinae (Micherdzinsky 1969, Juvara-Balş unpublished data). For example, *Occigamasus* has fewer solenidia when compared with *Aclerogamasus* (Juvara-Balş 1977). The leg chaetotaxy in the whole family needs a revision in order to evaluate the importance of this characteristic in taxonomy.

Another problem brought up during this study, is placement of the species *Cycetogamasus californicus* (Banks). This species was described as *Gamasus californicus* by Banks (1904),

then transferred by Essig (1958) to the genus *Parasitus* and subsequently included by Hennessey and Farrier (1988) in the genus *Cycetogamasus* Athias 1980, subfamily Parasitinae. The genus *Cycetogamasus* established by Athias-Henriot (1980) belongs to the "neogamasidienne series" (série néogamasidienne Athias-Henriot 1977, 1980) which is a different lineage from *Pergamasus*, *Parasitus* or even *Neogamasus*. The main characteristics of females of this genus are the separation of the podonotum and opisthonotum, presence of the cingulum, setae *ZV1* very small, and the movable digit with three teeth. Athias-Henriot (1980) added four other species to the genus *Cycetogamasus* and specified "je range dans ce nouveaux taxon une série, probablement hétéroclite d'espèces "néogamasidienne" (1980 pg. 289). The female of *C. californicus* differs from those species by the following characteristics: absence of the cingulum, four teeth on the movable digit of the chelicera and dorsal shield not separated, only with a partial transverse split behind setae *z6* (Figs. 17, 18, lectotype in Hennessey and Farrier 1988). The common features of female *C. californicus* with *Occigamasus* are: scutum peritrematal free posteriorly; on podonotum, 20–22 pairs of acicular setae, *s1* and *s2* tiny; 22 pairs of acicular setae on opisthonotum; opisthogaster with *ZV1* tiny and *gv2* double, eight pairs of opisthogastric setae; movable digit of female with four teeth; palpgenual *al* setae entire and palpfemoral *al* seta pectinate.

I conclude that *C. californicus* must be included in the new genus *Occigamasus* which belongs to the subfamily Pergamasinae Juvara-Balş.

Occigamasus californicus (Banks, 1904) **n. comb.**

Gamasus californicus Banks, 1904: p. 368

Parasitus californicus Essig, 1958: p. 35

Cycetogamasus californicus (Banks, 1904) Hennessey and Farrier, 1988 (lectotype designated: p. 8).

Another species which should be reviewed is *Neogamasus sakhalinus* Volonikhina, 1993. This species cannot be included in the genus *Neogamasus*, as defined by Athias-Henriot 1977, for the following reasons: lack of a cingulum in the female and two teeth on male's movable digit, versus one in *Neogamasus*. The description and figures are not very detailed and need to be augmented with the characteristics of legs chaetotaxy as well as those of the dorsal and ventral adenotaxy. *N. sakhalinus* belongs to the "neogamasidien series" sensu Athias-Henriot 1980, with genus placement yet to be determined.

The mites included in this heterogeneous group need more research in order to redefine the genus or subgenus categories.

I have had the opportunity to study other material, undescribed species, from the CNC from western and eastern Canada and also from Virginia State, U.S.A. The Parasitidae from North and South Carolina have been identified by Hennessey and Farrier (1989). There are great differences between the specimens from the east and those from the west. The species collected from the eastern Canada and northeastern U.S.A. are more similar with those of Europe. Hennessey and Farrier found in Carolina soils *Psilogamasus hurlbutti* Athias-Henriot (1969b) (subfamily Parasitinae) which was collected from Tanzania and *Phytogamasus primitivus* (Oudemans, 1904) (subfamily Pergamasinae) which is a Gondwanian species dispersed in the southern hemisphere (Juvara-Balş and Athias-Henriot 1972). At present the paucity of information on the systematics of Parasitidae mites does not permit a discussion of the relationships between *Occigamasus* of northwestern North America and other genera found on the east coast of Asia (Athias-Henriot 1977, 1980; Tseng 1995). Probably their closest relatives may be found on the east coast of Asia.

Acknowledgements

I would like to thank Evert. E. Lindquist for access to specimens of the Canadian National Collection and for the useful remarks and corrections for the manuscript. For reviewing the manuscript and helpful comments I kindly thank Kamila Hružová. I'm indebted to an

anonymous reviewer for valuable suggestions. I thank Christina Lehmann-Graber, Gilles Roth (MHNG) for scanning the illustrations and Jean Mariaux and the Department of Invertebrates (MHNG) for providing working space and laboratory facilities.

References

- Al-Atawi F., Klompen H., Moser J. C. 2002. Redescription of *Schizosthetus lyriformis* (McGraw and Farrier, 1969) (Parasitiformes: Parasitidae), with revision of the genus. *Internat. J. Acarol.* 28 (4): 341-360. doi:10.1080/01647950208684310
- Athias-Henriot C. 1967a. Observations sur les *Pergamasus*. I. sous-genre *Paragamasus* Hull, 1918 (Acariens anactinotriches: Parasitidae). *Mém. Mus. Hist. nat., Sér. A (Zool.)* 49 (1): 1-198.
- Athias-Henriot C. 1967b. Observations sur les *Pergamasus*. II. Sous-genre *Pergamasus* Berlese, 1904 (Acariens anactinotriches: Parasitidae). *Acarologia* 9 (4): 669-761.
- Athias-Henriot C. 1967c. Observations sur les *Pergamasus* III. Sous-genre *Amblygamasus* Berlese, 1903 (Acariens Anactinotriches : Parasitidae). *Acarologia* 9 :762-800.
- Athias-Henriot C. 1969a. Les organes cuticulaires sensoriels et glandulaires. Poroidotaxie et adénotaxie. *B. Soc. Zool. Fr.*, 94 : 485-492.
- Athias-Henriot C. 1969b. *Psilogamasus hurbutti* n. g. n. sp. gamaside nouveau de Tanzanie (Acariens Anactinotriches, Parasitidae). *Ann. Soc. Ent. Fr. (N.S.)*, 5 (2): 439-449.
- Athias-Henriot C. 1971. *Paragamasus (Tanygamasus) probsti* (Oudemans) (système, géographie) avec quelques mises au point synonymiques (Arachnides, Gamasides tocospermique, Parasitidae). *Zool. Meded. (Leyden)* 45 (16): 167-179.
- Athias-Henriot C. 1977. Untersuchungen über die Gattung *Neogamasus* Tikhomirov mit zwölf neuen Arten aus Korea (cari, Parasitiformes, Gamasida). *Anls. Hist.-Nat. Mus. Nat. Hung.*, 69: 311-341.
- Athias-Henriot C. 1980. Parasitidae nord-coréens (Parasitiformes) : apparentés au genre *Neogamasus* Tikh., avec la création de trois nouveaux genres. *Anls. Hist.-Nat. Mus. Nat. Hung.* 72: 285-294.
- Banks N. 1904. Some Arachnida from California. *Proceedings of the California Academy of Sciences. Zoology* 3 (13): 331-374.
- Banks N. 1910. New American mites. *Proceedings of the Entomological Society of Washington* 12: 2-12, pls. 1-3 doi:10.1093/besa/2.2.12
- Beaulieu F., Knee W., Nowell V., Schwarzfeld M., Lindo Z., Behan-Pelletier V., Lumley L., Young M.R., Smith I., Proctor H.C., Mironov S.V., Galloway T.D., Walter D.E., Lindquist E.E. 2019. Acari of Canada. *ZooKeys* 819: 77-168. doi:10.3897/zookeys.819.28307
- Berlese A. 1906. Monographia del genere *Gamasus* Latr. *Redia* 3: 66-304.
- Bhattacharyya S.K. 1963. A revision of the British mites of the genus *Pergamasus* s. lat. (Acari: Mesostigmata). *Bull. Br. Mus. Nat. Hist. (Zool.)* 11 (2): 131-242. doi:10.5962/bhl.part.4717
- Essig E.O. 1958. *Insects and Mites of Western North America*. The Macmillan Company, New York. pp. 1050
- Evans G.O. 1963. Observations on the chaetotaxy of the legs in the free-living Gamasina (Acari. Mesostigmata). *Bull. Br. Mus. Nat. History (Zool.)* 10: 275-303. doi:10.5962/bhl.part.20528
- Evans G.O., Till W.M. 1979. Mesostigmata mites of Britain and Ireland (Chelicerata: Acari, Parasitiformes). An introduction to their external morphology and classification. *Trans. zool. Soc. Lond.*, 35: 139-270. doi:10.1111/j.1096-3642.1979.tb00059.x
- Ewing H.E. 1909. New North American Acarina. *Trans. Acad. Scien. ST. Lois (Missouri)* 18: 53-77, pls. 8-11.
- Hennessey M.K., Farrier M.H. 1988. Systematic revision of thirty species of free-living, soil inhabiting Gamasine mites (Acari: Mesostigmata) of North America. *N.C. ARS, Tech. Bull.* 285: pp. 123.
- Hennessey M.K., Farrier M.H. 1989. Mites of the family Parasitidae (Acari: Mesostigmata) inhabiting forest soils of North and South Carolina. *N. C. ARS, Tech. Bull.* 292: pp.78.
- Hrúzová K., Fend'a P. 2018. The family Parasitidae (Acari: Mesostigmata)-history, current problems and challenges. *Acarologia* 58 (Suppl): 25-42.
- Hyatt K.H. 1980. Mites of the subfamily Parasitinae (Mesostigmata: Parasitidae) in the British Isles. *Bull. Br. Mus. Nat. Hist. (Zool.)*, 38 (5): 237-378.
- Johnston D. E., Moraza M.L. 1991. The idiosomal adenotaxy and poroidotaxy of Zerconidae (Mesostigmata: Zerconina). In: Dusbabek, F. & Bukva, V. (eds). *Modern Acarology*. SPB academic, The Hague, 2: 349-356.
- Juvara-Balş I. 1972. *Mixogamasus*, un noueae genre de Parasitidae (Acariens : Anactinotriches) de Roumanie. *Acarologia* 14: 3-14.
- Juvara-Balş I. 1977. Quelques remarques sur les *Aclerogamasus* de Roumanie *A. bicalliger* Athias et *A. motasi* n. sp. (Acarina : Gamasida Parasitidae). *Trav. Inst. Spéol. "Emil Racovitza"*, 16:37-45.
- Juvara-Balş I. 2002. A revision of the genus *Heteroparasitus* new status, with the description of *Heteroparasitus (Medioparasitus) athiasae* subgen. n., sp. n. from Spain and with a key to the genera of Pergamasinae (Acari, Gamasida, Parasitidae). *Rev. suisse Zool.*, 109 (19): 23-46. doi:10.5962/bhl.part.79579
- Juvara-Balş I., Athias-Henriot C. 1972. Définition de *Phityogamasus* n. g. et redescription de son espèce type, *Parasitus primitivus* Oudemans (Gamasides, Parasitidae). *Acarologia* 13: 446-459.
- Karg W. 1993. Acari (Acarina), Milben: Parasitiformes (Anactinotricha). *Cohors Gamasina Leach. Raubmilben. Tierwelt Dtl.*, Gustav Fischer Verlag, Jena, 59: 1-475

- Lindquist E.E. 1994. Some observations on the chaetotaxy of the caudal body region of gamasinae mites (Acari: Mesostigmata), with a modified notation for some ventro-lateral body setae. *Acarologia* 35: 323-326.
- Lindquist E.E., Evans G.O. 1965. Taxonomic concepts in Ascidae, with a modified setal nomenclature for the idiosoma of Gamasina (Acarina, Mesostigmata). *Mem. Entomol. Soc. Canada* 47: 1-64. [doi:10.4039/entm9747fv](https://doi.org/10.4039/entm9747fv)
- Lindquist E.E., Aiscough B.D., Clulow F.V., Funk R.C., Marshall V.G., Nesbitt H.H.J., OConnor B.M., Smith I.M., Wilkinson P.R. 1979. Acari. In: Danks H. V. (Ed.) *Canada and its Insect Fauna*. *Mem. Entomol. Soc. Canada* 108: 252-290.
- Micherdzinski W. 1969. *Die Familie Parsitidae Oudemans 1901 (Acarina, Mesostigmata)*. Kraków: Państwowe Wydawnictwo Naukowe. pp. 660.
- Oudemans A.C. 1901. Notes on Acari. Ser.3. *Tijdschr. nederl. dierk. Vereen., Leyden*: 50-88.
- Oudemans A.C. 1904. *Acarologische Aanteekeningen* X. *Ent. Ber.* 1: 140-141. [doi:10.5962/bhl.part.1124](https://doi.org/10.5962/bhl.part.1124)
- Richards L.A. 1976. Parasitid Mites associated with bumblebees in Alberta, Canada (Acarina: Parasitidae; Hymenoptera: Apidae). I. Taxonomy. *Uni. Kansas Scien. Bull.* 50, 13: 731-773.
- Scudder G.G.E. 1979. Present patterns in the fauna and flora of Canada. In: Danks H. V. (Eds) *Canada and its insect fauna*. *Mem. Entomol. Soc. Canada* 108: 87-179. [doi:10.4039/entm111108087-1](https://doi.org/10.4039/entm111108087-1)
- Tseng Y-H. 1995. A taxonomical study of free-living gamasine mite family Parasitidae Oudemans (Acari, Mesostigmata) from Taiwan. *J. Taiwan Mus.*, 48: 11-81.
- Volonikhina I.I. 1993. New species of Gamasid mites of the genus *Neogamasus* (Parasitiformes, Gamasina) from the South of the Far East. *Zool. Zh.*, 72 (8): 11-21.
- Witalinski W. 1972. Mites of the genus *Holoparasitus* Oudemans, 1936 (Acarina, Parasitidae). *Acta Zool. Cracov.*, 17 (9): 217-238.