Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Supporting agricultural research
for sustainable development

Acarologia is under **free license** and distributed under the terms of the
Creative Commons-BY

A new opilioacarid species (Parasitiformes: Opilioacarida) from Crete (Greece) with notes on its karyotype

Maria Magdalena Vázquez^a, Ivalú Macarena Ávila Herrera^b, Pavel Just^{b,c},
Azucena Claudia Reyes Lerma^b, Maria Chatzaki^d, Tim Lukas Heller^e, Jiří
Král^b

^a Division de Ciencias e Ingenierías, Universidad de Quintana Roo, Chetumal, Quintana Roo, Mexico.

^b Laboratory of Arachnid Cytogenetics, Department of Genetics and Microbiology, Faculty of Science, Charles University, Viničná 5, 128 44, Prague 2, Czech Republic.

^c Invertebrate Zoology Unit, Department of Zoology, Faculty of Science, Charles University, Viničná 7, 128 44, Prague 2, Czech Republic.

^d Department of Molecular Biology and Genetics, Democritus University of Thrace, GR - 68100, Alexandroupolis, Greece.

^e Ludwig-Maximilians-University, Munich, Germany.

Original research

ABSTRACT

A new species of opilioacarid mite, *Opilioacarus thaleri* n. sp., is described from mid-level elevation close to the seashore in Crete (Greece). It was found on moister sites, under stones or inside stone accumulations in phrygana vegetation. The peak of activity was found to be in early May. The diagnostic morphological characters include 7–8 eugenital setae (female), 5/7 setae on each sclerite at pregenital and genital areas, 11 setae on genital area (male), anterior dorsal shield with two pairs of eyes and 60–70 stout, ribbed setae, 5 (female) or 6 (male) leaf setae plus one pectinate seta on the palp tarsus, and the “simple” type of ovipositor with two pairs of glands, a pair of ducts plus a basal structure. The female karyotype comprises 16 monocentric chromosomes, predominantly with acrocentric morphology. Comparison with data of the ixodids and mesostigmatids indicates that a low number of chromosomes and predomination of acrocentric chromosomes could be plesiomorphies of the parasitiform mites.

Keywords chromosome; internal structure; Opilioacaridae; *Opilioacarus*; ovipositor

Zoobank <http://zoobank.org/BD5ACF5D-93C2-4BC8-ABDF-4EB51A9044B1>

Introduction

The suborder Opilioacarida is a small, a low-diversity acarine clade (Bernardi *et al.* 2018). Although these mites exhibit features of both acariform and parasitiform mites (Klompen *et al.* 2015), recent molecular analyses clearly support their placement among parasitiform mites (e.g. Klompen 2010). Opilioacarids are probably the most early-diverging lineage among the recent parasitiform mites (Walter and Harvey 2009, Pepato *et al.* 2010). They exhibit many morphological characters, which are considered ancestral within parasitiform mites and which are shared with other arachnid groups, e.g. two pairs of eyes and abdominal segmentation (Coineau and Legendre 1975; Vázquez and Palacios-Vargas 1988; Dunlop and Alberti 2008). Opilioacarids are the only parasitiform mites, which possess complete acarine ontogenetic sequence including a prelarva (Klompen 2000; Bernardi *et al.* 2013b). Due to the position of

Received 05 August 2020

Accepted 31 May 2021

Published 07 June 2021

Corresponding author
Maria Magdalena Vázquez:
marvazqu@uqroo.edu.mx

Academic editor
Tsolakis, Haralabos

DOI
10.24349/acarologia/20214449

ISSN 0044-586X (print)
ISSN 2107-7207 (electronic)

© Copyright
Vázquez M. M. *et al.*

Distributed under
Creative Commons CC-BY 4.0

the opilioacarids as a basal group of parasitiform mites, their characters can be used for the reconstruction of parasitiform phylogeny (Grandjean 1936; van der Hammen 1970, Klompen *et al.* 2007, Pepato *et al.* 2010). Despite the considerable importance of opilioacarids for reconstruction of arachnid phylogeny, there are only few studies aimed at understanding their phylogeny, ethology, and ecology. Opilioacarids are relatively large, free-living scavengers of arthropods and pollen and possibly occasional predators (Walter and Proctor 1998). A recent study of Vázquez *et al.* (2018) suggests a complex behaviour of opilioacarids, which is, however, not sufficiently understood and includes courtship, brood care and care for immature instars. Opilioacarids are usually found under stones or in leaf litter. Recent study from Belize also revealed opilioacarids under bark of fallen trees or under moss and lichens on tree bases (Vázquez *et al.* 2018). Despite their low diversity, the opilioacarids inhabit a wide range of elevations (from altitudes close to sea level up to 2,000m) and biotopes (from coastal plains and mangroves to uplands like altiplano and from semi-desert biotopes to wet tropical forests). They also occur in hypogean environments (Bernardi *et al.* 2012; Vázquez and Klompen 2015; Moraza *et al.* 2021).

The suborder Opilioacarida currently includes a single family Opilioacaridae comprising 13 genera, 52 species, and 2 subspecies (Araújo *et al.* 2020; Moraza *et al.* 2021). The first described opilioacarid was *Opilioacarus segmentatus* (the type species) from Algeria, which was described by Carl Johannes With in 1902 (With 1902). For the following 40 years, relatively little attention was paid to opilioacarids, the findings of which were limited to the Mediterranean region. During the last two decades, the number of studies dealing with opilioacarid taxonomy has considerably increased. Nowadays, opilioacarids are known from tropical and temperate regions of all continents except for Antarctica (Araújo *et al.* 2020). Remarkably, particular species are often endemic to relatively narrow regions. Opilioacarids were found in South America (Lehtinen 1980; Vázquez *et al.* 2014, 2015, 2018; Bernardi *et al.* 2012; 2013a, b; 2014; Araújo *et al.* 2018a), Mesoamerica and the Antilles (Juvara-Bals and Baltac 1977; Vázquez and Palacios-Vargas 1988; Vázquez and Klompen 2002, 2009, 2015), the southern part of the United States (Chamberlin and Mulaik 1942; van der Hammen 1966), sub-Saharan Africa including Madagascar (Naudo 1963; van der Hammen 1977; Vázquez and Klompen 2010), the Arabian peninsula (van der Hammen 1969), India (Das and Bastawade 2007), central (Krivolutsky 1965) and southeastern Asia (Leclerc, 1989), and Australia (Walter and Proctor 1998).

The type genus *Opilioacarus* is restricted to the Mediterranean region of Europe and North Africa (Vázquez and Klompen 2009). This genus includes six species, namely *O. segmentatus* With, 1902 (Algeria), *O. italicus* (With, 1904), *O. brignolii* Araújo and Di Palma, 2018 (Italy), *O. baeticus* Moraza, Prieto and Balanzategui, 2021 (Spain), and two fossil species from amber (Dunlop *et al.* 2010; Dunlop and Bernardi 2014). Previously, the genus *Opilioacarus* also included some New World species, which are currently placed into the genus *Neocarus*. The revision of various samples revealed that diagnostic characters of *Neocarus* are sufficiently robust to constitute a separate genus (Vázquez and Klompen 2002, 2009).

Specimens of *Opilioacarus* were also collected in Greece, namely Peloponnese peninsula and islands of Crete (Thaler and Knoflach 2002), Korfu (Silvestri 1905), Kassos, Karpathos, and Rhodes (Beron 1990). These specimens were classified as *O. segmentatus*. However, taxonomic revision of material from Greece is needed (Thaler and Knoflach 2002, Moraza *et al.* 2021). Our material collected on the island of Crete at the locality of Thaler and Knoflach (2002) allowed us to describe a new species of the genus *Opilioacarus* and to bring some information on its life history and karyotype. In contrast to the parasitiform suborders Ixodida and Mesostigmata, for opilioacarids the chromosome data are unknown.

Figure 1 Map of Greece, black triangle shows the type locality of *Opilioacarus thaleri* n. sp.

Material and methods

Material

Specimens were collected during four expeditions in Greece, western Crete, Georgioupolis, hill to the west of the town (35.372193° N, 24.256336° E) (Figure 1). Three trips took place in early spring (25.III.–2.IV.2015, 22.IV.–1.V.2017, 1.–8.V.2019), another in early autumn (15.IX.–24.IX.2013). Opilioacarids were collected under large stones on moister spots of phrygana vegetation or inside stone accumulations (Figure 2), which were also inhabited by spider species, which normally occur in caves [pholcid *Hoplopholcus labyrinthi* (Kulczyński, 1903), leptonetid *Sulcia cretica* Fage, 1945]. One or two opilioacarid specimens occurred on stones. However, in early May 2019 (beginning of the hot period) the distribution of specimens was different. During this period, opilioacarids were more abundant. Some stones were inhabited by more (up to eight) specimens.

After sampling, the specimens were reared in plastic containers with slightly moistened cellulose. Water as needed was provided in the form of droplets. Opilioacarids were fed by tiny granules of dry yeast, pollen, and crushed bodies of *Drosophila* and chironomids. They preferred the granules of dry yeast. Specimens were fed twice per week. Excess food was removed to prevent fungal growth.

Figure 2 Habitat of *Opilioacarus thaleri* n. sp.

Morphological studies

Some specimens used for study of morphology were slide-mounted. For this purpose, specimens were cleared in lactic acid for two days, washed in distilled water, dissected and mounted on slides in Hoyer's medium (Krantz and Walter 2009) with the aim to observe their structures under light microscopy. The main structures (idiosoma, gnathosoma, and legs) were mounted on separate slides and kept in the oven at 50 °C for a week. After that the slides were sealed.

Identification drawings were prepared using a Zeiss phase contrast microscope, connected to a drawing tube; measurements were taken using an ocular micrometer and are presented in micrometers (µm). Measurements of idiosomal length and width might be distorted, as they are based on slide-mounted specimens. The description of palps follows the nomenclature of Grandjean (1936) and Vázquez and Klompen (2002). Description of subcapitulum, chelicerae, and sternal area follows Klompen *et al.* (2015). The examined material is deposited in the collection of JK (Department of Genetics and Microbiology, Charles University, Prague, Czech Republic).

Chromosome preparations and their evaluation

Mitotic plates were obtained from the content of the idiosoma of five female specimens, using a spreading technique described by Dolejš *et al.* (2011). Briefly, the dissected tissues were hypotonized in 0.075M KCl for 20 min and then fixed in three rounds of freshly prepared ethanol:acetic acid (3:1) fixative (6, 10 and 20 min, RT). A piece of fixed tissue was suspended into a drop of 60% acetic acid using a pair of tungsten needles. Another tungsten needle was used to gradually spread the resulting cell suspension along the microscope slide placed on a heated histological plate (40°C). Preparations were stained for 30 min in 5% Giemsa solution in Sörensen buffer (pH 6.8). They were inspected using the BX 50 microscope (Olympus). Images were taken under immersion objective 100x using a DP 71 CCD camera (Olympus) and optimized in Corel PHOTO-PAINT X4 programme.

Taxonomic section

Suborder Opilioacarida (following Krantz and Walter, 2009)

Family Opilioacaridae With, 1904

Opilioacarus With, 1902

Type species *Opilioacarus segmentatus* With, 1902: by original designation

The genus currently includes four extant species, namely *O. segmentatus* With, 1902 (Mediterranean region of Algeria), *O. italicus* (With, 1904) (Sicily), *O. brignolii* Araújo and Di Palma, 2018 (southern part of the Italian peninsula and Sardinia), and *O. baeticus* Moraza, Prieto and Balanzategui, 2021 (Spain). *O. italicus* is a *nomen dubium* (Araújo et al. 2018b). Types of this species are missing, the original description is insufficient, and the recollecting efforts at the type locality were unsuccessful (Araújo et al. 2018b). The main differences between the species of *Opilioacarus* described by With are in the size of their legs. Information on chaetotaxy, a standard feature nowadays, was provided only for species described recently (Araújo et al. 2018b, Moraza et al. 2021) and partially for *O. segmentatus* (Grandjean 1936). Recently, two fossil species have been placed putatively into the genus *Opilioacarus*, namely the Eocene *O. aenigmus* Dunlop et al., 2010 from Baltic amber (Dunlop et al. 2010) and the Upper Cretaceous *O. groehni* Dunlop and Bernardi, 2014 from Burmese amber (Dunlop and Bernardi 2014). These findings indicate a considerable antiquity of the clade containing *Opilioacarus*.

Diagnosis: 4–6 foliate *d* setae (*d1*) and one slightly pectinate *d* seta (*d2*) on palp. With's organ membranous, discoid, hyaline. Two pairs of dorsolateral eyes, preanal region with 4–5 dorsal setae and 2 lateroventral setae. Dorsal segments between prodorsal shield and preanal region without setae. Leg I exhibits crown-like eupathidium ($\zeta 1$) close to main sensilla group.

Opilioacarus thaleri n. sp. Vázquez, Ávila and Just

Zoobank: 9E2813B3-3B51-424B-A57C-85DD2F0C95C4

Figures 3–10

Material examined — Specimens examined were collected manually by J. Král, 1.–8.V.2019 in Greece, western Crete, Georgioupolis, hill to the west of the town, 35.372193° N, 24.256336° E. To evaluate characters, the slides were prepared from holotype female, 3 paratype females, and 4 paratype males. Specimens dissected for chromosomes were preserved in 96% ethanol. Some of them (6 females, 4 males) were also slide-mounted. These specimens are also designated as paratypes. The type material is deposited in the collection of JK (Department of Genetics and Microbiology, Charles University, Prague, Czech Republic).

Etymology — This species is dedicated to Konrad Thaler and his wife Barbara Thaler-Knoflach, outstanding Austrian arachnologists, who collected and reported opilioacarids from several places in Greece (Thaler and Knoflach 2002).

Diagnosis — Palp genu with ~6 *p*-type setae, palp femur with 13 *p*-type setae. Palp tarsus with foliate setae *d1* with 3 rounded ending acute lobes each and 1 thin pectinate seta *d2*. Males and females differ by number of setae *d1*. While males exhibit 6 setae *d1*, females shows only 5 *d1* setae. Pregenital region in males with 3 short and thick, ribbed and blunt pregenital setae. Pregenital region in females without pregenital setae. It has 6 eugenital setae. Genital region in males with 11 thin and lightly serrated and tapering setae. Ovipositor of the simple type, folded with 2 pairs of glands and two pairs of ducts on the basis. Internal basal structure of the ovipositor smooth, without any spikes, bifurcated on the top.

Description*Measurements*

Measurements are summarized in Table 1.

Figure 3 *Opilioacarus thaleri* n. sp., dorsal view on an adult specimen in ethanol.

Gnathosoma

Chelicera (Figure 4A). Basal segment with 1 seta, fixed digit with 3 setae. All setae are lightly barbed. Dorsal and antiaxial lyrifissures well developed. Fixed digit with one tooth, movable digit with 1–3 teeth and a well developed terminal hook. Movable digit with large denticle and two very small denticles. Without sexual dimorphism.

Subcapitulum (Figure 4B). All four pairs of paralaial setae are present: *pl1* small, conical; With’s organ (*pl2*) membranous, hyaline, discoid and barbulated. Rutella (*pl3*) with 1 row of 5 teeth, inserted dorsolaterally; *pl4* small but distinct, inserted dorsally on subcapitulum. In addition, 4 circumbuccal (*cb*) and 5–7 medium and subcapitular, latero-ventral (ventral medium in part), latero-ventral medium, latero-dorsal medium, ventral posterior, and latero-ventral posterior setae. Females with 5 subcapitular setae, males with 7 subcapitular setae. Lateral lips with distinct channels.

Palp (Figures 5–7, Tables 2, 3). Trochanter with 4 ribbed, tapering setae (= *r*-type) in

Table 1 Comparative measurements of females and males of *Opilioacarus* spp.

	Length	Width	Position	Source
	♂	♂		
<i>O. thaleri</i> n. sp	1810–1915 µm	787–850 µm	at the level of stigmata	Present work
<i>O. brignolii</i>	824–1327 µm	181–224 µm	at the level of ocelli	Araújo <i>et al.</i> (2018)
<i>O. segmentatus</i>	1600–1900 µm	?		Grandjean (1936)
<i>O. baeticus</i>	1944–2000 µm	944–1111 µm		Moraza <i>et al.</i> (2021)
	♀	♀		
<i>O. thaleri</i> n. sp	2150–2400 µm	1240–1310 µm	at the level of stigmata	Present work
<i>O. brignolii</i>	910–1154 µm	183–228 µm	at the level of ocelli	Araújo <i>et al.</i> (2018)
<i>O. segmentatus</i>	2200–2750 µm	?		With (1904)
<i>O. segmentatus</i>	1600–1900 µm	?		Grandjean (1936)
<i>O. baeticus</i>	2161–2445 µm	1000–1138 µm		Moraza <i>et al.</i> (2021)

Figure 4 *Opilioacarus thaleri* n. sp., gnathosoma. A – chelicera, female. Fixed digit with 3 lightly barbed setae (a) and lyrifissure (b). Movable digit with one large tooth and 3 denticles (c). Basal segment with one seta (d); B – subcapitulum. Paralabial setae pl1, With's organ pl2, rutellum pl3, circumbucal setae cb, and channels ch.

females (Figure 5A), males with 5 *r*-type setae (Figure 6A); femur with 7 papilliform setae (= *p*-type) in females (Figure 5B), males with 13 *p*-type setae (Figure 6B); genu with 4 *p*-type setae and 19 *r*-type setae in females (Figure 5C), males with 6 *p*-type setae and 29 *r*-type setae (Figure 6C); tibia with 17 long lightly serrate and pointed setae (= *s*-type) and 20 *r*-type setae and two thin smooth sensilla (*ss*) in females (Figure 5D), males with 26 *s*-type setae and 36 *r*-type setae and 2 *ss* (Figure 6D). Tibia and tarsus partially fused. Tarsus with 6 foliate (*d1*-type) and 1 pectinate (*d2*-type) setae in males (Figure 7A–B). Females with 5 foliate *d1* setae and 1 pectinate *d2* (Figure 7C–D). Females with 6 *v1*, 3 *v2*, 3 *ss*, 14 *ch*, 5 *sm1*, 3 *sm2*. Males with 8 *v1*, 3 *v2*, 3 *ss*, 12–14 *ch*, 6 *sm1*, 3 *sm2* (Table 3). Lyrifissures *ia* and *ix* present (Figure 7A–B). Pretarsus with a pair of well-developed, smooth, sessile claws (Figure 7).

Idiosoma

Color. Light blue and violet body; combination of blue-purple and white stripes on the legs.

Sternogenital region (Figures 8–9, Table 4). Sternal verrucae in adults each with 3–4 large serrate and pointed and 1 composite (*st1*) setae. Remaining sternal region with 4–5 pairs of stout, ribbed, setae with blunt-tips, and 2 pairs of composite (*st1*) setae in both females and males. Pregenital capsules each with 1 long tapering seta (*st5*) and 4–5 stout, ribbed, blunt-tipped setae (Table 4).

Pregenital area in males with 3 short ribbed blunt-tips pregenital setae (Figure 8A).

Table 2 Comparative setal pattern for palp of adults of *Opilioacarus* spp. The data on *O. segmentatus* are not available. The nomenclature of setae follows Grandjean (1936) and Vázquez and Klompen (2002).

	Trochanter	Femur		Genu		Tibia		Source
	<i>r</i> -type	<i>p</i> -type	<i>r</i> -type	<i>p</i> -type	<i>r</i> -type	<i>s</i> -type	<i>r</i> -type	
<i>O. brignolii</i> ♀	3	7–8	11–12	4–7	20–28	3	38–43	Araújo <i>et al.</i> (2018)
<i>O. brignolii</i> ♂	3–4	5–9	12–19	1	43–54	3	43–54	Araújo <i>et al.</i> (2018)
<i>O. thaleri</i> ♀	4	7	10	4	19	17	20	Present work
<i>O. thaleri</i> ♂	5	13	13	6	29	26	36	Present work
<i>O. baeticus</i> ♀	6–8	15	12	9	41	4	48	Moraza <i>et al.</i> (2021)

Figure 5 *Opilioacarus thaleri* n. sp., female palp. A – trochanter; B – femur; C – genu; D – tibia. Axial (left) and antiaxial (right) view. E – particular types of setae.

Pregenital area in females without pregenital setae (Figure 8B, Table 4). It has 6 thin lightly serrated eugenital setae (Figure 9A, Table 4). Genital region in males with 11 thin lightly serrated genital setae (Figure 8A). Ovipositor folded, with two pairs of gland-like structures and a pair of ducts. Internal basal structure smooth, without any spikes, bifurcated on the top

Table 3 Comparative setal pattern for the palp-tibiotarsus of adults of *Opilioacarus* spp. Sex of *O. segmentatus* is not specified by Grandjean (1936).

	v1	v2	ss	d1	d2	ch	sm1	sm2	sm3	Source
<i>O. segmentatus</i>	?	?	3	5	1	16	8?	5?	0	Grandjean (1936)
<i>O. brignolii</i> ♀	4–5	2–3	3	4	1	14–15	10–11	?	0	Araújo <i>et al.</i> (2018)
<i>O. brignolii</i> ♂	4	2–3	3	4	1	11–14	8–12	?	0	Araújo <i>et al.</i> (2018)
<i>O. baeticus</i> ♀	4	4	3	5	1	23	14	?	0	Moraza <i>et al.</i> (2021)
<i>O. baeticus</i> ♂	4	3	3	5	1	16	14	?	0	Moraza <i>et al.</i> (2021)
<i>O. thaleri</i> n. sp. ♀	8	3	3	5	1	12–14	5	3	0	Present work
<i>O. thaleri</i> n. sp. ♂	6	3	3	6	1	14	6	3	0	Present work

Figure 6 *Opilioacarus thaleri* n. sp., male palp. A – trochanter; B – femur; C – genu; D – tibia. Axial (left) and antiaxial (right) view. E – particular types of setae.

(Figure 9B).

Dorsum. Prodorsal shield in adults with 60–70 stout ribbed setae and two pairs of lateral eyes. Dorsal idiosoma between the shield and the pre-anal segment without setae, but with numerous lyrifissures arranged in transverse rows. Pre-anal segment with 4–5 dorsal and two lateroventral short, stout and ribbed setae (Figure 9C). Anal plates with 12–13 papilliform setae. Dorsum without sexual dimorphism.

Taxonomic remarks

The material collected in Crete, Greece allowed us to describe a new species *O. thaleri* and to compare it with *O. segmentatus*, *O. brignolii* and *O. baeticus*. It should be underlined that the diversity of *Opilioacarus* could be higher than recognized. Specimens from other Greek localities [Peloponnese (Thaler and Knoflach 2002), island of Korfu (Silvestri 1905), islands of Kasos, Karpathos and Rhodes (Beron 1990)] were not revised yet. Moreover, data on *Opilioacarus* indicate that this genus occurs throughout all the coast of Mediterranean Sea forming many species, which inhabit a narrow geographic region.

The number of the setae on the penultimate segment is one of the main characteristics of

Figure 7 *Opilioacarus thaleri* n. sp., palp tarsus, male (A–B) and female (C–D). Ventral (left) and dorsal view (right). E – particular types of setae.

Figure 8 *Opilioacarus thaleri* n. sp., sternogenital area. A – male; B – female, C – particular types of setae. Pregenital region pg, genital region gen.

Figure 9 *Opilioacarus thaleri* n. sp., female. A – eugenital setae; B – ovipositor; C – setae on penultimate segment.

Table 4 Comparative setal pattern for the sternogenital region of adults of *Opilioacarus* spp. Sex of *O. segmentatus* is not specified by Grandjean (1936).

	Sternal verrucae		Sternogenital region		Genital verrucae		Pregenital region		Genital area		♀	Eugenital setae	Source
	stout setae	tapering setae	stout setae	tapering setae	stout setae	tapering setae	stout setae	tapering setae	stout setae	tapering setae			
<i>O. segmentatus</i>	?	?	?	?	3	1	?	?					Grandjean (1936)
<i>O. brignolii</i> ♀	3–3	1	4–6	2	5–7	1	0	0	0	0		4–9	Araújo <i>et al.</i> , (2018)
<i>O. brignolii</i> ♂	2–3	1	4–5	2	4–6	1	0–5	0–3		2–7			Araújo <i>et al.</i> , (2018)
<i>O. baeticus</i> ♀	3–5	1	2–3	2	5–7	1	0	0				5	Moraza <i>et al.</i> , (2021)
<i>O. baeticus</i> ♂	3–4	1	3	2	6–7			5		5			Moraza <i>et al.</i> , (2021)
<i>O. thaleri</i> n. sp. ♀	3–4	1	4–5	2	5	1	0	0				6	Present work
<i>O. thaleri</i> n. sp. ♂	3–4	1	4–5	2	5	1	3	0		11			Present work

the genus *Opilioacarus*. There are 4–5 dorsal and two lateroventral stout ribbed setae in both female and male. This characteristic is constant in the genus. The genus *Neocarus* has only one dorsal and two lateroventral stout ribbed setae on the penultimate segment.

There is some variation in the number of setae on the palp among *Opilioacarus* species. *Opilioacarus segmentatus* and *O. baeticus* have 5 foliated *d1* type setae on the palp (Grandjean 1936; Moraza *et al.* 2021) while *O. brignolii* has 4 foliated *d1* type setae (Araújo *et al.* 2018b). Males of *O. thaleri* present 6 foliated *d1* setae while females of this species have 5 foliated *d1* (this study). The lowest number of *ch*-type setae was found in *O. brignolii* (11–14) (Araújo *et al.* 2018b) and *O. thaleri* (12 in females and 14 in males) (this study). *Opilioacarus segmentatus* has 16 *ch* setae (Grandjean 1936) while *O. baeticus* exhibits 23 *ch* setae in females and 16 *ch* setae in males (Moraza *et al.* 2021).

In the genital area, the differences among *Opilioacarus* species are also significant. Pattern of eugenital setae in female pregenital area is often species-specific. *Opilioacarus segmentatus* and *O. baeticus* exhibits 5 eugenital setae (Grandjean 1936; Moraza *et al.* 2021). *Opilioacarus brignolii* presents 4–9 smooth eugenital setae (Araújo *et al.* 2018b) while *O. thaleri* shows 6 of these setae (this study). Pattern of setae in male pregenital and genital area is also species-specific. Males of *O. brignolii* have in trapezoidal area 2–7 acute ribbed genital setae (Araújo *et al.* 2018b). Males of *O. segmentatus* have 5 short tapering and smooth pregenital setae (Grandjean 1936). Males of *O. baeticus* have 5 pregenital and 5 genital setae (Moraza *et al.* 2021) while males of *O. thaleri* have 3 short ribbed blunt pregenital setae and 11 slightly serrated acute genital setae (this study), which is a very significant difference. Finally, there are significant interspecific differences in morphology of the ovipositor. In *O. thaleri* the ovipositor has 2 pairs of glands and 1 pair of ducts on the basis of the ovipositor. These structures were found neither in *O. brignolii* (Araújo *et al.* 2018b) or *O. baeticus* (Moraza *et al.* 2021).

Karyotype

In the present study, we reported for the first time chromosome data of opilioacarids. The female karyotype consists of 16 large monocentric chromosomes, predominantly with acrocentric morphology (Figure 10). The karyotype of *O. thaleri* n. sp. corresponds to the karyotypes of parasitiform suborders Ixodida and Mesostigmatida in terms of diploid number and chromosome morphology. While chromosomes of these groups were studied intensely due to their medical and economical importance, there are no cytogenetic data on Holothyrida, the fourth parasitiform suborder. Like in *O. thaleri*, 1) ixodids and mesostigmatids exhibit a low number of monocentric chromosomes (the diploid set of ixodids ranges from 12 to 32 chromosomes and diploid set of mesostigmatids from 6 to 26 chromosomes), and 2) karyotypes of most ixodids and mesostigmatids are predominated by acrocentric chromosomes (Oliver 1977, 1989; Norton *et al.* 1993). These features could be plesiomorphic for parasitiform mites.

Figure 10 *Opilioacarus thaleri* n. sp. ($2n = 16$), female mitotic metaphase stained by Giemsa. Note predominance of acrocentric chromosomes. One large pair is probably metacentric (arrowheads).

Acknowledgements

We are very grateful to B. Thaler-Knoflach (University of Innsbruck, Austria) for specification of the opilioacarid locality in the neighbourhood of Georgioupolis, Crete. Furthermore, we are thankful to N. Kordatzakis and her family (Georgioupolis, Greece), P. Pousková (Charles University, Prague), H. Čermáková (Czech Ministry of Education, Youth, and Sports), and B. Lejtnarová (embassy of the Czech Republic in Greece), taking part in the organization of JK's expeditions to Crete. These expeditions were supported by several scholarships, which were based on agreement between the Czech Ministry of Education, Youth, and Sports and the Greek Ministry of Education, Lifelong Learning, and Religious Affairs. Research stay of MMV at the Charles University (Prague) focused on opilioacarid taxonomy was supported by the Mobility Fund of this institution. Our study was also funded by the projects of the Czech Ministry of Education, Youth, and Sports (SVV 260314: ACRL, IMAH, JK; SVV 260571/2021: PJ; LTAUSA 19142: IMAH, JK) and Chilean National Agency for Research and Development (ANID) (IMAH).

References

- Araújo M.S., Bichuette M.E., Bauchan G.R., Ochoa R., Feres R.J.F. 2018a. A new species of cave dwelling *Neocarus* (Acari: Opilioacaridae) from Bahia state, Brazil, with remarks on taxonomic characters. *Zootaxa*, 4402(2): 303-322. doi:10.11646/zootaxa.4402.2.4
- Araújo M.S., Di Palma A., Feres R.J.F. 2018b. A new species of *Opilioacarus* With, 1902 (Acari: Opilioacaridae) from Italy, and a new diagnosis of the genus. *Zootaxa*, 4500(1): 135-145. doi:10.11646/zootaxa.4500.1.9

- Araújo, M.S., Di Palma, A., Feres, R.J.F. 2020. Catalog of the Opilioacarida (Acari: Parasitiformes). *Zootaxa*, 4895: 332-356. doi:10.11646/zootaxa.4895.3.2
- Bernardi L.F.O., Zacarias M.S., Ferreira R.L. 2012. A new species of *Neocarus* Chamberlin & Mulaik, 1942 (Acari: Opilioacarida) from Brazilian caves and karst areas. *Zootaxa*, 3416: 53-68. doi:10.11646/zootaxa.3416.1.5
- Bernardi L.F.O., Silva F.A.B., Zacarias M.S., Klompen H., Ferreira R.L. 2013a. Phylogenetic and biogeographic analysis of the genus *Caribeacarus* (Acari: Opilioacarida), with description of a new South American species. *Invertebr. Syst.*, 27: 294-306. doi:10.1071/IS12041
- Bernardi L.F.O., Klompen H., Zacarias M.S., Ferreira R.L. 2013b. A new species of *Neocarus* Chamberlin & Mulaik, 1942 (Opilioacarida: Opilioacaridae) from Brazil, with remarks on postembryonic development. *ZooKeys*, 358: 69-89. doi:10.3897/zookeys.358.6384
- Bernardi L.F.O., Klompen H., Ferreira R.L. 2014. *Neocarus caipora*, a new mite species (Parasitiformes: Opilioacarida: Opilioacaridae) from Brazilian Amazon caves. *Acarologia*, 54(1): 47-56. doi:10.1051/acarologia/20142113
- Bernardi L.F.O., Zacarias M.S., Borges-Filho E.L. 2018. *Neocarus spelaion* sp. n. (Parasitiformes, Opilioacaridae), a new species of cave dwelling *Neocarus* from Minas Gerais state, Brazil. *Subterr. Biol.*, 27: 1-16. doi:10.3897/subterbiol.27.25777
- Beron P.K. 1990. On the occurrence of *Opilioacarus segmentatus* With, 1903 (Arachnida, Opilioacarida) on the islands of Kassos, Karpathos and Rhodes (Greece). *Acta Zool. Bulg.*, 39: 64-66.
- Chamberlin R.V., Mulaik S. 1942. On a new family of Notostigmata. *Proc. Biol. Soc. Wash.*, 55: 125-132.
- Coineau Y., Legendre R. 1975. Sur un mode de régénération appendiculaire inédit chez les Arthropodes: la régénération des pattes marcheuses chez les Opilioacariens (Acari: Notostigmata). *C. R. Hebd. Séanc. Acad. Sci. Paris*, 280: 41-43.
- Das N.P.I., Bastawade D.B. 2007. The first report of the acarine suborder Opilioacarida from India, with description of new genus, *Indiacarus*, and a new species, *Indiacarus pratyushi*. *Acarologia*, 47(1-2): 3-11.
- Dolejš P., Kořínková T., Musilová J., Opatová V., Kubcová L., Buchar J., Král J. 2011. Karyotypes of central European spiders of the genera *Arctosa*, *Tricca* and *Xerolycosa* (Araneae: Lycosidae). *Eur. J. Entomol.*, 108: 1-16. doi:10.14411/eje.2011.001
- Dunlop J.A., Alberti G. 2008. The affinities of mites and ticks: a review. *J. Zool. Syst. Evol. Res.*, 46(1): 1-18.
- Dunlop J.A., Sempf C., Wunderlich J. 2010. A new opilioacarid mite in Baltic amber. In: Nentwig W., Entling M., Kropf C. (Eds). *European Arachnology 2008*. Bern: Natural History Museum. pp. 59-70.
- Dunlop J.A., Bernardi L.F.O. 2014. An opilioacarid mite in Cretaceous Burmese amber. *Naturwissenschaften*, 101(9): 759-763. doi:10.1007/s00114-014-1212-0
- Grandjean F. 1936. Un acarien synthétique: *Opilioacarus segmentatus* With. *Bull. Soc. Hist. Nat. Afr. Nord*, 27: 413-444.
- Juvara-Bals I., Baltac M. 1977. Deux nouvelles espèces d'*Opilioacarus* (Acarina: Opilioacarida) de Cuba. In: Orghidan T., Núñez Jiménez A., Decou V., Negrea S., Viña Bayés N. (Eds). *Résultats des Expéditions Biospéologiques Cubano-Roumaines à Cuba*. Bucuresti: Academiei Republicii Socialiste Romania. pp. 169-184.
- Klompen H. 2000. Prelarva and larva of *Opilioacarus (Neocarus) texanus* (Chamberlin and Mulaik) (Acari: Opilioacarida) with notes on the patterns of setae and lyrissures. *J. Nat. Hist.*, 34: 1977-1992. doi:10.1080/00222930050144819
- Klompen H. 2010. Holothyrids and ticks: new insights from larval morphology and DNA sequencing, with the description of a new species of *Diplothyridus* (Parasitiformes: Neothyridae). *Acarologia*, 50(2): 269-285. doi:10.1051/acarologia/20101970
- Klompen H., Lekveishvili M., Black W.C. 2007. Phylogeny of parasitiform mites (Acari) based on rRNA. *Mol. Phylogenet. Evol.* 43: 936-951. doi:10.1016/j.ympev.2006.10.024
- Klompen H., Vázquez M.M., Bernardi L.F.O. 2015. Post-embryonic development in the mite suborder Opilioacarida, with notes on segmental homology in Parasitiformes. *Exp. Appl. Acarol.*, 67(2): 183-207. doi:10.1007/s10493-015-9939-7
- Krantz G.W., Walter D.E. 2009. *A manual of acarology*. 3rd Edition. Lubbock, TX: Texas Tech University Press.
- Krivolutsky D.A. 1965. A new finding of Opilioacarina in USSR. *Russ. J. Zool.*, 44: 1413.
- Leclerc P. 1989. Considerations paléogéographiques à propos de la découverte en Thaïlande d'Opilioacariens nouveaux (Acari-Notostigmata). *C. R. Séances. Soc. Biogéogr.*, 65(4): 162-174.
- Lehtinen P.T. 1980. A new species of Opilioacarida (Arachnida) from Venezuela. *Acta Biol. Venez.*, 10(2): 205-214.
- Moraza M.L., Prieto C.E., Balanzategui I. 2021. A new species of the genus *Opilioacarus* With, 1902 (Acari: Opilioacarida) for the Iberian Peninsula. *Acarologia*, 61: 128-147. doi:10.24349/acarologia/20214422
- Naudo M.H. 1963. Acariens Notostigmata de l'Angola. *Publicações Cult. Co. Diam. Angola*, 63: 13-24.
- Norton R.A., Kethley J.B., Johnston D.E., O Connor B.M. 1993. Phylogenetic perspectives on genetic systems and reproductive modes of mites. In: Wrensch D.L., Ebbert, M.A. (Eds). *Evolution and diversity of sex ratio in insects and mites*. New York: Chapman & Hall. pp. 8-99. doi:10.1007/978-1-4684-1402-8_2
- Oliver J.H.jr. 1977. Cytogenetics of mites and ticks. *Ann. Rev. Entomol.*, 22: 407-429. doi:10.1146/annurev.en.22.010177.002203
- Oliver J.H.jr. 1989. Biology and systematics of ticks (Acari: Ixodida). *Ann. Rev. Ecol. Syst.*, 20: 397-430. doi:10.1146/annurev.es.20.110189.002145

- Pepato A.R., da Rocha C.E.F., Dunlop J.A. 2010. Phylogenetic position of the acariform mites: sensitivity to homology assessment under total evidence. *BMC Evol. Biol.*, 10: 235. doi:10.1186/1471-2148-10-235
- Silvestri F. 1905. Note Aracnologiche III. *Redia*, 2: 239-261.
- Thaler K., Knoflach B. 2002. Neue *Opilioacarus*-Funde (Acari, Notostigmata) in Peloponnes und Ägäis (Griechenland). *Entomolog. Nachr. Ber.*, 46: 271-277.
- Van der Hammen L. 1966. Studies on Opilioacarida (Arachnida) I. Description of *Opilioacarus texanus* (Chamberlin, Mulaik) and revised classification of the genera. *Zool. Verh.*, 86: 1-80.
- Van der Hammen L. 1969. Studies on Opilioacarida (Arachnida) III. *Opilioacarus platensis* Silvestri, and *Adenacarus arabicus* (With). *Zool. Meded., Leiden*, 44(8): 113-131.
- Van der Hammen L. 1970. La Phylogénèse des Opilioacarides, et leurs affinités avec les autres Acariens. *Acarologia*, 12(3): 465-473.
- Van der Hammen L. 1977. Studies on Opilioacarida (Arachnidea). IV The genera *Panchaetes* Naudou and *Salfacarus* gen. nov. *Zool. Meded., Leiden*, 51(4): 43-78.
- Vázquez M.M., Palacios-Vargas J.G. 1988. Algunas observaciones sobre el comportamiento de los acaros opilioacaridos (Acarida: Notostigmata). *Rev. Nicar. Entomol.*, 6: 1-6.
- Vázquez M.M., Klompen H. 2002. The family Opilioacaridae (Acari: Parasitiformes) in North and Central America, with description of four new species. *Acarologia*, 42(4): 299-322.
- Vázquez M.M., Klompen H. 2009. New species of New World Opilioacaridae (Acari: Parasitiformes) with the description of a new genus from the Caribbean region. *Zootaxa*, 2061: 23-44. doi:10.11646/zootaxa.2061.1.2
- Vázquez M.M., Klompen H. 2010. The genus *Salfacarus* (Acari: Opilioacarida) in Madagascar. *Zootaxa*, 2482: 1-21. doi:10.11646/zootaxa.2482.1.1
- Vázquez M.M., Araújo M.S., Feres R.J.F. 2014. A new genus and two new species of Opilioacaridae (Acari: Parasitiformes) from Amazonia, Brazil with a key to the world genera. *Zootaxa*, 3814(2): 151-176. doi:10.11646/zootaxa.3814.2.1
- Vázquez M.M., Klompen H. 2015. The family Opilioacaridae (Parasitiformes: Opilioacarida) in Mexico, description of two new species and notes on biology and geographical distribution. *Zootaxa*, 3957(5): 535-552. doi:10.11646/zootaxa.3957.5.3
- Vázquez M.M., Araújo M.S., Feres R.J.F. 2015. *Brasilacarus cocaris* (Acari: Opilioacaridae), a new genus and species from Amazonia, Brazil. *Zootaxa*, 3915(3): 375-389. doi:10.11646/zootaxa.3915.3.3
- Vázquez M.M., May D., Alamilla E., Klompen H. 2018. A new species of Opilioacaridae (Parasitiformes: Opilioacarida) from Belize with some observations on life history and behavior. *Syst. Appl. Acarol.*, 23(1): 132-144. doi:10.11158/saa.23.1.11
- Walter D.E., Proctor H.C. 1998. Feeding behaviour and phylogeny: Observations on early derivative Acari. *Exp. Appl. Acarol.*, 22(1): 39-50. doi:10.1023/A:1006033407957
- Walter D.E., Harvey M.S. 2009. Order Opilioacarida. In: Krantz G.W., Walter D.E. (Eds.). *A manual of acarology*, Lubock, TX: Texas Tech University Press. pp. 104-106.
- With C.J. 1902. A new acaride *Opilioacarus segmentatus*. *Forhandlingar vid Nordiska Naturforskareog Lakaremotet i Helsingfors den 7 till 12 Juli 1902 (Comptes Rendus du Congrès des Naturalistes et Médecins du Nord tenu à Helsingfors), Sektionen för Zoologi*, 20: 4-5.
- With C.J. 1904. The Notostigmata, a new suborder of Acari. *Vidensk. Meddel. Naturhist. Foren. Kjøbenhavn.*, 137-192.