Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Supporting agricultural research
for sustainable development

**Acarologia is under free license and distributed under the terms of the
Creative Commons-BY**

Oldest determined record of a mesostigmatic mite (Acari: Mesostigmata: Sejidae) in Cretaceous Burmese amber

Omid Joharchi ^a, Dmitry D. Vorontsov^b, David Evans Walter^c

^a Institute of Environmental and Agricultural Biology (X-BIO), Tyumen State University, Tyumen, Russia.

^b Institute of Developmental Biology, Russian Academy of Sciences, Moscow, Russia.

^c School of Science, Technology and Engineering, University of the Sunshine Coast, 90 Sippy Downs Drive, Sippy Downs, Queensland 4556, Australia.

Original research

ABSTRACT

This paper describes the first record of the Sejida (= Sejina) family Sejidae (superfamily Sejoidea) from Burmese amber, based on deutonymphs. Specimens were found in the amber piece along with a wasp (Hymenoptera: Scelionidae), flies (Diptera: Ceratopogonidae) and thrips (Thysanoptera: Adiheterothripidae). This record traces the Sejida (= Sejina) back to the mid-Cretaceous (ca. 100 Ma) and represents not only the oldest valid record of Mesostigmata, but it is also one of the oldest examples of the entire Parasitiformes clade.

Keywords fossil record; Myanmar; Sejida; Parasitiformes

Introduction

The Sejida (= Sejina) can be subdivided into two superfamilies: Heterozeroconioidea and Sejoidea, the latter comprising four families (Sejidae, Ichthyostomatogasteridae, Uropodellidae, Regimacharlotiidae), ten genera and over 60 described species of mites generally associated with woody substrates, and often with deutonymphs that are phoretic on insects (Walter 2013; Lekveishvili and Klompen 2004). The fossil record of Mesostigmata is surprisingly sparse. Only nine fossil families from the lineages Gamasina and Uropodina (Mesostigmata: Monogynaspida) have been recorded and no valid fossils of the seemingly more early derivative Sejida and Trigynaspida have been found (Dunlop *et al.* 2018; Sidorchuk 2018). Koch and Berendt (1854) reported a sejid fossil species from Baltic amber, but Dunlop *et al.* (2018) recently determined that it was a misinterpretation of an unknown species of cohort Anystina (Acariformes: Prostigmata). All recorded fossil mesostigmatans are from the Cenozoic period [from Palaeogene (Eocene, ca. 44–49 Ma) to Neogene (Miocene, ca. 16 Ma)] and among them only three species were named formally (Dunlop *et al.* 2013; Dunlop *et al.* 2014; Dunlop *et al.* 2018; Sidorchuk 2018). Mesostigmata (both Gamasina and Uropodina) are listed from a number of Cretaceous amber localities, beginning with the Lower Cretaceous Lebanese amber (Rasnitsyn *et al.* 2016), but neither of them was described. Here, we illustrate and describe two deutonymphs from mid-Cretaceous Burmese amber (Kachin) that belong to the Sejida (= Sejina) family Sejidae (superfamily Sejoidea). They thus become the oldest described mesostigmatic mites.

Received 10 June 2021
Accepted 18 August 2021
Published 14 September 2021

Corresponding author
Omid Joharchi
j.omid2000@gmail.com

Academic editor
Faraji, Farid

[https://doi.org/
10.24349/goj5-BZms](https://doi.org/10.24349/goj5-BZms)

ISSN 0044-586X (print)
ISSN 2107-7207 (electronic)

Copyright
Joharchi O. *et al.*

Distributed under
Creative Commons CC-BY 4.0

Figure 1 General view, focus-stacked micrographs of two deutonymph specimens (PIN 5620-26 F & E) of Sejidae in Cretaceous Burmese amber; A – Piece of amber, dorsal and caudal views; B – lateral view of the specimen E; C – dorsal view of both specimens.

Material and methods

The piece of Cretaceous amber containing the fossil mites described herein was mined in the Hukawng Valley (Kachin State, northern Myanmar). The amber locality is radiometrically dated to 98.79 ± 0.62 Ma based on U–Pb zircon dating of the volcanoclastic matrix and shown to be earliest Cenomanian in age (Shi *et al.* 2012; Smith and Ross 2018; Zhang *et al.* 2018; Yu *et al.* 2019).

Amber preparation

The amber piece containing the mite was cut from a larger piece of amber and fine-polished following the protocols of Sidorchuk (2011, 2013) and tools described by Sidorchuk and Vorontsov (2018). The piece was shaped as a parallelepiped ca. $0.5 \times 0.5 \times 0.25$ mm (Figure 1A). We polished the piece from six sides, trying to leave not more than $100 \mu\text{m}$ of amber between the mites and the surface in the dorso-ventral direction, as the thicker layer of semi-transparent amber did not let us see enough details while the close polishing allowed us to use high-resolution water and oil immersion optics. From the six available views, only three were found useful: dorsal, ventral and one of the lateral views.

Imaging

To image the mites two compound microscopes, a Nikon E-800 with water (40x and 60x) and oil (100x) immersion optics and a Zeiss AxioImager A2 with oil immersion optics (100x) were used, with brightfield illumination and differential interference contrast (DIC). To image deeper into the piece, in addition to the standard technique with immersion oil separated from the amber by a coverslip, we used a saturated fructose solution, as described in Khaustov *et al.* (2021). Stacks of images, comprising multiple focal planes (up to 200 in a single stack), were obtained with an Olympus OM-D E-M10-II digital camera. Figures herein contain images assembled from multiple focal planes using Helicon Focus 7.5 software, using algorithms A and B with subsequent manual addition of significant details from the individual focal planes. Some of the resulting original stacks are available through Figshare (<https://doi.org/10.6084/m9.figshare.14707077>). We also provide a link to the source image stacks of the type material studied by Dunlop *et al.*, (2018): <https://doi.org/10.6084/m9.figshare.c.4066367>.

Terminology

The nomenclature used for anatomical structures follows Evans and Till (1979).

Material and repository

The piece containing the two mites is housed in the Öhm-Kühnle collection maintained permanently at the A.A. Borissiak Paleontological Institute, Russian Academy of Sciences (PIN RAS), Moscow. The piece of amber is stored in a plastic vial filled with water; thymol and myramistin (benzyl dimethyl [3-(myristoilamino) propyl] ammonium chloride) are added to the water to limit fungal and bacterial growth. Due to the minute size of the piece (0.5 mm), extreme care is needed when manipulating it.

Two mites (catalog number PIN 5620-26, inclusions E and F) lie very close to each other in a thin layer of darkened amber (probably, at the edge of amber layers) (Figure 1). Each mite is curved (L- shaped), with the dorsal side outwards (see Figure 1B, lateral view). Between the mites there is a structure of presumably plant origin, resembling stellate hairs (leaf trichomes) (Figure 1C). The amber is full of microscopic particles and bubbles that obscure the view of mites.

Arthropod syninclusions that were present in the original piece along with the mites:

PIN 5620-26 A Hymenoptera Scelionidae

PIN 5620-26 B, C, D Diptera Ceratopogonidae

PIN 5620-26 G Thysanoptera Adiheterothripidae

Systematic paleontology

Parasitiformes Reuter, 1909

Mesostigmata G. Canestrini, 1891

Sejida Kramer, 1885

Sejoidea Berlese, 1885

Sejidae Berlese, 1885

Description

Deutonymph (n= 2)

(Figures 1–5).

Dorsal Idiosoma — Oval in outline, dorsal surface smooth, without any distinct ornamentation; podonotal, mesonotal and pygidial shields indistinct, all dorsal setae smooth and relatively long, setae J5/Z5 insert on a small tubercles posteriorly (Figure 2A, D).

Figure 2 Details of the fossil mite, selected images of specimens PIN 5620-26 F & E; A – general view of idiosoma (specimen F), selectively combined from several image stacks (the images were taken from the dorsal side through the whole mite, and some ventral structures are shown); B, C – leg IV of specimen E, manual focus-stacking, together with respective interpretative drawing (trochanter-tarsus) to show intercalary sclerite, note that the leg is bent under the dorsal shield and viewed from the dorsal side through the shield (its border is shown in dotted line); D – posterior part with a focus on tubercles (specimen E). Abbreviations: *ao*, anal opening; *at*, apicotarsus; *gt?*, gnathotectum; *ics*, intercoxal shield; *is*, intercalary sclerite; *per*, peritreme; *st*, stigma; *tub*, tubercles.

Figure 3 Gnathosoma of the fossil mite; A, B – subcapitulum of specimen E, manual focus-stacking, together with respective interpretative drawing; C, D – subcapitulum of specimen F, manual focus stacking, together with respective interpretative drawing. Abbreviations: *ch*, chelicera; *cor*, corniculus; *im*, internal mala; *lab*, labrum; *pc*, palpcoxal seta; *ts*, tritosternum; *ss?*, salivary stylet.

Figure 4 Ventral idiosoma of specimen F; A – interpretative drawing; B – focus-stacked image of intercoxal shield. Abbreviations: *per*, peritreme; *st*, stigma; *st1-4*, sternal setae 1-4.

Ventral idiosoma — Tritosternal laciniae free, base expanded laterally, anchor-shaped (Figure 3A–D), presternal shields absent, intercoxal shield present, somewhat sclerotized, bearing sternal setae *st1-4*, ventrianal shield interlocks with intercoxal shield (Figure 4A, B), anal opening distinct posteriorly (Figure 2A), peritreme on ventral margin of shield extending from between coxae III–IV to mid-level of coxa II (Figures 2A and 4A), peritrematic shields absent.

Gnathosoma — (Figure 3A–D). Chelicerae short and robust, digits indistinct, corniculi appressed anteriorly and flattened, gnathotectum indistinct.

Legs — Leg setae smooth and simple, a distal subdivision (apicotarsus or acrotarsus) and ambulacrum of tarsus I present (Figure 5 A–D); tarsus IV with an intercalary sclerite bearing pair of ventral setae (*av4*, *pv4*) (Figure 2B, C).

Figure 5 Distal portion of tarsus I; A, B – specimen E, manual focus-stacking, together with respective interpretative drawing; C, D – specimen F, manual focus stacking, together with respective interpretative drawing to show apicotarsus (distal subdivision of the tarsus). Abbreviations: *amb*, ambulacrum; *apo*, apotele (pretarsus); *at*, apicotarsus (acrotarsus).

Remarks

The mites are both deutonymphs because of the lack of genital shields and presence of an intercoxal shield bearing four pairs of setae. This species is assigned to the Sejidae (Mesostigmata: Sejida) by the following combination of characters: (1) a distal subdivision (apicotarsus or acrotarsus) and ambulacrum of tarsus I present (Figure 5 A–D), (2) tarsus IV with an intercalary sclerite that bears a pair of ventral setae (*av4*, *pv4*) (Figure 2B, C), (3) dorsal setae *J5/Z5* insert on small tubercles (Figure 2A, D), (4) tritosternum with a broad base, anchor-shaped (Figure 3A, B). Unfortunately, we were unable to see the modification of anterior hypostomal setae (*h1*) as an apomorphic character (it is typically membranous and often highly reduced in Sejidae), nevertheless the above-mentioned diagnostic character states present enough arguments to consider these mites members of Sejidae. We are unable to claim a certain genus for two reasons: (1) modern genera are based mostly on adult females and immature stages have not been well studied, (2) sejid genera are not well studied or resolved. For example, here we would need adult females to check whether the sternal platelets bearing *st1* and *st2* and the posterior mesonotal shields are fused together or not (neither are fused in *Sejus*). The small posterior tubercles associated with setae *J5/Z5* support an association with the genus *Sejus*, which usually have similar but larger tubercles in the adults, to which we tentatively assign these specimens. The genus *Adenosejus* Lekveishvili & Krantz is unlikely because of the absence of thorn-like leg setae; the bifurcate gnathotectum characteristic of *Epicroseius* Berlese cannot be seen; however, an unknown modern or ancient genus cannot be ruled out in this poorly studied group.

Discussion

Fossil mesostigmatid mites are extremely rare. A few cases of Mesostigmata had been reported from the Cretaceous (e.g. Poinar 1998), but none of them have been formally confirmed as Mesostigmata. Our present discovery of the family Sejidae from Burmese amber is significant for representing the oldest and first determined example of the Sejida from the Cretaceous. Deutonymphs and adults of many families of Mesostigmata have established close phoretic relationships with other arthropods, and phoresy by deutonymphs is a common phenomenon among modern Sejidae, typically on beetles associated with dead wood, and it is possible that this pair of mites were dislodged from an insect that was ensnared by the amber. However, we were unable to see an anal disk or a remnant pedicel, and the only insects in this amber piece are very small, so how these mites became entrapped remains a mystery.

Acknowledgements

We cordially thank Christoph Öhm-Kühnle (Herrenberg, Germany) for donating the amber specimen from his collection to PIN RAS, which allowed us to study it. We are grateful to Alexander Rasnitsyn and Alexey Shmakov (PIN RAS) for identification of syninclusions. We express our sincere appreciation to Owen D. Seeman (Queensland Museum, South Brisbane, Queensland, Australia) and another anonymous reviewer for their invaluable comments on an earlier draft of this paper.

References

- Berlese, A. 1885. Acarorum systematis specimen. Bollettino della Società Entomologica Italiana, 17: 121-135.
- Canestrini G. 1891. Abbozzo del sistema Acarologico. Atti 1st. Ven, 38: 699-725.
- Evans G.O., Till W.M. 1979. Mesostigmatic mites of Britain and Ireland (Chelicerata: Acari-Parasitiformes). An introduction to their external morphology and classification. Transactions of the Zoological Society of London, 35: 145-270. <https://doi.org/10.1111/j.1096-3642.1979.tb00059.x>

- Dunlop J.A., Kontschán J., Zwanzig, M. 2013. Fossil mesostigmatid mites (Mesostigmata: Gamasina, Microgyniina, Uropodina), associated with longhorn beetles (Coleoptera: Cerambycidae) in Baltic amber. *Naturwissenschaften*, 100: 337-334. <https://doi.org/10.1007/s00114-013-1031-8>
- Dunlop J.A., Kontschán J., Walter D.E., Perrichot V. 2014. An ant associated mesostigmatid mite in Baltic amber. *Biology Letters*, 10(9): 20140531. <https://doi.org/10.1098/rsbl.2014.0531>
- Dunlop J.A., Walter D.E., Kontschán J. 2018. A putative fossil sejid mite (Parasitiformes: Mesostigmata) in Baltic amber re-identified as an anystine (Acariformes: Prostigmata). *Acarologia*, 58: 665-672. <https://doi.org/10.24349/acarologia/20184263>
- Khaustov A.A., Vorontsov D.D., Perkovsky E.E., Lindquist E.E. 2021. Review of fossil heterostigmatic mites (Acari: Heterostigmata) from late Eocene Rovno Amber. I. Families Tarsocheylidae, Dolichocybidae and Acarophenacidae. *Systematic & Applied Acarology*, 26(1): 33-61. <https://doi.org/10.11158/saa.26.1.3>
- Koch C.L., Berendt G.C. 1854. Die im Bernstein befindlichen Myriapoden, Arachniden und Apteren der Vorwelt. In: Berendt G.C. (Ed.) *Die in Bernstein befindlichen organischen Reste der Vorwelt gesammelt in Verbindung mit mehreren bearbeitet und herausgegeben* 1. Berlin, Nicolai, 124 pp.
- Kramer P. 1885. Über Halarachne Halichoeri Allm. *Zeitschrift für die Gesamten Naturwissenschaften*, 58: 46-74.
- Lekveishvili M., Klompen H. 2004. Phylogeny of infraorder Sejina (Acari, Mesostigmata). *Zootaxa*, 629: 1-19. <https://doi.org/10.11646/zootaxa.629.1.1>
- Poinar Jr G.O. 1998. Fossils explained 22: Palaeontology of amber. *Geology Today*, July-August 1998: 154-160. <https://doi.org/10.1046/j.1365-2451.1998.014004154.x>
- Rasnitsyn A.P., Bashkuev A.S., Kopylov D.S., Lukashovich E.D., Ponomarenko A.G., Popov Y.A., Rasnitsyn D.A., Ryzhkova O.V., Sidorchuk E.A., Sukatsheva I.D. 2016. Sequence and scale of changes in the terrestrial biota during the Cretaceous (based on materials from fossil resins). *Cretaceous Research*, 61: 234-255. <https://doi.org/10.1016/j.cretres.2015.12.025>
- Reuter E. 1909. Zur Morphologie und Ontogenie der Acariden mit besonderer Berücksichtigung von *Pediculopsis graminum*. *Acta Soc. Sci. Fenn.*, 36: 1-288. <https://doi.org/10.5962/bhl.title.4705>
- Shi G., Grimaldi D.A., Harlow G.E., Wang J., Wang J., Yang M., Lei W., Li Q., Li X., 2012. Age constraint on Burmese amber based on U-Pb dating of zircons. *Cretaceous Research* 37, 155 e163. <https://doi.org/10.1016/j.cretres.2012.03.014>
- Sidorchuk E.A. 2011. Preparation of six-sided micro-samples of Baltic amber for study of organismal inclusions. In: Kostyashova, Z.V. (Eds.), *Amber Mining and Processing in Sambia*, International Symposium Materials, 12-14 May 2010, Kaliningrad, pp. 47-53.
- Sidorchuk, E.A. 2013. New technique for preparation of small-sized amber samples with application to mites. In: Azar D., Engel M.S., Jarzembowski E., Krogmann L., Nel A., Santiago-Blay J.A. (Eds.), *Insect evolution in an Amberiferous and Stone Alphabet*. Proceedings of the 6th International Congress on Fossil Insects, Arthropods and Amber; Brill: Leiden-Boston. pp. 189-201.
- Sidorchuk, E.A. 2018. Mites as fossils: forever small?. *International Journal of Acarology*, 44(8): 349-359. <https://doi.org/10.1080/01647954.2018.1497085>
- Sidorchuk E.A., Vorontsov D.D. 2018. Preparation of small-sized 3D amber samples: state of the technique. *Palaeontology*, 1: 80-90. <https://doi.org/10.11646/palaeontology.1.1.10>
- Smith R.D.A., Ross A.J., 2018. Amberground pholadid bivalve borings and inclusions in Burmese amber: implications for proximity of resin-producing forests to brackish waters, and the age of the amber. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 107 (2-3): 239-247. <https://doi.org/10.1017/S1755691017000287>
- Walter D.E. 2013. A new genus and family of sejine mites (Acari, Parasitiformes, Mesostigmata, Sejoidea) based on new species from Lord Howe Island and Brazil, and a redescription of *Sejus americanus* (Banks, 1902). *Zootaxa*, 3691: 301-323. <https://doi.org/10.11646/zootaxa.3691.3.1>
- Yu T.T., Kelly R., Mu L., Ross A., Kennedy J., Broly P., Xia F.Y., Zhang H.C., Wang B., Dilcher D. 2019. An ammonite trapped in Burmese amber. *Proceeding of the National Academy of Sciences of United States of America*, 116 (23): 11345-11350. <https://doi.org/10.1073/pnas.1821292116>
- Zhang Q., Rasnitsyn A.P., Wang B., Zhang H.C. 2018. Hymenoptera (wasps, bees and ants) in mid-Cretaceous Burmese amber: A review of the fauna. *Proceedings of the Geologists' Association*, 129: 736-747. <https://doi.org/10.1016/j.pgeola.2018.06.004>