Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Supporting agricultural research
for sustainable development

**Acarologia is under free license and distributed under the terms of the
Creative Commons-BY**

Teneriffia hajiqaanbari sp. nov. (Acari: Trombidiformes: Teneriffiidae), first record of the genus from Iran, with a key to world species of *Teneriffia*

Saeid Paktinat-Saeij ^{a,b}, Shahrooz Kazemi ^c^a Department of Plant Protection, Faculty of Agriculture, University of Tehran, Karaj, Iran.^b Zeraat Gostar Poursoleyman Company, Babol, Mazandaran province, Iran.^c Department of Biodiversity, Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman, Iran.

Original research

ABSTRACT

Teneriffia hajiqaanbari n. sp. is described based on female, male, and deutonymph specimens collected in mangrove forests in Qeshm Island, in the eastern part of the Persian Gulf, southern Iran. This new species can be distinguished by following characters: gnathosoma without “clasps” ventrally; palpal onychophysis absent; venter with 17–20 pairs of setae; genital plates with seven pairs of setae (female); dorsal body setae minutely barbed; opisthosomal setae d_1 , e_1 and f_1 long and reaching or surpassing the base of next seta; solenidotomy of tarsi I–IV: 3-3-0-1. Additionally, an identification key to known species of *Teneriffia* is updated.

Keywords Prostigmata; predatory mites; littoral mites; taxonomy; Hara forests**Zoobank** <http://zoobank.org/891A36CE-2BFC-4474-AC8D-95CA8AC0C673>

Introduction

Members of the family Teneriffiidae are red, yellow, or brownish long-legged predators that are fast-moving and medium-sized mites (Walter *et al.* 2009; Luxton 1993). Teneriffiids occupy many different habitats, ranging from the sea level to altitudes about 2,000 m in terrestrial and marine ecosystems, including intertidal sandbanks, undersides of rocks, caves, arid desert, plants, soil, and forest litter (Walter *et al.* 2009; Khanjani *et al.* 2011; Ueckermann and Durucan 2020).

The Teneriffiidae comprises about 27 species in the nine genera: *Austroteneriffia* Womersley, 1935 (9 or 10 species); *Heteroteneriffia* Hirst, 1925 (3 species); *Himalteneriffia* Schmölzer, 2002 (1 species); *Mesoteneriffia* Irk, 1939 (2 species); *Mesoteneriffiola* Schmölzer, 1956 (1 species); *Neoteneriffiola* Hirst, 1924 (5 species); *Parateneriffia* Thor, 1911 (1 species); *Sinoteneriffia* Yin *et al.*, 1994 (1 species) and *Teneriffia* Thor, 1911 (4 species) (Schmölzer 2002; Ueckermann and Durucan 2020; Zmudzinski *et al.*, 2021). The genus *Teneriffia* was erected by Thor (1911), with *Teneriffia quadripapillata* Thor, 1911, as the type species collected from Spain (Canary Islands). Until the present study, four species of this genus have been described from the world. The genus *Teneriffia* can be distinguished from other genera of the family by having a simultaneously prodorsal shield, coxae I–IV at least with five setae each and strongly bipectinated claws on tarsi I–II (Schmölzer 2002; Ueckermann and Durucan 2020).

During studies on mite fauna of the mangrove forests of southern Hormozgan province, a new species of *Teneriffia* was discovered. Herein, we describe *T. hajiqaanbari* n. sp., the first

Received 21 November 2021

Accepted 10 February 2022

Published 21 February 2022

Corresponding author

Academic editor
Auger, Philippe<https://doi.org/10.24349/jkco-c4nu>ISSN 0044-586X (print)
ISSN 2107-7207 (electronic)

Paktinat-Saeij S. and Kazemi S.

Licensed under
Creative Commons CC-BY 4.0

How to cite this article Paktinat-Saeij S. and Kazemi S. (2022), *Teneriffia hajiqaanbari* sp. nov. (Acari: Trombidiformes: Teneriffiidae), first record of the genus from Iran, with a key to world species of *Teneriffia*. *Acarologia* 62(1): 262–269. <https://doi.org/10.24349/jkco-c4nu>

species of *Teneriffia* recorded from Iran. Also, an updated key to the world species of *Teneriffia* is presented.

Material and methods

Mites were extracted from littoral and soil samples collected in mangrove forests in southern Iran using Berlese-Tullgren funnels, cleared in Nesbitt's fluid and mounted in Hoyer's medium. Morphological observations, measurements and illustrations were made using a compound microscope equipped with differential interference contrast and phase contrast optical systems and a drawing tube (Olympus BX51). Initial pencil line drawings were then scanned and cleaned using Adobe Illustrator CS6. Additional images were prepared using the automated Z-stacking feature of the Nikon NIS Elements package on a Nikon Eclipse 90i (Melville, NY) compound microscope with a PC controlled Ds-5M-U1 digital camera at the Acarology Laboratory of the Ohio State University (OSAL). Measurements are in micrometers (μm). The body length and width of the specimens were measured from the suture between the gnathosoma and idiosoma to the posterior margin of the idiosoma and at its broadest level, respectively. The length of the gnathosoma was taken from the base to the tip of the subcapitulum. The leg lengths were measured from the ventral insertion of the coxae to the base of the pretarsi. Measurements of the paratypes are given in brackets following that of the holotype. Terminology generally follows that of Ueckermann and Durucan (2020) and Kethley (1990).

Results

Family Teneriffiidae Thor, 1911

Genus *Teneriffia* Thor, 1911

Type species *Teneriffia quadripapillata* Thor, 1911

Teneriffia hajiqaanbari n. sp.

Zoobank: [BB4B55F2-503B-4B4F-AC96-25C3CC99F88D](https://zoobank.org/BB4B55F2-503B-4B4F-AC96-25C3CC99F88D)

(Figures 1–3)

Diagnosis

Gnathosoma without “clasps” ventrally; palpal oncofysis absent; venter with 17–20 pairs of setae; genital plates with seven pairs of setae (female); dorsal body setae minutely barbed; opisthosomal setae d_1 , e_1 and f_1 long and reaching or surpassing the base of next seta; solenidotaxy of tarsi I–IV: 3-3-0-1.

Description

Female (n=4)

Body length 782 (728–748), width 534 (446–510); length of subcapitulum 168 (164–188).

Dorsal idiosoma — (Figures 1A, 1B). Color red in life, idiosoma oval. Integument striated. Dorsum with one clear propodosomal shield 314 (317–324) long, 178 (173–183) wide, with broken longitudinal striae and two pairs of minutely barbed setae (*ve*, *sce*) and two pairs of ciliated trichobothria (*vi*, *sci*), the bases of which have a rosette pattern, trichobothrium *vi* on naso and trichobothrium *sci* near seta *ve* (Figures 1A, 1B). Two pair of eyes presented anteriad setae c_2 . Stigmatic opening anterior to setae *vi* and near to base of chelicera.

Hysterosoma with seven pairs of setae (c_1 , c_2 , d_1 , e_1 , f_1 , h_1 and h_2); all setae minutely barbed (Figures 1A, 1B). Setae c_2 much longer than other dorsal setae. Three pairs of cupules (*ia*, *im* and *ip*) present. Lengths of dorsal setae: *vi* 84 (87–92), *ve* 114 (106–116), *sci* 124 (124–126), *sce* 124 (129–136), c_1 67 (67–69), c_2 178 (168–186), d_1 111 (92–104), e_1 101

Figure 1 *Teneriffia hajiqaanbari* n. sp., female, A – dorsal view of idiosoma; B – seta *c*₁ and trichobothria *sci*; C – ventral view of idiosoma; D – subcapitulum and palp; E – peritreme; F – chelicera.

Figure 2 *Teneriffia hajiqaanbari* n. sp., female, A – leg I; B – leg II; C – leg III; D – leg IV.

(94–109), f_1 126 (114–129), h_1 126 (106–124), h_2 69 (75–84). Distances between dorsal setae: $ve-ve$ 153 (136–156), $sci-sci$ 144 (126–144), $sce-sce$ 149 (126–137), c_1-c_1 148 (119–139), c_1-c_2 64 (52–55), c_2-c_2 277 (235–248), c_1-d_1 106 (65–74), d_1-d_1 198 (158–198), d_1-e_1 99 (79–90), e_1-e_1 156 (125–134), e_1-f_1 104 (89–92), f_1-f_1 74 (69–77), f_1-h_1 104 (74–89), h_1-h_1 37 (42–63), h_1-h_2 38 (59–77), h_2-h_2 139 (141–186).

Ventral idiosoma — (Figures 1C). Ventral surface ornamented with striations. Coxae I–II separated from coxae III–IV. Genital plates with seven pairs of setiform genital setae (g_{1-7}), about 20–25 in length (measurement range in holotype and paratypes). Ventral region with 17–20 pairs of aggenital setae (ag) about 27–42 in length (measurement range in holotype and

paratypes), all serrate. Three pairs of genital papillae present. Anal plates with four pairs of pseudoanal setae (ps_{1-4}). Lengths of setae: ps_1 50 (45–57), ps_2 42 (47–59), ps_3 59 (50–52), ps_4 50 (45–52).

Gnathosoma — (Figures 1D–1F). Subcapitulum finely lineated, bearing one pair of subcapitular setae m 67 (62–69) and three pairs of adoral setae (or_{1-3}), or_1 35 (34–37), or_2 12 (11–12), or_3 15 (15–16) (Figure 1D). Peritreme located at cheliceral base (Figure 1E). Chelicera 198 (193–198) long; bearing two dorsal setae cha and chb (Figure 6). Palp (Figure 1F) 188 (198–202) long; five-segmented; trochanter short, without setae; femur and genu each with one seta; tibia with one robust terminal spur (o_1) 54 (53–56), two subterminal spurs (o_{2-3}), 17 (15–16) and 15 (14–16) long, plus one serrated seta; tarsus with one solenidion and eight barbed or smooth setae. Oncophysis on genu absent.

Legs — (Figures 2A–2D). Leg IV much longer than body; each side of claws on tarsus I with 21–27 pectinations and tarsus II with 16–20 pectinations, tarsi III–IV with smooth tarsal claws and one median empodium. Measurements of leg segments as follows: Ta I 176 (183–189), Ti I 144 (131–141), Ge I 104 (101–105), TF I 99 (94–101), BF I 106 (99–104), Tr I 82 (74–77), Cx I 198 (171–193), leg I 909 (858–900); Ta II 173 (153–178), Ti II 129 (124–126), Ge II 97 (87–92), TF II 101 (87–89), BF II 99 (87–90), Tr II 84 (79–84), Cx II 181 (173–176), leg II 864 (809–825); Ta III 220 (228–230), Ti III 149 (131–141), Ge III 99 (97–99), TF III 104 (94–99), BF III 94 (87–92), Tr III 99 (94–104), Cx III 178 (170–186), leg III 943 (908–942); Ta IV 277 (255–278), Ti IV 176 (178–183), Ge IV 129 (124–127), TF IV 124 (116–126), BF IV 87 (89–94), Tr IV 104 (116–119), Cx IV 193 (173–188), leg IV 1090 (1064–1086).

Chaetotaxy of legs I–IV: coxae 6/7/8-6/7-6/7-5; trochanters 1-2-2-2; basifemora 5-6-4-4; telofemora 5-5-4-4; genua 8(+1 σ)-8(+1 σ)-7(+1 σ)-6; tibiae 12(+1 ϕ +1 k)-12(+1 ϕ)-12(+1 ϕ)-11(+1 ϕ); tarsi 27(+3 ω)-27(+3 ω)-22+1Tr-23(+1 ω)+1Tr.

Male (n=4)

(Figures 3A–3D)

Body length 598–756, width 402–471; length of subcapitulum 164–168.

Dorsal idiosoma — Color red in life, idiosoma oval. Integument striated. Dorsum with one clear propodosomal shield 257–302 long, 149–173 wide, with broken longitudinal striae and two pairs of minutely barbed setae (ve , sce) and two pairs of slightly bipectinated trichobothria (vi , sci), trichobothrium vi on naso and trichobothrium sci near seta ve . Two pairs of eyes presented anteriad of setae c_2 . Stigmatic opening anterior to setae vi and near to base of chelicera.

Hysterosoma with seven pairs of setae (c_1 , c_2 , d_1 , e_1 , f_1 , h_1 and h_2); all setae minutely barbed. Setae c_2 much longer than other dorsal setae. Three pairs of cupules (ia , im and ip) present. Lengths of dorsal setae: vi 77–92, ve 82–107, sci 99–116, sce 99–124, c_1 62–65, c_2 141–173, d_1 75–86, e_1 77–104, f_1 89–116, h_1 99–128, h_2 58–79. Distances between dorsal setae: ve – ve 114–129, sci – sci 109–124, sce – sce 92–125, c_1 – c_1 104–125, c_1 – c_2 50–52, c_2 – c_2 203–265, c_1 – d_1 50–84, d_1 – d_1 116–158, d_1 – e_1 69–98, e_1 – e_1 104–122, e_1 – f_1 74–94, f_1 – f_1 57–75, f_1 – h_1 54–87, h_1 – h_1 32–40, h_1 – h_2 54–69, h_2 – h_2 124–156.

Ventral idiosoma — Ventral surface ornamented with striations. Coxae I–II separated from coxae III–IV. Genital plates with six to eight pairs of setiform genital setae (g), about 20–25 in length (measurement range in paratypes) and internal male genitalia. Ventral region with 17–20 pairs of aggenital setae (ag) about 25–41 in length (measurement range in paratypes), all serrate. Three pairs of genital papillae present. Anal plates with four pairs of pseudoanal setae (ps_{1-4}). Lengths of setae: ps_1 52–59, ps_2 42–54, ps_3 40–45, ps_4 42–52.

Gnathosoma — (Figures 3A–3B). Subcapitulum finely lineated, bearing one pair of subcapitular setae m 57–65 and three pairs of adoral setae (or_{1-3}), or_1 32–35, or_2 11–12, or_3 12–15. Peritreme located at cheliceral base. Chelicera 173–181 long; bearing two dorsal setae cha and chb . Palp 159–198 long; five-segmented; trochanter short, without setae; femur and genu each with one seta; tibia with one robust terminal spur (o_1) 46–53, two subterminal spurs (o_{2-3}), 15–16 and 14–15 long, plus one serrated seta; tarsus with one solenidion and eight barbed or smooth setae. Oncophysis on genu absent.

Figure 3 *Teneriffia hajiqanbari* n. sp., male, A – ventral view of gnathosoma; B – dorsal view of gnathosoma; C – tarsus of leg I; D – tarsi of leg III and IV.

Legs — (Figures 3C–3D). Leg IV longer than body; each side of claws on tarsus I with 20–26 pectinations and tarsus II with 18–21 pectinations, tarsi III–IV with smooth tarsal claws and one median empodium. Measurements of leg segments as follows: Ta I 151–178, Ti I 106–134, Ge I 79–101, TF I 79–99, BF I 76–101, Tr I 70–74, Cx I 129–168, leg I 691–855; Ta II 151–193, Ti II 101–129, Ge II 76–92, TF II 64–87, BF II 69–89, Tr II 70–74, Cx II 151–178, leg II 692–840; Ta III 183–225, Ti III 106–126, Ge III 79–94, TF III 74–101, BF III 69–80, Tr III 74–87, Cx III 146–173, leg III 743–886; Ta IV 223–282, Ti IV 134–173, Ge IV 94–114, TF IV 76–111, BF IV 69–82, Tr IV 84–99, Cx IV 129–176, leg IV 819–1037.

Chaetotaxy of legs I–IV: coxae 6/7-6/7/8-5/6/7-5/6; trochanters 1-2-2-2; basifemora 5-6-4-4; telofemora 5-5-4-4; genua 8/10(+1 σ)-8(+1 σ)-7(+1 σ)-6; tibiae 12/13/14(+1 ϕ +1 k)-12(+1 ϕ)-11/12(+1 ϕ)-11(+1 ϕ); tarsi 27(+3 ω)-27(+3 ω)-22+1Tr-22/23(+1 ω)+1Tr.

Deutonymph (n=1)

Body length 594, width 436; length of subcapitulum 164; palp 174; chelicera 172;

propodosomal shield length 277, width 156. Lengths of dorsal setae: vi 77, ve 87, sci 94, sce 119, c_1 59, c_2 166, d_1 101, e_1 106, f_1 129, h_1 111, h_2 74. Distances between dorsal setae: $ve-ve$ 129, $sci-sci$ 129, $sce-sce$ 124, c_1-c_1 111, c_1-c_2 57, c_2-c_2 230, c_1-d_1 87, d_1-d_1 161, d_1-e_1 67, e_1-e_1 114, e_1-f_1 74, f_1-f_1 69, f_1-h_1 74, h_1-h_1 35, h_1-h_2 50, h_2-h_2 129. Lengths of setae: ps_1 52, ps_2 50, ps_3 42, m 47; or_1 27, or_2 10, or_3 11, o_1 44, o_2 14 and o_3 13.

Deutonymph like female but differs in following characters: venter with two pairs of genital setae, about 18 in length and 11 pairs of aggenital setae about 25–32 in length. Anal plates with three pairs of pseudoanal setae (ps_{1-3}).

Measurements of leg segments as follows: Ta I 149, Ti I 114, Ge I 77, TF I 74, BF I 79, Tr I 74, Cx I 131, leg I 698; Ta II 139, Ti II 94, Ge II 74, TF II 69, BF II 77, Tr II 67, Cx II 129, leg II 649; Ta III 188, Ti III 111, Ge III 74, TF III 71, BF III 69, Tr III 74, Cx III 141, leg III 728; Ta IV 200, Ti IV 134, Ge IV 84, TF IV 82, BF IV 69, Tr IV 87, Cx IV 131, leg IV 787. Each side of claws on tarsus I with 28 pectinations and tarsus II with 23 pectinations.

Chaetotaxy of legs I–IV: coxae 6-6-5-4; trochanters 1-2-2-2; basifemora 5-6-4-3; telofemora 5-5-4-4; genua 8(+1 σ)-7/8(+1 σ)-7(+1 σ)-6; tibiae 12(+1 ϕ +1 k)-12(+1 ϕ)-11/12(+1 ϕ)-11(+1 ϕ); tarsi 25(+1 ω)-25(+2 ω)-19+1Tr-19(+1 ω)+1Tr.

Larvae, Protonymph and Tritonymph: Unknown.

Remarks

At first glance, the new species closely resembles *Teneriffia sebahatae* Ucekerman & Durucan, 2020 in having gnathosoma without “clasps” ventrally, palpal onychophysis absent, and solenidotaxy on legs. However, *T. hajiqaanbari* n. sp. can be easily distinguished from *T. sebahatae* by the following characters: (1) ventral idiosoma with 17–20 pairs of setae (vs. 23 pairs in *T. sebahatae*); (2) genital plates (female) with seven pairs of setae (vs. six pairs in *T. sebahatae*); and (3) opisthosomal setae d_1 , e_1 and f_1 long and reaching or surpassing the base of next seta (vs. short and not reaching in *T. sebahatae*). Also, length of setae ve 114 (106–116), d_1 111 (92–104), e_1 101 (94–109), f_1 126 (114–129) and h_1 126 (106–124) instead of length of setae ve , d_1 , e_1 , f_1 and h_1 = 58–78 in *T. sebahatae*.

Etymology

The species is named in honor of Dr. Hamidreza Hajiqaanbar (17 August 1973–18 Oct 2021), a distinguished and well-known Iranian Acarologist who has made outstanding contributions to Acarology, especially Heterostigmatina.

Type materials

Type material was collected in mangrove forests in Qeshm Island, in the eastern part of the Persian Gulf, southern Hormozgan province, Iran, collected by S. Saberi.

Holotype: female, from soil and broken cockleshells in littoral zone (26°53'25" N, 55°42'57" E), 07 March 2020, will be deposited in the Acarology Collection, Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman, Iran (ACISTE). **Paratypes:** two females and one deutonymph, from soil in littoral zone (26°54'58" N, 55°35'33" E), 03 March 2020, will be deposited in ACISTE; one female, from soil and broken cockleshells in littoral zone (26°52'36" N, 55°35'06" E), 03 March 2020, will be deposited in the Acarological Collection of the Jalal Afshar Zoological Museum, Department of Plant Protection, Faculty of Agriculture, University of Tehran, Karaj, Iran (JAZM); four males, from soil in littoral zone (26°53'37" N, 55°35'26" E), 02 May 2020, will be deposited in ACISTE; one male, from soil in littoral zone (26°53'37" N, 55°35'26" E), 02 May 2020, deposited in Ohio State University Acarology Collection (OSAL: 156027); two males, from soil and broken cockleshells in littoral zone (26°55'27" N, 55°43'42" E), 29 December 2019, will be deposited in ACISTE; one male and one deutonymph, from soil and broken cockleshells in littoral zone (26°49'50" N, 55°43'40" E), 07 March 2020, will be deposited in ACISTE; one male, from littoral soil (26°51'01" N, 55°44'27" E), 11 January 2020, deposited in JAZM.

Key to the females of *Teneriffia* (After Ueckermann and Durcan, 2020)

1. Gnathosoma with “clasps” ventrally *T. quadripapillata* Thor, 1911
— Gnathosoma without “clasps” 2
2. Palpal oncophysis present; venter with seven pairs of setae
..... *T. aethiopica* Zmudzinski *et al.*, 2021
— Palpal oncophysis absent; venter at least with 17 pairs of setae 3
3. Venter with more than 30 pairs of setae; tarsi III and IV with 1–2 and 2–3 solenidia, respectively *T. mexicana* McDaniel *et al.* 1976
— Venter with 17–23 pairs of setae; tarsi III and IV with 0 and 1 solenidium, respectively 4
4. Venter with 17–20 pairs of setae; seven pairs of genital setae present
..... *T. hajiqaanbari* n. sp.
— Venter with 23 pairs of setae; six pairs of genital setae present
..... *T. sebahatae* Ueckerman & Durcan, 2020

Acknowledgements

We wish to thank Sepideh Saberi for her valuable help with this project, especially in sampling the material used in this study. We also thank Hans Klompen for use of the equipment/facilities at the Ohio State University. This work was supported by INSF, under grant number 98015867.

References

- Hirst S. 1924. On three new Acari belonging to the superfamily Trombidioidea (Erythraeidae and Teneriffiidae). *Proc. Zoo. Soc. Lond.*, 94: 1078-1080. <https://doi.org/10.1111/j.1096-3642.1924.tb03329.x>
- Hirst S. 1925. On some new genera and species of Arachnida. *Proc. Zoo. Soc. Lond.*, 1278-1280. <https://doi.org/10.1111/j.1469-7998.1925.tb07436.x>
- Irk V. 1939. Drei neue Milbenarten aus dem Tiroler Hochgebirge. *Zool. Anz.*, 128: 217-223.
- Kethley J. 1990. Acarina: Prostigmata (Actiniedida). In: Dindal D.L. (Ed). *Soil Biology Guide*. New York: John Wiley and Sons. p. 667-756.
- Khanjani M., Asali Fayaz B., Ueckermann E.A. 2011. A new species of the genus *Austroteneriffia* (Acari: Anystina: Teneriffiidae) from western Iran. *Internat. J. of Acarol.*, 37(6): 550-555. doi:10.1080/01647954.2010.528798 <https://doi.org/10.1080/01647954.2010.528798>
- Luxton M. 1993. The genus *Heteroteneriffia* Hirst, 1925 (Acarina: Prostigmata: Teneriffiidae). *Zool. Anz.*, 230: 103-109.
- McDaniel B., Morihara D., Lewis J.K. 1976. The family Teneriffiidae Thor, with new species from Mexico. *Ann. Ent. Soc. Amer.*, 69(3): 522-537. <https://doi.org/10.1093/aesa/69.3.527>
- Schmölzer K. 1956. Landmilben aus dem Dauphiné (Acarina terrestria). *Österr. Zool. Z.*, 6(3/5): 542-565.
- Schmölzer K. 2002. On Teneriffiidae and the description of a new genus and species from the Himalaya (Acarina, Trombidiformes). *Ber. Naturwiss.-med. Ver. Innsb.*, 89: 126-132.
- Thor S. 1911. Eine neue Acarinenfamilie (Teneriffiidae) und zwei neue Gattungen, die eine von *Teneriffia*, die andre aus Paraguay. *Zool. Anz.*, 38, 171-179.
- Ueckermann E.A., Durcan F. 2020. *Teneriffia sebahatae* sp. nov. (Acari: Trombidiformes: Teneriffiidae), the first teneriffiid mite from Turkey. *Syst. Appl. Acarol.*, 25(6): 1139-1146. <https://doi.org/10.11158/saa.25.6.15>
- Walter D.E., Lindquist E.E., Smith I.M., Cook D.R., Krantz G.W. 2009. Order Trombidiformes. In: Krantz G.W., Walter D.E. (Eds). *A Manual of Acarology 3rd Edition*. Lubbock, USA: Texas Tech University Press. p. 233-420.
- Womersley H. 1935. The occurrence in Australia of Acarina of the family Teneriffiidae (Trombidioidea). *Rec. South Aust. Mus.*, 5, 333-338.
- Yin, S., Bei, N. & Li, X. (1994) A new genus and new species of Teneriffiidae from China (Acari: Prostigmata). *Acta Zootaxonomica Sinica*, 19(4), 433-437.
- Zmudzinski, M., Skoracki, M. & Friedrich, S. (2021) A new species of Teneriffiidae (Acariformes: Prostigmata) from Ethiopia. *Int. J. Acarology*, 47(4): 317-326. <https://doi.org/10.1080/01647954.2021.1908422>