

# New data on some *Antennoseius* Berlese species (Acarai: Ascidae) from Russia

Omid Joharchi <sup>a</sup>, Irina I. Marchenko  <sup>b</sup>, Maria K. Stanyukovich<sup>c</sup><sup>a</sup> Institute of Environmental and Agricultural Biology (X-BIO), Tyumen State University, Semakova Str. 10, 625003 Tyumen, Russia.<sup>b</sup> Institute of Systematics and Ecology of Animals, Novosibirsk, Russia.<sup>c</sup> Zoological Institute of Russian Academy of Sciences, Saint Petersburg, Russia.

## Original research

### ABSTRACT

This paper reports on four species of the genus *Antennoseius* Berlese phoretic on ground beetles (Coleoptera: Carabidae) from Russia: *Antennoseius (Vitzthumia) bregetovae* Chelebiev; *Antennoseius (Antennoseius) bullitus* Karg; *A. (A.) pannonicus* Willmann and *A. (A.) ponticus* Trach and Makarova. Moreover, we provides additional morphological information and new illustrations for two species of *Antennoseius* Berlese – *A. (V.) hyperboreus* Nikolsky and *A. (V.) koroljevae* Chelebiev based on the type series to facilitate species delimitation.

**Keywords** Gamasina; Parasitiformes; Ascoidea; phoretic mite; beetle; Asian Russia

## Introduction

A remarkable diversity of phoretic and parasitic mites is found on ground beetles, representing three major groups: the Mesostigmata, Prostigmata, and Astigmatina (Olynyk and Freitag 1979; Hunter and Rosario 1988; Felska *et al.* 2018). Deutonymphs and adults of many families of Mesostigmata have established close phoretic relationships with ground beetles, and phoresy by adult females is common phenomenon among two genera of family Ascidae – *Antennoseius* Berlese and *Anystipalpus* Berlese (Lindquist *et al.* 2009; Trach 2013). *Antennoseius* is well known as a group of predatory mites generally found in soil, litter, moss in meadow, and forests, as well as in nests of rodents and birds but many species are also recorded on carabid beetles as phoretic (Ryke 1962; Costa 1969; Bregetova 1977; Karg 1993; Halliday *et al.* 1998; Eidelberg 2000; Beaulieu *et al.* 2008; Trach and Makarova 2008; Lindquist and Moraza 2009; Trach 2013; Moraza and Kazemi 2009; Kazemi and Moraza 2013; Faraji *et al.* 2017; Kazemi 2018). The genus comprises about 58 nominal species that are recorded worldwide (Moraes *et al.* 2016; Kazemi 2018; Bahrami and Kazemi 2019). Of these species, about 23 species of *Antennoseius* are phoretic on ground beetles (Faraji *et al.* 2017). Before this study, only five species regarded to belong to *Antennoseius* had been reported phoretic on ground beetles from Russia (Bregetova 1977; Faraji *et al.* 2017; Belova and Makarova 2017): *A. (A.) bullitus* Karg, 1969; *A. (V.) multisetus* Eidelberg, 2000; *A. (V.) ovaliscutalis* Eidelberg, 2000; *A. (A.) pseudospinosus* Eidelberg, 1990; *A. (A.) sabulicola* Bregetova, 1977. During a survey of ground beetle-associated gamasid mites in Russia, *Antennoseius (Vitzthumia) bregetovae* Chelebiev; *Antennoseius (Antennoseius) bullitus* Karg; *A. (A.) pannonicus* Willmann and *A. (A.) ponticus* Trach and Makarova were collected. The aim of this paper is to review the morphology of these species, and to add more details to their original descriptions. Moreover, we provides additional morphological information and new illustrations for *A. (V.) bregetovae* Chelebiev; *A. (V.) hyperboreus* Nikolsky and *A. (V.) koroljevae* Chelebiev, species known only from the type series to facilitate species delimitation.

Received 23 February 2022

Accepted 24 March 2022

Published 28 March 2022

Corresponding author

Omid Joharchi 

j.omid2000@gmail.com

Academic editor

Faraji, Farid

[https://doi.org/  
10.24349/4vni-nutt](https://doi.org/10.24349/4vni-nutt)ISSN 0044-586X (print)  
ISSN 2107-7207 (electronic)Joharchi O. *et al.*Licensed under  
Creative Commons CC-BY 4.0**How to cite this article** Joharchi O. *et al.* (2022), New data on some *Antennoseius* Berlese species (Acarai: Ascidae) from Russia. *Acarologia* 62(2): 378-395. <https://doi.org/10.24349/4vni-nutt>

## Material and methods

This study is based on phoretic *Antennoseius* species on ground beetles that were collected during several biological expeditions in Russia over a period of two years (2020–2021) and also on examination of type series of some specimens deposited in the collection of the Institute of Systematics and ecology of animals (ISEA, Novosibirsk), and collection of the Zoological Institute of the Russian Academy of Sciences (ZIN, Saint Petersburg). Host beetles were collected from different habitats by individual hand picking and placed individually in vials with 96% ethanol. Alcohol sediments from the vials were inspected for phoretic mites. Mites were removed from the beetles and alcohol sediments, cleared in lactic acid solution and mounted in Hoyer's medium (Walter and Krantz 2009). The line drawings and examinations of the specimens were performed with Zeiss Axio Imager A2 and Leica DM 2500 compound microscopes equipped with drawing tubes and differential interference contrast and phase contrast optical systems, attached to cameras AxioCam ICc 5 and ICC50 HD, respectively. Most images were captured in stacks (with focal depth manually controlled). Selected images were combined using Helicon Focus 7.6.4 Pro (Helicon Soft Ltd., 2000). Digital drawings were prepared using Adobe Photoshop CS2 software based on the original pencil line drawings. Images and morphological measurements were taken via ZEN 2012 software (version 8.0) and Leica Application Suite (LAS) software (version 4.2, Live and Interactive Measurements modules). Photomicrographs were taken with an AxioCam 506 camera (Carl Zeiss, Germany). Measurements of structures are expressed as ranges (minimum–maximum) in micrometres ( $\mu\text{m}$ ). Podonotal and opisthonotal shields length were taken from anterior to posterior margins as midline and their width, respectively, from lateral margins at the level of dorsal setae *r2* and at level of setae *S1*. Length and width of sternal shield measured at midline and level of *st2*, respectively. The length of the genital shield was measured along the midline from the anterior margin of the hyaline extension to the posterior margin of the shield, and its width where maximal (at level of setae *st5*). Anal shield was measured on midline length from anterior to posterior margins including cribrum and its width at the broadest point. Leg length was measured from the base of the coxa to the apex of the tarsus (excluding the pre-tarsus). The nomenclature used for the dorsal idiosomal chaetotaxy follows that of Lindquist and Evans (1965), the notations for leg and palp setae follow those of Evans (1963a, b), and other anatomical structures mostly follow Evans and Till (1979). Notations for idiosomal pore-like structures (gland pores and poroids/lyrifissures) and peritrematal shield follow mostly Athias-Henriot (1971, 1975). The notations for pore-like structures on the sternal shield and for the peritrematal shield region also follow modifications and additions by Johnston and Moraza (1991).

## Taxonomy

### Genus *Antennoseius* Berlese, 1916

Type species: *Antennoseius delicatus* Berlese, 1916, by original designation. *Diagnosis*: The diagnosis of *Antennoseius* used here is based on that of Moraes *et al.* (2016).


### *Antennoseius (Vitzthumia) bregetovae* Chelebiev, 1984

*Antennoseius (Vitzthumia) bregetovae* Chelebiev, 1984: 1629.


*Antennoseius (Vitzthumia) bregetovae* Lindquist and Walter, 1989: 1293; Beaulieu *et al.*, 2008: 54; Lindquist and Moraza, 2009: 33; Moraza and Kazemi, 2009: 64; Moraes *et al.*, 2016: 67.

(Figures 1-2)

**Specimens examined** — Type material: slide number 1443 with one adult female (paratype), Saran, Karaganda province, Kazakhstan, 14.4.1973, K.A. Chelebiev coll., in nest of *Lagurus lagurus* (Pallas) (Rodentia: Cricetidae); deposited in the ZIN, Saint Petersburg. Other



**Figure 1** *Antennoseius (Vitzthumia) bregetovae* Chelebiev, 1984, adult female: A – Idiosoma, dorsal view, enlarged sections not to scale; B – Idiosoma, ventral view; C – Subcapitulum; D – Epistome; E – Chelicera; F – Leg I (trochanter-tibia).



**Figure 2** DIC micrographs of *Antennoseius (Vitzthumia) bregetovae* Chelebiev, 1984, adult female: A – Idiosoma, dorsal view; B – Idiosoma, ventral view.

material: two adult females, Aktash, Ulagansky District, Altai Republic, Russia, 50°14'28.0"N 87°41'57.0"E, 30 July 2020, O. Joharchi coll., on *Harpalus* sp. (Coleoptera: Carabidae).

**Redescription — Female** — two specimens measured.

Dorsal idiosoma (Figures 1A & 2A) — Idiosoma oval (510–515 long × 305–328 wide); dorsal shield divided and lineate-reticulate over entire surface (Figures 1A & 2A). Podonotal shield (279–284 long × 262–275 wide) with 19 pairs of setae (*j1*–6, *z1*–6, *s1*–6, *r2*), *r3*–6 off shield, of which four pairs (*j3*–5 and *z4*) short (16–18) strongly thickened, spinelike, with a short thin tip (Figures 1A & 2A) and nine pairs of pore-like structures, including four pairs of gland openings (*gd1*, *gd2*, *gd4*, *gd5*) and five pairs of poroids (*id1*–2, *id4*–6). Setae *j1* (11–13), *j2* (14–16), *z1* (7–9) smooth and short, *j1* slightly spinelike, other podonotal shield setae sub-equal (20–28) and mostly smooth, occasionally with 1–2 small barbs in *s* and *r* series (Figures 1A & 2A). Opisthonotal shield (229–232 long × 262–275 wide), evenly rounded posteriorly, with 15 pairs of setae (19–25) (*J1*–5, *Z1*–5, *S1*–5), setae in *Z* and *S* series with 1–2 small barbs, *Z5* (36–38) longest, and 13 pairs of pore-like structures, including three pairs of gland openings (*gd6*, *gd8*, *gd9*) and ten pairs of poroids (*idm1*–*idm6*, *is1*, *idl1*, *idl3*, *idx*)

(Figures 1A & 2A); transverse interval between setae *J5* about two times shorter than those between *J4* and *Z5*. Lateral soft cuticle with one pair descramble poroid (*idr6*) and 21–22 pairs of setae, mostly sparsely barbed, including setae *r3-r6* (18–21) and 17–18 pairs of setae *R* and *UR* (12–19). Muscle-marks (sigillae) visible mostly on podonotal shield (Figures 1A & 2A).

Ventral idiosoma (Figures 1B & 2B) — Tritosternum with paired sparsely pilose laciniae (90–94), fused basally (13–15), columnar base 20–22 × 13–16 wide; presternal area with pair of narrow indistinct platelets merged to anterior margin of sternal shield, sternal shield length 121–126, width 88–92, fused to endopodal elements between coxae I and II, with a median notch in its anterior margin, posterior margin more or less straight, lineate on lateral margins, with three pairs of smooth subequal setae *st1–3* (28–32), and three pairs of poroids (Figures 1B & 2B). Metasternal setae *st4* (29–31) smooth, inserted on soft cuticle. Endopodal platelets between coxae II–III and III–IV present, small and free. Genital shield drop-shaped length 127–133, width 55–57, faintly lineated laterally, anterior margin of shield convex, not overlapping posterior margin of sternal shield, posterior margin evenly rounded, setae *st5* (23–25) on lateral margins of shield, paragenital poroids *iv5* located on soft cuticle laterad to shield near setae *st5* (Figures 1B & 2B). Anal shield obtriangular, length 100–103, width 75–77, with lineate-reticulate over entire surface, paraanal setae (18–20) shorter than post-anal seta (28–30), cribrum goatee-shaped, consisting of a terminal tuft with 3–4 irregular rows of spicules, restricted to area behind postanal seta (Figures 1B & 2B); anal gland pores (*gv3*) on anterolateral margin of anal shield, posterior to paranal setae, a pair of postanal sigillae present. Soft opisthogastric cuticle with pair of sub-oval metapodal plates (26–28 long × 9–11 wide), nine pairs of smooth setae (*Jv1–5, Zv1–4*) (15–26), three pairs of poroids (two *ivo*; *ivp*) and a gland pores *gv2*, laterad to coxae IV. Peritrematal shield fused with exopodal elements along coxae II to IV, anteriorly fused to podonotal shield, with longitudinal lines, evenly rounded posteriorly behind coxae IV, with two pairs of glandular pores and two pairs of poroids; gland pore *gp1* located on shield's ventral edge near abutment of endopodal extension of sternal shield between coxae I and II, poroid *ip1* and adjacent gland pore *gp2* inserted at level between coxae II and III, and poroid *ip2* next to posterior edge of stigma (Figures 1B & 2B). Peritremes long, extending from stigmata at mid-level of coxae IV to anterior margin of coxae I (Figures 1B & 2B).

Gnathosoma (Figures 1C, 1D & 1E) — Anterior margin of epistome convex, irregularly denticulate, mid surface of epistome with transverse line of denticles, slightly concave medially (Figure 1D). Hypostomal groove with seven transverse rows of denticles, with smooth anterior and posterior transverse lines, all connected laterally, rows 1–5 similar in width, each bearing one or two median denticles, 6<sup>th</sup> and 7<sup>th</sup> rows slightly widened and with four denticles (Figure 1C). Hypostome with four pairs of smooth setae, *h1* (31–33) > *h3* (25–27) > *pc* (18–20) > *h2* (16–18). Corniculi robust and horn-like, slightly shorter than smooth internal malae. Supralabral process not distinguishable. Labrum blade-like, slightly shorter than internal malae. All palp setae smooth and setiform, palptarsal claw two-tined. Fixed digit of chelicera with an offset distal tooth (gabelzhan), followed by seven small sized teeth, a minute setiform pilus dentilis, and a hyaline flap at basal area, dorsal cheliceral seta thick, prostrate, movable digit with two large teeth medially and a minute tooth subapically, with a ventral projection, arthrodial membrane with a rounded flap, cheliceral lyrifissures distinct (Figure 1E).

Legs — Legs II (328–332) and III (335–342) short, legs I (430–438) and IV (418–423) longer.

Legs I to IV with chaetotactic formulae of femora, genua, tibiae as described for genus. Coxae II–IV lineate on posterior inner surfaces (Figure 1B); both setae of coxa I and posterior seta of coxa II modified as spines, pointed abruptly at tip, *av* on coxa II and setae of coxae III–IV normal, slender (Figure 1B). Leg I with setae *pd* of trochanter and *pd2* of femur stout, pointed, spinelike, and with *pd3* of genu stout, spinelike, blunt (Figure 1F); *pd1* on femur slightly thickened, other setae normal, slender (Figure 1F). Legs II–IV without modified setae and smooth. Tibia III having eight setae (2–1/1–2/1–1).

Remarks — *Antennoseius (V.) bregetovae* was described from Kazakhstan (Chelebiev 1984). It has been found in nest of steppe vole, *Lagurus lagurus* (Pallas) (Rodentia: Cricetidae). The description of this species is brief and both the description and illustrations lack many important details. Since that time, there has been no other formal occurrence or recording of this species in the world, and is now recorded in Russia for the first time, on ground beetle *Harpalus* sp. (Coleoptera: Carabidae). Lindquist and Moraza (2009) suspected this species to be a senior synonym of *Antennoseius (Vitzthumia) koroljevae* Chelebiev due to both smooth-morph (phoretic) and granular-morph (free-living) occurred in the same locality and habitat, were discussed by Lindquist and Walter (1989). But according to Chelebiev (1984) and our observations on labels of type series of both species, these two species have been collected from two different nest of Rodents in two different regions and habitats (see provided information in specimens examined section for both species in the current study). In the meantime, *A. (V.) koroljevae* was collected about two years after *A. (V.) bregetovae*. Therefore, we believe these are two distinct species. The diagnosis given above is based primarily on specimens from Russia, but also in comparison with the type material. The species is easily recognized by the podonotal shield with four pairs of spur-like setae (*j3–5* and *z4*), both setae of coxa I and posterior seta of coxa II modified as spines (pointed abruptly at tip), and leg I with setae *pd* of trochanter and *pd2* of femur stout, pointed, spinelike, and with *pd3* of genu stout, spinelike, blunt.

### ***Antennoseius (Antennoseius) bullitus* Karg, 1969**

*Antennoseius bullitus* Karg, 1969: 398.


*Antennoseius bullitus* Karg, 1971: 301; 1977: 5; 1993: 306; Gwiazdowicz, 2007: 55; Kontschán, 2007: 101; Belova and Makarova, 2017: 978.

*Antennoseius (Antennoseius) bullitus* Bregetova, 1977: 251; Moraes *et al.*, 2016: 67; Faraji *et al.*, 2017: 338.

(Figure 3)

**Specimens examined** — Four females, vicinity of lake Kuchak, Tyumen Province, RUSSIA, 57°21'N, 66°03'E, 23 September 2021, O. Joharchi coll., on *Harpalus* sp. (Coleoptera: Carabidae).

Remarks — *Antennoseius (A.) bullitus* was described from Vogtland, Germany (Karg 1969) where it was found on grass. The species has been recorded as phoretic on different species of ground beetles in Europe (Hungary, Ukraine, Poland, Romania; see Faraji *et al.* 2017). Belova and Makarova (2017) collected the species on seven different carabid species from Southern Taiga Forest of Vologda, European Russia. Evidently, this species may be widely distributed in Central and Eastern Europe where associated with a diversity of ground beetles. It is now recorded in North Asia for the first time, on *Harpalus* sp. (Coleoptera: Carabidae). The species is easily recognized by podonotal shield with setiform setae, except seta *j1* distally expanded (Figure 3A), sternal shield with a mid-longitudinal delineation (Figure 3B), coxal setae on legs I–IV slender and setiform (Figure 3B), genital shield drop-shaped, with lateral margins constricted anterior to genital setae and posterior margin more or less bluntly pointed (Figure 3B), anal shield with only circumanal setae (Figure 3B), hypostomal groove with seven transverse rows of denticles, rows 1–5 each with one or two median denticles, 6<sup>th</sup> and 7<sup>th</sup> rows bearing four and seven denticles, respectively (Figure 3C), anterior margin of epistome convex, sparsely denticulate, mid surface of epistome with transverse line of denticles, slightly concave medially (Figure 3D), fixed digit of chelicera with an offset distal tooth (gabelzhan), followed by 12–13 small sized teeth, movable digit with two large teeth medially and a minute tooth subapically (Figure 3E).



**Figure 3** DIC micrographs of *Antennoseius (Antennoseius) bullitus* Karg, 1969, adult female: A – Idiosoma, dorsal view, enlarged sections not to scale; B – Idiosoma, ventral view; C – Subcapitulum; D – Epistome; E – Chelicera.

**Antennoseius (Vitzthumia) hyperboreus Nikolsky, 1988**

*Antennoseius (Vitzthumia) hyperboreus* Nikolsky, 1988: 30.

*Antennoseius (Vitzthumia) hyperboreus* Beaulieu *et al.*, 2008: 55; Moraza and Kazemi, 2009: 65; Moraes *et al.*, 2016: 69.

(Figures 4-6)


**Specimens examined** — Type material: slide number 75 with two adult females (holotype and paratype), Gydanskiy Peninsula, vicinity of Nosok village, Krasnoyarskii Krai, tundra, 24.07.1977, B.S. Yudin coll. on *Microtus oeconomus* (Pallas) (Rodentia: Cricetidae); deposited in ISEA, Novosibirsk.

**Redescription — Female** — Holotype and one paratype measured.


Dorsal idiosoma (Figures 4A, 5A & 5C) — Idiosoma oval (595–599 long × 341–345 wide); dorsal shield divided and harshly lineate-reticulate over entire surface (Figures 4A & 5A). Podonotal shield (320–324 long × 323–326 wide) with 20 pairs of setae (*j1*–6, *z1*–6, *s1*–6, *r2*, and one extra set of paired setae *x* between *s3*–*s4*), setae *r2* and *x* off shield on the left side (in holotype), and nine pairs of pore-like structures, including four pairs of gland openings (*gd1*, *gd2*, *gd4*, *gd5*) and five pairs of poroids (*id1*–2, *id4*–6) (Figures 4A & 5A). Setae *j1* (32–34) longest, slightly thicker, sparsely barbed, *z1* (16–18) shortest, other podonotal shield setae sub-equal (27–35) and smooth (Figures 4A, 5A & 5C). Opisthonotal shield (273–275 long × 302–306 wide), evenly rounded posteriorly, with 15 pairs of setae (27–35) (*J1*–5, *Z1*–5, *S1*–5), *R1* on the right side of anterolateral corner of the shield in holotype (Figures 4A & 5A), setae *Z5* (40–42) longest, slightly thicker, sparsely barbed, and 13 pairs of pore-like structures, including three pairs of gland openings (*gd6*, *gd8*, *gd9*) and ten pairs of poroids (*idm1*–*idm6*, *is1*, *idl1*, *idl3*, *idx*) (Figures 4A & 5A). Lateral soft cuticle with 35–36 pairs of setae (including ventral setae), including *r3*–6 and 31–32 pairs of hypertrichous series *R* and *UR*, and two pairs of poroids. Muscle-marks (sigillae) visible mostly on podonotal shield (Figures 4A & 5A).

Ventral idiosoma (Figures 4B, 5B, 5D–E) — Tritosternum with paired pilose laciniae (100–103), fused basally (20–22), columnar base 24–26 × 20–23 wide; presternal area with a pair of narrow and faint platelets, sternal shield length 134–137, width 117–121, fused to endopodal elements between coxae I and II, with a median notch in its anterior margin, posterior margin irregularly straight, slightly lineate on lateral margins, region anterior to *iv1* poorly sclerotized, with three pairs of smooth setae (*st1* 35–37, *st2* 33–35, *st3* 29–31) (*st3* off shield on right side in holotype), and three pairs of poroids (*iv3* absent on left side in holotype) (Figures 4B, 5B & 5E). Metasternal setae *st4* (28–30) smooth, inserted on soft cuticle. Endopodal platelets between coxae II–III and III–IV present, small and free. Genital shield drop-shaped length 174–176, width 73–76, faintly lineated anteriorly, anterior margin of shield convex, not overlapping posterior margin of sternal shield, posterior margin evenly rounded, setae *st5* (27–29) on lateral margins of shield, paragenital poroids *iv5* located on soft cuticle laterad to shield near setae *st5* (Figures 4B, 5B & 5D). Anal shield subtriangular, length 113–116, width 92–95, with lineate-reticulate over entire surface, para-anal setae and post-anal seta subequal in length (20–22), cribrum large, goatee-shaped, restricted to area behind postanal seta (Figures 4B, 5B & 5D); anal gland pores (*gv3*) on anterolateral margin of anal shield, posterior to paranal setae, a pair of postanal sigillae present. Soft opisthogastric cuticle with pair of sub-oval metapodal plates (20–22 long × 9–12 wide), ten pairs of setae, *Jv1*–3 and *Zv1*–4 smooth, *Jv4*–5, and *Zv5* slightly thicker, with 1–2 small barbs, and four pairs of poroids (three *ivo*; *ivp*) and a gland pores *gv2*, laterad to coxae IV. Peritrematal shield fused with exopodal elements along coxae II to IV, anteriorly fused to podonotal shield, with longitudinal lines, evenly rounded posteriorly behind coxae IV, with one pair of glandular pore and two pairs of poroids; poroid *ip1* and adjacent gland pore *gp2* inserted at level between coxae II and III, and poroid *ip2* next to posterior edge of stigma (Figures 4B & 5B). Peritremes long, extending from stigmata at mid-level of coxae IV to anterior margin of coxae I (Figures 4B & 5B).


Gnathosoma (Figures 4C–D & 5F–G) — Anterior margin of epistome convex, irregularly denticulate, mid surface of epistome with transverse line denticles, concave medially (Figure



**Figure 4** *Antennoseius (Vitzthumia) hyperboreus* Nikolsky, 1988 (holotype), adult female: A – Idiosoma, dorsal view, enlarged sections not to scale; B – Idiosoma, ventral view; C – Epistome; D – Chelicera.



**Figure 5** DIC micrographs of *Antennoseius (Vitzthumia) hyperboreus* Nikolsky, 1988 (holotype), adult female: A – Idiosoma, dorsal view; B – Idiosoma, ventral view; C – General view of some podonotal setae; D – General view of opisthogastric area; E – Sternal shield; F – Subcapitulum ; G – Epistome.



**Figure 6** *Antennoseius (Vitzthumia) hyperboreus* Nikolsky, 1988 (holotype), adult female: A – Leg I (trochanter-tibia); B – Leg II; C – Leg III; D – Leg IV.

4C & 5G). Hypostomal groove with seven transverse rows of denticles, each row with 4–9 small denticles, with smooth anterior and posterior transverse lines, all connected laterally, 6<sup>th</sup> and 7<sup>th</sup> rows slightly convex and concave, respectively (Figure 5F). Hypostome with four pairs of smooth setae, *h1* (44–46) > *pc* (33–35) > *h2* ≈ *h3* (27–30). Corniculi robust and horn-like, slightly shorter than smooth internal malae. Supralabral process not distinguishable. Labrum blade-like, slightly shorter than internal malae. All palp setae smooth and needle-like except *d1* on palp femur with some barbs, palp-tarsal claw two-tined. Fixed digit of chelicera with an offset distal tooth (gabelzhan), followed by 19–20 small sized teeth, a minute setiform pilus dentilis, dorsal cheliceral seta thickened, thorn-like, movable digit with two large teeth medially, with a ventral projection, arthrodial membrane with a rounded flap, cheliceral lyrifissures distinct (Figure 4D).

Legs (Figure 6) — Legs II (506–512) and III (480–486) short, legs I (694–700) and IV (606–612) longer. Legs I to IV with chaetotactic formulae of femora, genua, tibiae as described for genus (Figure 6). Coxae II–IV lineate on posterior inner surfaces and setae of coxae I–IV normal, slender (Figure 4B). Legs I–IV setae unmodified, some setae sparsely barbed (more distinct on legs I and IV, Figure 6), not elongated as macrosetae, except four subapical setae on tarsus I elongated (ratio of subapical setae / tarsus I length ≈ 0.65). Tibia III having nine setae, *pl2* present. Complement of setae on segments of legs I-II-III-IV: coxae 2–2–2–1; trochanters 6–5–5–5; femora 2–3/1–2/2–2; 2–3/1–2/2–1; 1–2/1–1/0–1; 1–2/0–1/1–1; genua 2–3/1–3/2–2; 2–3/1–2/1–2; 2–2/1–2/1–1; 2–2/0–3/1–1; tibiae 2–3/1–3/2–2; 2–2/1–2/1–2; 2–1/1–2/1–2; 2–1/1–3/1–2 and tarsi II-IV 3–3/2–1/1–3/2–3.

**Remarks** — *Antennoseius (V.) hyperboreus* was described from tundra in Northern Russia (Nikolsky 1988). It has been found on tundra vole, *Microtus oeconomus* (Pallas) (Rodentia: Cricetidae). Since that time, there has been no other formal occurrence or recording of *A. (V.) hyperboreus* in the world. The original description of the species is brief and its illustrations are incomplete, lacking many important details. We herein redescribe *A. (V.) hyperboreus* on the basis of type series to complement the original description which provided by Nikolsky (1988). The species is easily recognized by the podonotal shield with 20 pairs of setae, none of the setae enlarged (in distinction from others), setae *j1* and *Z5* sparsely barbed and slightly thicker than other setae, sternal shield with strong anterior medial notch (reaching to level of *iv1*), lateral soft cuticle with 35–36 pairs of setae (including ventral setae), coxal setae on legs I–IV slender and setiform, four long subapical setae on tarsus I (ratio of subapical setae / tarsus I length ≈ 0.65).

#### ***Antennoseius (Vitzthumia) koroljevae Chelebiev, 1984***

*Antennoseius (Vitzthumia) koroljevae* Chelebiev, 1984: 1631.

*Antennoseius (Vitzthumia) koroljevae* Lindquist and Walter, 1989: 1293; Beaulieu *et al.*, 2008: 55; Moraza and Kazemi, 2009: 63; Moraes *et al.*, 2016: 70; Joharchi *et al.*, 2020: 477.


*Antennoseius (Vitzthumia) koreljevae* (*sic*) Lindquist and Moraza, 2009: 3

(Figure 7)

**Specimens examined** — Type material: slide number 1792 with one adult female (paratype), vicinity of Nura River, Tokarevka, Karaganda province, Kazakhstan, 07.06.1975, K.A. Chelebiev coll., in nest of *Microtus gregalis* (Pallas) (Rodentia: Cricetidae); deposited in the ZIN, Saint Petersburg.

**Remarks** — *Antennoseius (Vitzthumia) koroljevae* is known only from its original description, being five females collected in nest of narrow-headed vole, *Microtus gregalis* (Pallas) (Rodentia: Cricetidae), in Kazakhstan (Chelebiev 1984). The description of the species is brief and both the description and illustrations lack many important details. We herein provide additional morphological characters to complement the previous description (Chelebiev 1984) of *A. (V.) koroljevae* on the basis of type series:

Dorsal shields densely granulate-tuberculate, lacking reticulate patterning (Figure 7A), podonotal shield with 21 pairs of setae including three pairs of *sx* setae and four unpaired seta



**Figure 7** DIC micrographs of *Antennoseius (Vitzthumia) koroljevae* Chelebiev, 1984 (holotype), adult female: A – Idiosoma, dorsal view; B – Idiosoma, ventral view; C – Sternal shield; D – General view of opisthogastric area; E – Epistome; F – Subcapitulum.

*x* were present between setae *j3–j5* (Figure 7A), setae *r2, r3, r4* off the shield (Figure 7A), opisthonotal shield with 17 pairs of setae including two pairs of *Sx* setae and three unpaired seta *X* were present between *Z* and *S* series (Figure 7A), all dorsal setae moderately barbed (Figure 7A), setae *j1* and *Z5* longer and thicker than other setae; presternal area with pair of narrow distinct platelets (Figure 7B & 7C), sternal shield with strong anterior medial notch reaching half-way between *st1* and *st2* (Figure 7B & 7C); genital shield drop-shaped, its lateral margins widened posteriorly, posterior margin evenly rounded (Figure 7B & 7D); anal shield densely granulate-tuberculate, lacking reticulate patterning (Figure 7B & 7D); anterior margin of epistome convex, irregularly denticulate (Figure 7E), hypostomal groove with seven transverse rows of denticles, each row with 10–13 small denticles, with smooth anterior transverse line (Figure 7F); all palp setae smooth and needle-like, palp-tarsal claw two-tined, fixed digit of chelicera multidentate; almost all leg setae moderately barbed, setae *pd* on trochanter and *pd2* on femur of leg I slightly thickened. Tibia III having eight setae (2–1/1–2/1–1).

### ***Antennoseius (Antennoseius) pannonicus* Willmann, 1951**

*Antennoseius pannonicus* Willmann, 1951: 109.

*Antennoseius pannonicus* Athias-Henriot, 1961: 461; Ryke, 1962: 662; Karg, 1971: 298; 1977: 4; 1993: 305.

*Antennoseius (Antennoseius) pannonicus* Bregetova, 1977: 248; Beaulieu *et al.*, 2008: 47; Lindquist and Moraza, 2009: 34; Moraes *et al.*, 2016: 73; Faraji *et al.*, 2017: 339; Bahrami and Kazemi, 2019: 357.

(Figure 8)

**Specimens examined** — Two females, vicinities of Uspenka, Tyumen Province, Russia, 57°04'N, 65°04'E, 18 October 2021, Vladimir A. Khaustov coll., on *Harpalus* sp. (Coleoptera: Carabidae).

**Remarks** — *Antennoseius (A.) pannonicus* was described from Austria (Willmann 1951) where it was found in meadow soil. The species has since been recorded as phoretic on different species of ground beetles in Europe (see Faraji *et al.* 2017) and Iran (see Bahrami and Kazemi, 2019). It is now recorded in Russia for the first time, on *Harpalus* sp. (Coleoptera: Carabidae).

Faraji *et al.* (2017) have redescribed this species based on newly collected specimens from the France. Our specimens agree very well with both the original description and redescription of the species given by Willmann (1951) and Faraji *et al.* (2017), respectively. This species can be readily recognized by the presence of four pairs of spur-shaped setae on podonotal shield (including *j5*) (Figure 8A), sternal shield with a prominent brownish crown-shape configuration between *st1* setae (Figure 8B), setae *st3* off sternal shield on soft cuticle (Figure 8B) and coxal setae on legs I-IV setiform (Figure 8B).

### ***Antennoseius (Antennoseius) ponticus* Trach and Makarova, 2008**


*Antennoseius ponticus* Trach and Makarova, 2008: 181.

*Antennoseius (Antennoseius) ponticus* Lindquist and Moraza, 2009: 33; Moraes *et al.*, 2016: 73; Faraji *et al.*, 2017: 338.

(Figure 9)


**Specimens examined** — Three females, near the Moinakskoye Lake, Yevpatoria, Western Crimea, Russia, 45°11'31.0"N 33°19'59.0"E, 13 June 2021, O. Joharchi coll., on *Harpalus* sp. (Coleoptera: Carabidae).

**Remarks** — *Antennoseius (A.) ponticus* was originally described from southern Ukraine (Odessa and Mykolaiv Regions) where it was found on carabids (*Harpalus seripes* (Quensel), *Amara* sp.) and leaf beetle, *Chrysolina gypsophilae* (Küster) (Coleoptera: Chrysomelidae) (Trach and Makarova 2008), and afterward has been recorded from Eastern Ukraine (Trach 2013). It is now recorded in Russia for the first time, on *Harpalus* sp. (Coleoptera: Carabidae). Our specimens agree very well with the description given by Trach and Makarova 2008. The



**Figure 8** DIC micrographs of *Antennoseius (Antennoseius) pannonicus* Willmann, 1951, adult female: A – Idiosoma, dorsal view; B – Idiosoma, ventral view.

species may be recognized primarily by the podonotal shield with five pairs of spikelike (with a short thin tip) setae (including z4) (Figure 9A), presternal region with one pair of oblique narrow distinct platelets (Figure 9B), ventral posterior seta on coxae I and II modified as spines, pointed abruptly at tip (Figure 9B), fixed digit of chelicera with six teeth and movable digit with two large teeth medially and a minute tooth subapically (Figure 9D). The illustration and descriptions of the hypostomal groove in Trach and Makarova's (2008) shows six transverse rows of denticles, each row with 6–7 small denticles. Based on the specimens collected in Crimea, we found hypostomal groove with seven transverse rows of denticles, with smooth anterior and posterior transverse lines, rows 1–6 each bearing 2–4 small denticles, 7<sup>th</sup> row with only one median denticle (Figure 9C).



**Figure 9** DIC micrographs of *Antennoseius (Antennoseius) ponticus* Trach and Makarova, 2008, adult female: A – Idiosoma, dorsal view; B – Idiosoma, ventral view; C – Subcapitulum; D – Chelicera.

## Acknowledgements

The present research was supported by the grant from the Russian Science Foundation, project number 20–64–47015. We cordially thank Dr. Alexander Khaustov, Mr. Vladimir A. Khaustov and Mr. Roman Latyntsev (Tyumen State University, Russia) for their help in collecting samples, and in logistics, respectively. We are deeply grateful to Zoological Institute of the Russian Academy of Sciences (ZIN, Saint Petersburg), and Institute of Systematics and ecology of animals (ISEA, Novosibirsk) for their hospitality of the first author during his visit for examining the mite collections.

## References

- Athias-Henriot C. 1961. Mesostigmates (Urop. Excl.) édaphiques méditerranéens (Acaromorpha, Anactinotrichida) (collect. Prof. H. Franz et C. Athias-Henriot). Première Série. *Acarologia*, 3(4): 381–509. [in French].
- Athias-Henriot C. 1971. La divergence néotaxique des Gamasides (Arachnides). *Bulletin Sc Bourgogne*, 28: 93–106. [in French].
- Athias-Henriot C. 1975. Nouvelles notes sur les Amblyseiini. 2. Le relevé organotaxique de la face dorsale adulte (gamasides, protoadéniques, Phytoseiidae). *Acarologia*, 17: 20–29. [in French].
- Bahrami F., Kazemi S. 2019. First record of *Antennoseius pannonicus* Willmann (Mesostigmata: Ascidae) from Iran. *Persian J. Acarol.*, 8(4): 357–359. <https://doi.org/10.22073/pja.v8i4.53561>
- Belova Yu.N., Makarova O.L. 2017. Mites (Acari) phoretic on ground beetles (Coleoptera, Carabidae) in a southern taiga forest in the environs of Vologda. *Entomol. Rev.*, 97 (7): 975–983. <https://doi.org/10.1134/S0013873817070120>
- Berlese A. 1911. Alcuni Acari entomofili nuovi. *Redia*, 7: 183–186.
- Berlese A. 1916. Centuria terza di Acari nuovi. *Redia*, 12: 289–338.
- Beaulieu F., Déchêne A.D., Walter D.E. 2008. Phase morphs and phoresy: new species of *Antennoseius (Vitzthumia)* mites (Acari: Mesostigmata: Ascidae) associated with pyrophilous carabids (Carabidae: *Sericoda* spp.) in Alberta, Canada. *Zootaxa*, 1961: 37–57. <https://doi.org/10.11646/zootaxa.1961.1.4>
- Bregetova N.G. 1977. Family Antennoseiidae Karg, 1965. In: Ghilyarov, M.S., Bregetova, N.G. (Eds). Key to the soil-inhabiting mites, Mesostigmata. Leningrad: Nauka. p. 244–253. [in Russian].
- Chelebiev K.A. 1984. Mites of the genus *Antennoseius* (Parasitiformes: Mesostigmata) from central Kazakhstan. *Zool. Zh.*, 63, 1629–1633. [in Russian].
- Costa M. 1969. *Antennoseius bytinskii* sp. nov., with notes on the genus *Antennoseius* Berlese (Acari: Mesostigmata) in Israel. *Israel J. Entomol.*, 4: 217–226.
- Eidelberg M.M. 2000. Three new mite species of the family Antennoseiidae (Parasitiformes, Gamasina). *Zool. Zh.*, 79(12): 1396–1401. [in Ukrainian].
- Evans G.O. 1963a. Observations on the chaetotaxy of the legs in the free-living Gamasina (Acari: Mesostigmata). *Bull. Br. Mus. (Nat. Hist.), Zool.*, 10: 277–303. <https://doi.org/10.5962/bhl.part.20528>
- Evans, G.O. 1963b. Some observations on the chaetotaxy of the pedipalps in the Mesostigmata (Acari). *Ann. Mag. Nat. Hist., Ser.* 13, 6: 513–527. <https://doi.org/10.1080/00222936308651393>
- Evans G.O., Till W.M. 1979. Mesostigmatic mites of Britain and Ireland (Chelicera: Acari Parasitiformes). An introduction to their external morphology and classification. *Transactions of the Zoological Society of London*, 35: 145–270. <https://doi.org/10.1111/j.1096-3642.1979.tb00059.x>
- Faraji F., Dehelean S.B., Vuyk S.M., Bakker, F. 2017. Four new species records of *Antennoseius* and *Anystipalpus* (Acari: Mesostigmata: Ascidae) phoretic on Carabidae beetles from France. *Acarologia*, 57(2): 337–353. <https://doi.org/10.1051/acarologia/20174159>
- Felska M., Wohltmann A., Mąkol J. 2018. A synopsis of host-parasite associations between Trombidioidea (Trombidiformes: Prostigmata, Parasitengona) and arthropod hosts. *Syst. Appl. Acarol.*, 23: 1375–1479. <https://doi.org/10.11158/saa.23.7.14>
- Halliday R.B., Walter D.E., Lindquist E.E. 1998. Revision of the Australian Ascidae (Acarina: Mesostigmata). *Invertebr. Taxon.*, 12: 1–54. <https://doi.org/10.1071/IT96029>
- Hunter P.E., Rosario R.M.T. 1988. Associations of Mesostigmata with other arthropods. *Annu. Rev. Entomol.*, 33: 393–417. <https://doi.org/10.1146/annurev.en.33.010188.002141>
- Johnston D.E., Moraza M.L. 1991. The idiosomal adenotaxy and poroidotaxy of Zerconidae (Mesostigmata: Zerconina). In: Dusbábek F., Bukva V. (Eds). *Modern Acarology*. Vol. 2. Academia: Prague. p. 349–356.
- Karg W. 1969. Untersuchungen zur Kenntnis der Ascaloidea Karg, 1965 (Acarina: Parasitiformes) mit der Beschreibung von acht neuen Arten. *Zool. Anzeiger*, 182: 393–406.
- Karg W. 1971. Acari (Acarina), Milben, Unterordnung Anactinochaeta (Parasitiformes): Die freilebenden Gamasina (Gamasides), Raubmilben. In: *Die Tierwelt Deutschlands und der angrenzenden Meeresteile*, 59. Teil. Gustav Fischer Verlag, Jena. 475p.
- Karg W. 1977. Die Milbengattung *Antennoseius* Berlese, 1916 (Acarina, Parasitiformes) mit einer neuen Art aus dem Leutrat bei Jena (DDR). *Abhandlungen und Berichte des Naturkundemuseums Görlitz*, 50(4): 1–7.
- Karg W. 1993. Acari (Acarina), Milben Parasitiformes (Anactinochaeta) Cohors Gamasina Leach, Raubmilben. In: *Die Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und ihrer Lebensweise*. 59. Teil. 2 überarbeitete Auflage, Gustav Fischer Verlag, Jena, 523 pp.

- Kazemi S. 2018. Redescription of *Antennoseius (Antennoseius) avius* Karg (Acari: Mesostigmata: Ascidae), a senior synonym of *A. (A.) alexandrovi* Bregetova and *A. (A.) arvensis* Kalúz, with additional information for *Antennoseius* Berlese, and a key to the Iranian species of the genus. *Internat. J. Acarol.*, 44: 171-179. <https://doi.org/10.1080/01647954.2018.1471101>
- Kazemi S., Moraza M.L. 2013. Mites of the genus *Antennoseius* Berlese (Acari: Mesostigmata: Ascidae) from Iran. *Persian J. Acarol.*, 2(2): 217-234.
- Kontschán J., 2007. New and rare mesostigmatid mites to the fauna of Hungary. *Folia Historico Naturalia Musei Matraensis*, 31: 99-106.
- Lindquist E.E., Evans G.O. 1965. Taxonomic concepts in the Ascidae, with a modified setal nomenclature for the idiosoma of the Gamasina (Acarina: Mesostigmata). *Mem. Entomol. Soc. Can.* 47: 1-64. <https://doi.org/10.4039/entm9747fv>
- Lindquist E.E., Moraza M.L. 2009. *Anystipalpus*, *Antennoseius* and *Vitzthumia*: a taxonomic and nomenclatural conundrum of genera (Acari: Mesostigmata: Dermanyssina), with description of four species of *Anystipalpus*. *Zootaxa*, 2243: 1-39. <https://doi.org/10.11646/zootaxa.2243.1.1>
- Lindquist E.E., Walter D.E. 1989. *Antennoseius (Vitzthumia) janus* n.sp. (Acari: Ascidae), a mesostigmatic mite exhibiting adult female dimorphism. *Can. J. Zool.*, 67: 1291-1310. <https://doi.org/10.1139/z89-184>
- Lindquist E.E., Krantz G.W., Walter D.E. 2009. Order Mesostigmata. In: Krantz G.W., Walter, D.E. (Eds). *A Manual of Acarology*. 3<sup>rd</sup> Edition. Texas: Texas Tech University Press, Lubbock. p. 1-807.
- Moraes G.J., Britto E.P.J., Mineiro J.L. de C., Halliday B. 2016. Catalogue of the mite families Ascidae Voigts & Oudemans, Blattisociidae Garman and Melicharidae Hirschmann (Acari: Mesostigmata). *Zootaxa*, 4112 (1): 1-299. <https://doi.org/10.11646/zootaxa.4112.1.1>
- Moraza M.L., Kazemi S. 2009. A new species of *Antennoseius (Vitzthumia)* Thor (Acari: Mesostigmata, Ascidae), associated with carabid beetles in Iran and a key to species. *Internat. J. Acarol.*, 35(1): 59-65. <https://doi.org/10.1080/01647950902884538>
- Nikolsky V.V. 1988. New species of gamasid mite of the genus *Antennoseius* Berlese, 1916 (Parasitiformes, Ctamasina (sic)) from Siberia. In: Zolotarenko G.S. (Eds). *Taksonomiya Zhivotnykh Sibiri. Novie i Maloizvestnye Vidy Fauny Sibiri*. Akademiya Nauk SSSR: Sibirskoe Otdelenie. Biologicheskiy Institut. 20: p. 30-32. [in Russian].
- Olynyk J.E., Freitag R. 1979. Some phoretic associations of ground beetles (Coleoptera: Carabidae) and mites (Acarina). *Can. Entomol.*, 111: 333-335. <https://doi.org/10.4039/Ent111333-3>
- Ryke P.A.J. 1962. The genus *Antennoseius* Berlese (Acarina: Rhodacaridae). *Ann. Mag. Nat. Hist.*, 13(4): 657-663. <https://doi.org/10.1080/00222936108651190>
- Trach V.A., Makarova O.L. 2008. A new gamasid mite species of the genus *Antennoseius* (Parasitiformes, Ascidae) from the southwest of Ukraine. *Vest. Zool.*, 42(2): 181-184.
- Trach V.A. 2012. A new species of mites of the genus *Anystipalpus* (Mesostigmata, Ascidae) from the Eastern Ukraine. *Vest. Zool.*, 46(1): 73-77. <https://doi.org/10.2478/v10058-012-0005-1>
- Trach V.A. 2013. On the fauna of gamasid mites of the genera *Anystipalpus* and *Antennoseius* (Mesostigmata, Ascidae) of the Eastern Ukraine. *Vest. Zool.*, 47(5): 387-393. <https://doi.org/10.2478/vzoo-2013-0037>
- Walter D.E., Krantz G.W. 2009. Collecting, rearing and preparing specimens. In: Krantz G.W., Walter D.E. (Eds). *A Manual of Acarology*. Third Edition, Texas Tech University Press, Lubbock, Texas, p. 83-95.
- Willmann C. 1951. Untersuchungen über die terrestrische Milbenfauna im pannonicischen Klimagebiet Österreichs. *Sitzungsberichte der Österreichischen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Abteilung I*, 160: 91-176.